

On Haglund’s Conjecture 4 for the Riemann Ξ approximants

Kunal Tyagi*

October 3, 2026

Abstract

Let Ξ_k be the k -th partial sum of Riemann’s expansion $\Xi = \sum_{n \geq 1} \Phi_n$ of the Ξ -function in incomplete gamma functions. Haglund conjectured (his Conjecture 4) that for every $k \geq 1$ the imaginary part of each non-real zero of $\Xi_k + t\Phi_{k+1}$ decreases as t goes from 0 to 1. We prove that this pencil is Ξ minus a level that is positive on the real axis and decreases with t ; that real zeros stay real if and only if every real critical point of Ξ_{k+1}/Φ_{k+1} with value in $(0, 1]$ is a non-degenerate maximum; and that a simple zero β of Ξ above the real axis with $\text{Im } \Xi'(\beta) > 0$ would give, for every large k , a zero of the pencil whose imaginary part strictly increases. If the zeros of Ξ are real and simple and the level is frozen at a constant, every non-real zero descends and lands at the top of its own lobe; this is known in substance. In floating point, not interval arithmetic, and with two independent programs for $k \leq 3$, the imaginary part decreased at every step of every branch followed, in complete windows for $k \leq 8$ and around the landings for $k = 15, 20, 26, 27, 35, 50$; the real-axis criterion holds on $[0, 4(k+2)^2 + 40]$ for $k \leq 50$. A heuristic in three regimes explains why the true, slowly varying level leaves the frozen picture intact.

1 The conjecture and what is shown here

1.1 The pencil

Write $s = \frac{1}{2} + iz$ and let $\Xi(z) = \frac{1}{2}s(s-1)\pi^{-s/2}\Gamma(s/2)\zeta(s)$, which is even, entire and real on the real axis. Riemann’s series $\Xi = \sum_{n \geq 1} \Phi_n$ and Haglund’s approximants $\Xi_N := \sum_{n \leq N} \Phi_n$ are defined in [Hag09, (1)–(14), pp. 1–3] (page and equation numbers refer to the arXiv version); see also [Tya26, (3)]. From Haglund’s definition (his (6) with $z \in \mathbf{C}$, and his (14)), and on the real axis also from [Tya26, Theorem 3.3 and its proof],

$$\Phi_n(z) = 2 \int_0^\infty \tilde{\varphi}_n(v) \cos(zv) dv \quad (z \in \mathbf{C}), \quad \tilde{\varphi}_n(v) = 2y(2y-3)e^{v/2-y}, \quad y = \pi n^2 e^{2v}, \quad (1)$$

and for $n \geq 2$ the kernel $\tilde{\varphi}_n$ is positive, strictly decreasing and strictly convex on $[0, \infty)$ [Tya26, proof of Theorem 3.3]; for $n = 1$ it is positive but not monotone at 0. Put $Q_N := \sum_{n > N} \Phi_n = \Xi - \Xi_N$, an entire function (the series converges locally uniformly [Hag09, (12)]). For $k \geq 1$ the pencil is

$$F_t := \Xi_k + t\Phi_{k+1}, \quad 0 \leq t \leq 1, \quad u := 1 - t,$$

so that $F_0 = \Xi_k$ and $F_1 = \Xi_{k+1}$. Haglund writes [Hag09, p. 11]: “Another interesting phenomena occurs when we consider the zeros of $\Xi_k(z) + t\Phi_{k+1}(z)$. As we let t vary continuously from 0 to 1,

*hello.jay.tyagi@gmail.com. *Use of AI.* The mathematics in this paper — the derivations, the computations, the verification suite, and the text — was produced by Claude (Anthropic) working under the author’s direction; the author set the objectives, made the technical and editorial decisions, and is responsible for the content. Claude is a tool and is not an author. The complete working record is public: <https://github.com/x67ai/riemann-rh-program>.

computations indicate that the imaginary parts of the non-real zeros decrease monotonically (i.e. continuously), in a very regular manner, until, for high enough k , they collide with the corresponding zero (from Schwartz reflection) in the fourth quadrant, and arrive on the real line, where they remain.” He then states:

Conjecture 4 (Haglund [Hag09, p. 11]). *“For $k \geq 1$, the imaginary part of each non-real zero of $\Xi_k(z) + t\Phi_{k+1}(z)$ decreases monotonically (i.e. continuously) as t goes from 0 to 1.”*

The journal version [Hag11, p. 310] states it as Conjecture 5.1 in the same words, and the author’s later copy [Hag11w, p. 11] leaves the conjecture and its lead-in unchanged. We use two statements about the pencil for a given k :

- (D) along every branch of zeros of F_t in the open upper half-plane, $\text{Im } z$ does not increase with t ;
- (R) a real zero of F_t stays real as t increases.

Statement (D) is the conjecture; (R) is the last clause of the lead-in. A conjugate pair that leaves the real axis at some t_0 violates (R), and also (D) just after t_0 , since the imaginary part of its upper member rises from 0.

1.2 What is shown here

Each item is labeled by its kind. *Proved:* the pencil is Ξ minus a level, $F_t = \Xi - L_t$ with $L_t = u\Phi_{k+1} + Q_{k+1} > 0$ on the real axis and $\partial_t F_t = \Phi_{k+1} > 0$ there (Lemma 2.2); its real zeros lie in the positive lobes of Ξ , an even number in each and an odd number in the central one, and (R) holds for k if and only if every critical point of $S_k := \Xi_{k+1}/\Phi_{k+1}$ on the real axis with value in $(0, 1]$ is a non-degenerate maximum (Theorem 2.4); a simple zero β of Ξ with $\text{Im } \beta > 0$ and $\text{Im } \Xi'(\beta) > 0$ would give, for every large k , a non-real zero of the pencil whose imaginary part strictly increases on the whole interval $0 \leq t \leq 1$ (Proposition 4.2); and along the pencil the coefficient of $1/x^2$ on the real axis is negative and decreases in size (Lemma 6.1). *Proved under the hypothesis that every zero of Ξ is real:* a pair can leave the real axis only where the level satisfies an explicit inequality (Proposition 2.5); if the zeros are also simple and the level is frozen at a constant that decreases, every non-real zero descends along its own curve and lands at the top of its own lobe (Proposition 3.2, known in substance [CS00]). *Numerical, floating point, not interval arithmetic:* a census of the branches by two independent programs, and a test of the real-axis criterion (Section 5). *Heuristic:* why the true, slowly varying level does not change the frozen picture, in three regimes (Section 6). Neither (D) nor (R) is proved here for any k ; Baccaro’s preprint [Bac26] treats $k = 1$ (Section 7).

2 The level form and the real axis

Lemma 2.1. *For $n \geq 2$ and real x , $\Phi_n(x) > 0$.*

Proof. This is the Pólya argument of [Tya26, Theorem 3.1] applied to the single kernel $\tilde{\varphi}_n$. For $x \neq 0$, $\int_0^\infty \tilde{\varphi}_n(v) \cos(xv) dv = x^{-1} \int_0^\infty (-\tilde{\varphi}'_n(v)) \sin(xv) dv$ with $-\tilde{\varphi}'_n$ positive and strictly decreasing, so the pieces of the sine integral over consecutive half-periods alternate in sign with strictly decreasing sizes, the first positive. At $x = 0$ the integral is that of a positive function. $\square$

Lemma 2.2 (the pencil is Ξ minus a positive level). *For $k \geq 1$ and all z ,*

$$F_t(z) = \Xi(z) - L_t(z), \quad L_t := u\Phi_{k+1} + Q_{k+1} = 2 \int_0^\infty \kappa_t(v) \cos(zv) dv, \quad \kappa_t := u\tilde{\varphi}_{k+1} + \sum_{n>k+1} \tilde{\varphi}_n.$$

For $0 \leq u \leq 1$ the kernel κ_t is positive, strictly decreasing and strictly convex, so $L_t(x) > 0$ for real x ; and $\partial F_t / \partial t = \Phi_{k+1}$, which is positive on the real axis.

Proof. $\Xi_k = \Xi - Q_k$ and $Q_k = \Phi_{k+1} + Q_{k+1}$. The kernel properties hold termwise, since all indices are at least 2; then the argument of Lemma 2.1 applies to κ_t , and Lemma 2.1 itself gives $\Phi_{k+1} > 0$. $\square$

Lemma 2.3 (the quotient form). *Let $S_k := \Xi_{k+1} / \Phi_{k+1}$, a meromorphic function that is real on the real axis, where $\Phi_{k+1} > 0$. Then $F_t(z) = 0$ if and only if $S_k(z) = u$, at every z with $\Phi_{k+1}(z) \neq 0$; at a common zero of Ξ_{k+1} and Φ_{k+1} the pencil vanishes for every t . Along a branch of simple zeros $z(t)$ with $\Phi_{k+1}(z) \neq 0$, $dz/dt = -\Phi_{k+1}(z) / \partial_z F_t(z) = -1/S'_k(z)$, so $d(\operatorname{Im} z)/dt = \operatorname{Im} S'_k(z) / |S'_k(z)|^2$. Hence (D) for k is the statement that $\operatorname{Im} S'_k(z) \leq 0$ at every z with $\operatorname{Im} z > 0$ and $S_k(z) \in (0, 1)$, together with the behavior of the branches at multiple zeros.*

Proof. $F_t = \Xi_{k+1} - u\Phi_{k+1}$; differentiate $S_k(z(t)) = 1 - t$. $\square$

Let $0 < \gamma_1 < \gamma_2 < \dots$ be the positive real zeros of Ξ ; a *positive lobe* is an interval (γ_j, γ_{j+1}) on which $\Xi > 0$, and the *central lobe* is $(0, \gamma_1)$.

Theorem 2.4. *Let $k \geq 1$ and $0 \leq t \leq 1$.*

- (a) *For real x , $F_t(x) = \Xi(x) - L_t(x)$ with $L_t(x) > 0$, and $t \mapsto F_t(x)$ is strictly increasing.*
- (b) *Every real zero of F_t lies in an open interval on which $\Xi > 0$. Counted with multiplicity, each positive lobe (γ_j, γ_{j+1}) contains an even number of zeros of F_t and the central lobe an odd number; the number of zeros of F_t in $(0, \infty)$ is finite and odd.*
- (c) *The sets $P_t := \{x \in \mathbf{R} : F_t(x) > 0\}$ increase with t : $P_s \subset P_t$ for $s < t$, and every boundary point of P_s lies in the interior of P_t .*
- (d) *Let x_0 be a real zero of F_{t_0} of multiplicity m , with $F_{t_0}(x) = c(x - x_0)^m(1 + o(1))$. If m is odd, F_t has an odd number of real zeros near x_0 for t near t_0 , counted with multiplicity; for $t \neq t_0$ the m zeros near x_0 are $x_0 + (-(t - t_0)\Phi_{k+1}(x_0)/c)^{1/m}(1 + o(1))$ (the m values of the root), exactly one of them real and $m - 1$ non-real, so for odd $m \geq 3$ a conjugate pair lands at t_0 and leaves the axis at once. The same expansion gives, for even $m \geq 4$, two real and $m - 2$ non-real zeros on the side of t_0 where $F_t(x_0)$ has the sign opposite to c , and m non-real zeros on the other side. If $m = 2$ and $F_{t_0} \leq 0$ near x_0 (a local maximum with value 0), then for t slightly above t_0 there are two simple real zeros near x_0 and for t slightly below there are none: a conjugate pair arrives, a landing. If $m = 2$ and $F_{t_0} \geq 0$ near x_0 (a local minimum with value 0), then for t slightly below t_0 there are two real zeros near x_0 and for t slightly above there are none: a conjugate pair leaves the axis, a lift-off.*
- (e) *Hence (R) holds for k if and only if, for every $0 \leq t < 1$, every multiple real zero of F_t is a double zero at which F_t is locally ≤ 0 ; equivalently, every critical point of $x \mapsto S_k(x)$ with value in $(0, 1]$ is a non-degenerate local maximum ($S''_k < 0$). The weaker conditions “no F_t , $0 \leq t < 1$, has a real zero of even multiplicity at which it is locally non-negative”, “no two components of P_t merge as t increases” and “ S_k has no local minimum with value in $(0, 1]$ ” are equivalent to one another and necessary for (R); they are sufficient when no F_t , $0 \leq t < 1$, has a real zero of multiplicity at least 3 (no critical point of S_k with $S''_k = 0$ and value in $(0, 1]$), since at such a zero at least one conjugate pair leaves the axis just after t_0 , by (d).*

Proof. (a) Lemma 2.2. (b) Where $\Xi \leq 0$, $F_t = \Xi - L_t < 0$; in particular $F_t(\gamma_j) < 0$ for every j . At the origin $F_t(0) \geq \Xi_1(0) > 0$, because $F_t - \Xi_1 = \sum_{2 \leq n \leq k} \Phi_n + t\Phi_{k+1} \geq 0$ at real points (Lemma 2.1)

and $\Xi_1(0) = \Xi(0) - Q_1(0) \geq 0.4971 - c_1 > 0$, where $c_1 = \sum_{n \geq 2} (4\pi n^2 - 1)e^{-\pi n^2} \geq Q_1(0)$ [Tya26, proof of Corollary 3.5]. For large x , $x^2 F_t(x)$ tends to a negative limit: $\Xi(x)$ decays exponentially, and two integrations by parts, as in the proof of [Tya26, Proposition 3.4] applied to the kernel κ_t , give $\lim_{x \rightarrow \infty} x^2 L_t(x) = -2\kappa'_t(0) > 0$. So the zeros of F_t in $[0, \infty)$ lie in a bounded set and are finite in number. The parity statements follow from the signs at the ends of each lobe. (c) $F_s(x) > 0$ implies $F_t(x) > F_s(x) > 0$; a boundary point of P_s has $F_s = 0$ there, so $F_t > 0$ there. (d) $F_t(x) = F_{t_0}(x) + (t - t_0)\Phi_{k+1}(x)$ with $\Phi_{k+1}(x_0) > 0$; near x_0 this is $c(x - x_0)^m(1 + o(1)) + (t - t_0)\Phi_{k+1}(x_0)(1 + o(1))$, and the cases are read off; Weierstrass preparation gives exactly m zeros near x_0 , real or in conjugate pairs. (e) From (d), since $F_t(x) = \Phi_{k+1}(x)(S_k(x) - u)$ on the real axis. $\square$

Proposition 2.5 (conditional on the zeros of Ξ being real). *Assume every zero of Ξ is real. If F_{t_0} has a lift-off at x_0 (Theorem 2.4(d)), then $-(\log L_{t_0})''(x_0) \geq \sum_{\gamma > 0} [(x_0 - \gamma)^{-2} + (x_0 + \gamma)^{-2}]$.*

Proof. At a lift-off $\Xi = L$, $\Xi' = L'$ and $\Xi'' \geq L''$ at x_0 , with $L = L_{t_0} > 0$; so $(\log \Xi)''(x_0) = \Xi''/\Xi - (\Xi'/\Xi)^2 \geq L''/L - (L'/L)^2 = (\log L)''(x_0)$. With only real zeros, $\Xi(z) = \Xi(0) \prod_{\gamma > 0} (1 - z^2/\gamma^2)$ (Ξ is even, of order 1, and $\sum \gamma^{-2}$ converges), so $(\log \Xi)''(x) = -\sum_{\gamma > 0} [(x - \gamma)^{-2} + (x + \gamma)^{-2}]$. $\square$

So, under that hypothesis, (R) follows from the strict inequality $-(\log L_t)''(x) < \sum_{\gamma > 0} [(x - \gamma)^{-2} + (x + \gamma)^{-2}]$ at the real points where $\Xi = L_t$. The right side is at least $2 \sum \gamma^{-2} = 0.0462 \dots$ in the central lobe and at least $8(\gamma_{j+1} - \gamma_j)^{-2}$ in the lobe (γ_j, γ_{j+1}) (the two endpoint terms, smallest at the midpoint); the left side is, heuristically, of the order of the squared width of κ_t , about $1/(2\pi^2(k+1)^4)$. A proved bound for $-(\log L_t)''$ would give (R) for a range of k under the hypothesis of Proposition 2.5; with a hypothesis on the zeros of Ξ up to a finite height only, the product formula would need a remainder for the zeros above that height. The inequality is tabulated in Section 5.

3 The frozen level

By Lemma 2.2 the zeros of F_t are the solutions of $\Xi(z) = L_t(z)$. If the level were a constant $c > 0$ that decreases with t , (D) and (R) would be statements about the solutions of $\Xi = c$. When the zeros of Ξ are real these are known in substance, from the formula for $\text{Im } f'/f$ in Titchmarsh [Tit39, §8.52, p. 266] and from the level-curve statements of Csordas and Smith [CS00, (2.5), p. 604; (3.3)–(3.4), p. 609]; we state them in the form needed here, with a short proof.

Lemma 3.1 (Titchmarsh [Tit39, §8.52, p. 266]). *Let f be a real entire function of the form $f(z) = Cz^m e^{-az^2 + bz} \prod_j (1 - z/a_j) e^{z/a_j}$ with $a \geq 0$, b and the a_j real, $\sum_j a_j^{-2} < \infty$, and at least one zero or $a > 0$. Then $\text{Im } f'(z)/f(z) < 0$ for $\text{Im } z > 0$. If $c \neq 0$ is real and $z(c)$ is a branch of simple solutions of $f(z) = c$ in the open upper half-plane, then $d(\text{Im } z)/d|c| > 0$: as $|c|$ decreases, the solution moves toward the real axis.*

Proof. $f'/f = m/z - 2az + b + \sum_j [1/(z - a_j) + 1/a_j]$, and each of the terms m/z , $-2az$, $1/(z - a_j)$ has non-positive imaginary part for $\text{Im } z > 0$, with strict inequality for at least one. Then $dz/dc = 1/f'(z) = (1/c)(f/f')(z)$ because $f(z) = c$, and $\text{Im}(f/f') = -\text{Im}(f'/f)/|f'/f|^2 > 0$. $\square$

Proposition 3.2 (the solutions of $\Xi = c$ when the zeros of Ξ are real; in substance [CS00]). *Assume every zero of Ξ is real and simple, and let M_m be the maximum of Ξ over the m -th positive lobe $(\gamma_{2m}, \gamma_{2m+1})$, $m \geq 1$.*

- (i) For every $y > 0$ the continuous argument of Ξ along the horizontal line through iy , normalized by $\arg \Xi(iy) = 0$, is strictly decreasing in x and tends to $-\infty$; so for each $m \geq 1$ there is exactly one point $x_m(y) + iy$ with $x_m(y) > 0$ and $\arg \Xi = -2\pi m$. The curves $C_m := \{x_m(y) + iy : y > 0\}$ are disjoint real-analytic graphs over the y -axis, and $\{z : \operatorname{Re} z > 0, \operatorname{Im} z > 0, \Xi(z) > 0\} = \bigcup_{m \geq 1} C_m$.
- (ii) Along C_m the positive number Ξ is strictly increasing in y ; as $y \rightarrow 0$ the curve tends to the critical point of Ξ in the m -th positive lobe, and $\Xi \rightarrow M_m$.
- (iii) Hence, for a constant $c > 0$, the zeros of $\Xi - c$ in the open first quadrant are in bijection with the positive lobes with $M_m < c$: one zero on each such C_m , none on the others, which carry two real zeros of $\Xi - c$ in the lobe when $M_m > c$. As c decreases each such zero moves down its own curve C_m , strictly monotonically in $\operatorname{Im} z$, and reaches the real axis exactly when $c = M_m$, at the critical point of the lobe.

Proof. (i) Under the hypothesis $\Xi(z) = \Xi(0) \prod (1 - z^2/\gamma^2)$, and by Lemma 3.1 $\partial_x \arg \Xi(x + iy) = \operatorname{Im} \Xi'/\Xi \leq -\sum \gamma y/((x - \gamma)^2 + y^2)$, whose integral over x diverges because there are infinitely many γ ; $\Xi(iy) = 2 \int_0^\infty \sum_n \tilde{\varphi}_n(v) \cosh(yv) dv > 0$; the implicit function theorem gives the graphs. (ii) By Lemma 3.1, $\Xi' \neq 0$ off the real axis and Ξ , real on C_m , increases with y along it (that such level curves have no horizontal tangent is [CS00, (3.3)–(3.4)]). Near a real point that is neither a critical point nor a zero of Ξ , and near a simple zero, Ξ is injective and maps the real axis onto real values; so C_m can approach the real axis only at a critical point where $\Xi > 0$. As $y \rightarrow 0$ the argument tends, locally uniformly away from the zeros, to its boundary values ($-\pi$ per simple zero passed: $-2\pi m$ on the m -th positive lobe, $-2\pi m \pm \pi$ on the adjacent negative lobes), so every limit point of C_m is a critical point of Ξ in the closed lobe, not an endpoint. Under the hypothesis $(\Xi'/\Xi)'(x) = -\sum \gamma [(x - \gamma)^{-2} + (x + \gamma)^{-2}] < 0$, so Ξ'/Ξ falls strictly from $+\infty$ to $-\infty$ on each lobe: there is exactly one critical point, where $\Xi'' = \Xi \cdot (\Xi'/\Xi)' < 0$. (iii) For $y \geq 2\pi e^2$, $\sigma := \frac{1}{2} + y$ and $x \geq 0$, $\Xi(x + iy) = \xi(\sigma + ix)$ and $\arg \xi(\sigma + ix) \geq (x/2) \log(\sigma/(2\pi e)) - \pi/2$ (from $\arg \Gamma(s/2) = \frac{1}{2} \int_0^x \operatorname{Re} \psi((\sigma + i\tau)/2) d\tau$, $\operatorname{Re} \psi(a + ib) \geq \psi(a) \geq \log a - 1/a$, $\arg s(s-1) \in [0, \pi)$ and $|\arg \zeta(s)| < \pi/2$ for $\sigma \geq 2$); so $x_m(y) \leq 4\pi m + \pi$, and $|\xi(\sigma + ix)| \rightarrow \infty$ uniformly for $0 \leq x \leq 4\pi m + \pi$ as $\sigma \rightarrow \infty$. Thus Ξ maps C_m increasingly onto (M_m, ∞) ; the two real zeros in a lobe with $M_m > c$ come from its single critical point. $\square$

What this says about the pencil. If L_t were a constant that decreases with t , (D) and (R) would be exactly Proposition 3.2(iii): every non-real zero descends, each lands at the top of its own lobe when the level passes that top, and landed zeros stay real. The order in which the lobes are reached is the order of their heights M_m , not of their positions. This is how Haglund's Conjecture 1 (for every N , the zeros of Ξ_N in the closed first quadrant, listed by increasing real part, have nondecreasing imaginary parts [Hag09, p. 3]) fails at $N = 27$ [Tya26]: a low lobe to the left of a higher one. It is compatible with (D): the zero over the low lobe is still descending when the zeros of the higher lobe to its right have landed. The true level $L_t(z)$ is not constant: on the real axis it varies on the scale $2\pi(k+1)^2$, and off the axis it has a small phase (Section 6).

The exact criterion, with no hypothesis. At a simple zero z of F_t with $\operatorname{Im} z > 0$ that is not a common zero of Ξ and L_t , $\Xi(z) = L_t(z) \neq 0$ and $dz/dt = -\rho(z)/\Lambda(z)$ with $\rho := \Phi_{k+1}/L_t$ and $\Lambda := \Xi'/\Xi - L'_t/L_t$, so the zero descends if and only if $\operatorname{Im}(\rho/\Lambda) > 0$. In the frozen model $\rho > 0$ and $\Lambda = \Xi'/\Xi$.

4 A zero of Ξ off the real axis

Lemma 4.1 (Φ_n is nearly constant on a fixed disc). *Let $R > 0$ and $n \geq 2$ with $\pi n^2 \geq R + \frac{1}{2}$. Then for $|z| \leq R$,*

$$\left| \frac{\Phi_n(z)}{\Phi_n(0)} - 1 \right| \leq \frac{3.1 R^2}{\pi^2 n^4}.$$

Proof. Put $X = \pi n^2$. For $v \geq 0$, $|\cos(zv) - 1| \leq \cosh(Rv) - 1 \leq \frac{1}{2} R^2 v^2 e^{Rv}$. With $y = X e^{2v}$ (so $v = \frac{1}{2} \log(y/X)$, $dv = dy/2y$) one has $\tilde{\varphi}_n(v) dv = (2y - 3)(y/X)^{1/4} e^{-y} dy$. Hence $\int_0^\infty \tilde{\varphi}_n dv \geq \int_X^\infty (2y - 3) e^{-y} dy = (2X - 1) e^{-X}$. With $w = y - X$: $\log(y/X) \leq w/X$ and $(y/X)^{1/4 + R/2} \leq e^{(1/4 + R/2)w/X} \leq e^{w/2}$ because $X \geq \frac{1}{2} + R$; so $\int_0^\infty \tilde{\varphi}_n(v) \cdot \frac{1}{2} R^2 v^2 e^{Rv} dv \leq (R^2/8X^2) e^{-X} \int_0^\infty (2X + 2w) w^2 e^{-w/2} dw = (R^2/8X^2)(32X + 192) e^{-X}$. The quotient is at most $(R^2/8X^2)(32X + 192)/(2X - 1) \leq 3.1 R^2/X^2$ for $X \geq 4\pi$. $\square$

Consequences. For a fixed disc $|z| \leq R$ and $k \rightarrow \infty$, uniformly in $0 \leq t \leq 1$: $\Phi_{k+1}(z)/\Phi_{k+1}(0) \rightarrow 1$; $\sup |L_t| \leq \sum_{n \geq k} \sup |\Phi_n| \leq \Phi_{k+1}(0)(1 + o(1)) \rightarrow 0$, because the ratios $\Phi_{n+1}(0)/\Phi_n(0)$ tend to 0, since $(2X - 1)e^{-X} \leq \frac{1}{2} \Phi_n(0) \leq (2X + 3)e^{-X}$ for $X = \pi n^2 \geq 4\pi$ (the upper bound from $2y - 3 \leq 2y$ and $(y/X)^{1/4} \leq e^{w/4X}$, which give $e^{-X}[2X/(1-\varepsilon) + 2/(1-\varepsilon)^2]$ with $\varepsilon = 1/4X \leq 0.02$); and $L'_t \rightarrow 0$ on any smaller disc, by Cauchy's estimate.

Proposition 4.2. *Let β be a simple zero of Ξ with $\text{Im } \beta > 0$ and $\text{Im } \Xi'(\beta) \neq 0$. Then there are a disc D about β and k_0 such that for every $k \geq k_0$ and every $t \in [0, 1]$ the function $F_t = \Xi_k + t\Phi_{k+1}$ has exactly one zero $z_k(t)$ in D , it is simple, $t \mapsto z_k(t)$ is continuously differentiable, and $\text{sign } d(\text{Im } z_k(t))/dt = \text{sign } \text{Im } \Xi'(\beta)$ for all $t \in [0, 1]$. In particular, if Ξ had a simple zero β in the open upper half-plane with $\text{Im } \Xi'(\beta) > 0$, then for every $k \geq k_0$ the pencil would have a non-real zero whose imaginary part strictly increases on the whole interval $0 \leq t \leq 1$, and (D) would fail for all large k .*

Proof. Choose D with closure in the open upper half-plane, containing no other zero of Ξ and no zero of Ξ' , and let $\delta = \min_{\partial D} |\Xi| > 0$. By the consequences of Lemma 4.1, $\sup_{\overline{D}} |L_t| < \delta$ for $k \geq k_1$ and all t , so by Rouché's theorem $F_t = \Xi - L_t$ has exactly one zero $z_k(t)$ in D , and it is simple; it depends continuously differentiably on t by the implicit function theorem, with $dz_k/dt = -\Phi_{k+1}(z_k)/(\Xi'(z_k) - L'_t(z_k))$. As $k \rightarrow \infty$, uniformly in t : $z_k(t) \rightarrow \beta$ (the zero of $\Xi - L_t$ in D tends to the zero of Ξ as $\sup |L_t| \rightarrow 0$), $\Xi'(z_k) - L'_t(z_k) \rightarrow \Xi'(\beta) \neq 0$ and $\Phi_{k+1}(z_k)/\Phi_{k+1}(0) \rightarrow 1$. Therefore $\Phi_{k+1}(0)^{-1} dz_k/dt \rightarrow -1/\Xi'(\beta)$ uniformly on $[0, 1]$, and $\text{Im}(-1/\Xi'(\beta)) = \text{Im } \Xi'(\beta)/|\Xi'(\beta)|^2 \neq 0$ has the sign of $\text{Im } \Xi'(\beta)$. $\square$

Remark 4.3. (1) The mirror pencil $\Xi + L_t$ (the weights $1, \dots, 1, 2 - t, 2, 2, \dots$ on $\Phi_1, \Phi_2, \dots$, still a series of Haglund's form [Hag09, (53)] with non-negative coefficients) has $dz/dt = +\Phi_{k+1}/(\Xi' + L'_t)$, so the sign is reversed: a simple off-axis zero with $\text{Im } \Xi'(\beta) \neq 0$ produces an ascending branch, for all large k , in exactly one of the two pencils. (2) Not covered: $\text{Im } \Xi'(\beta) = 0$, and multiple zeros off the axis. For a zero of multiplicity $m \geq 2$ the m nearby zeros of F_t sit at β plus the m -th roots of $L_t(\beta)/c$, $c = \Xi^{(m)}(\beta)/m!$, and approach β radially as t increases ($|L_t(\beta)|$ decreases with u) and as k grows; at least one approaches from below, and so ascends in t , when $m \geq 3$, and when $m = 2$ unless c is a positive real number. This is a sketch, not a proof. (3) Proposition 4.2 concerns large k only; it says nothing about a fixed k . (4) By the reflection $\Xi(-\bar{z}) = \overline{\Xi(z)}$, the zero $-\bar{\beta}$ has $\Xi'(-\bar{\beta}) = -\overline{\Xi'(\beta)}$, with the same sign of the imaginary part; the two mirror zeros behave alike.

Read in the other direction, Proposition 4.2 says that (D) for all large k excludes the simple zeros of Ξ off the real axis with $\text{Im } \Xi'(\beta) > 0$; simple ones with $\text{Im } \Xi'(\beta) < 0$ descend in this pencil and are not excluded, and multiple ones are not treated.

5 The census

Everything in this section is numerical: floating point, not interval arithmetic. Two programs were written independently. *Program A* evaluates Φ_n and S_k in Arb ball arithmetic [Joh17, FLINT], using a value only when its relative radius is below 2^{-40} ; it follows each branch as the level curve $\text{Im } S_k = 0$ (Lemma 2.3) from a zero of Ξ_k to a landing or to a zero of Ξ_{k+1} , and counts zeros by the argument principle. *Program B*, written in mpmath [mpm] from Haglund’s definitions without sight of program A, follows each branch by Newton continuation on a grid in $\tau = -\log(1 - t)$; it reproduces Haglund’s 19 printed zeros of Ξ_1 to $7 \cdot 10^{-22}$, the largest real zeros of $\Xi_1, \dots, \Xi_4$ in his table, and the certified zeros of Ξ_{27} of [Tya26].

k	$\text{Re } z$	$\text{Im } z$	real	non-real	B	L	E	X	Δ^+	m	prog.
1	[0, 86.55]	[0, 41]	1/7	15/11	15	3	11	1	$-3.7 \cdot 10^{-7}$	0.708	A, B
2	[0, 130.53]	[0, 50]	7/15	23/18	23	4	18	1	$-3.1 \cdot 10^{-7}$	0.806	A, B
3	[0, 187.08]	[0, 59]	15/31	35/25	35	8	25	2	$-3.3 \cdot 10^{-7}$	0.812	A, B
4	[0, 256.19]	[0, 68]	31/53	47/34	47	11	34	2	$-3.2 \cdot 10^{-7}$	0.797	A
5	[0, 337.88]	[0, 79]	53/79	63/47	63	13	47	3	$-3.0 \cdot 10^{-7}$	0.828	A
6	[0, 432.12]	[0, 90]	79/113	82/62	82	17	62	3	$-3.0 \cdot 10^{-7}$	0.832	A
7	[0, 538.94]	[0, 102]	113/155	103/80	103	21	80	2	$-3.1 \cdot 10^{-7}$	0.842	A
8	[0, 658.32]	[0, 115]	155/207	127/98	127	26	98	3	$-3.0 \cdot 10^{-7}$	0.850	A
15	[984, 1196]	[0, 55]	37/145*	74/16	74	54	16	4	$-3.0 \cdot 10^{-7}$	0.880	A
20	[1724, 1976]	[0, 61]	40/196*	101/17	101	78	17	6	$-3.0 \cdot 10^{-7}$	0.876	A
26	[2876, 3176]	[0, 69]	45/263*	133/17	133	109	17	7	$-3.0 \cdot 10^{-7}$	0.868	A
27	[3096, 3404]	[0, 70]	48/276*	137/16	137	114	16	7	$-3.0 \cdot 10^{-7}$	0.851	A
35	[5144, 5516]	[0, 80]	53/363*	182/19	182	155	19	8	$-3.0 \cdot 10^{-7}$	0.876	A
50	[10364, 10856]	[0, 99]	58/546*	274/19	274	244	19	11	$-3.0 \cdot 10^{-7}$	0.857	A

Table 1: Branches of zeros of F_t , $0 \leq t \leq 1$, in the window W given by the ranges of $\text{Re } z$ and $\text{Im } z$. Upper block: complete windows, $\text{Re } z \leq 2\pi(k+2)^2 + 30$; lower block: windows around the landings, branches that start in them. Real, non-real: zeros of Ξ_k / Ξ_{k+1} in W (*: real zeros in the window). B: branches followed; L: landings; E: ends at a non-real zero of Ξ_{k+1} ; X: exits through the side of W . Δ^+ : the largest change of $\text{Im } z$ over one step, all branches (negative: $\text{Im } z$ decreased at every step). m : the smallest margin $-\text{Im } S'_k / |S'_k|$ above height 10^{-6} . Floating point, not interval arithmetic.

Readings (each a statement about its window). In every row of Table 1 the imaginary part decreased at every step of every branch and the margin stays above 0.7; the landings are half the difference of the real counts; and every non-real zero of Ξ_{k+1} in the window ends a followed branch (for $k = 6$ one branch reaches $t = 1$ at the zero $432.1548 + 63.0618i$ of Ξ_7 , just outside the window, and is counted as an exit). For $k = 1, 2, 3$ the two programs (B with window heights 45, 45, 60) agree on every count, and on all 15 landings to the digits B printed: $|\Delta x^*| \leq 3.9 \cdot 10^{-7}$, $|\Delta \tau^*| \leq 4.8 \cdot 10^{-6}$; B finds $\text{Im}(dz/dt) < 0$ at all 3,605, 6,886 and 11,974 grid values of its branches. The real counts of the upper block agree with Haglund’s table [Hag11w, p. 4], which has 31 at $N = 4$ (the arXiv and journal versions print 32 [Hag09, p. 4]).

The real axis. For $k = 1, \dots, 50$, on $[0, 4(k+2)^2 + 40]$, program A finds that S_k has no local minimum with value in $(0, 1)$ and that its local maxima with value in $(0, 1)$ are $(R_{k+1} - R_k)/2$ in number, where R_N is the number of real zeros of Ξ_N found on that interval (grid of $1/20$ of the mean zero spacing $2\pi/\log(x/2\pi)$, the part beyond $4(k+1)^2 - 40$ scanned again at $1/40$). Each R_N with $2 \leq N \leq 50$ is found twice, on the intervals of the pencils $k = N - 1$ and $k = N$, with the

same value. For $N = 1, \dots, 51$: 1, 7, 15, 31, 53, 79, 113, 155, 207, 263, 327, 401, 483, 575, 673, 781, 899, 1027, 1163, 1307, 1463, 1631, 1807, 1991, 2189, 2393, 2611, 2839, 3075, 3323, 3581, 3851, 4133, 4423, 4727, 5037, 5363, 5699, 6045, 6405, 6771, 7157, 7549, 7951, 8369, 8797, 9237, 9689, 10155, 10629, 11117 — all odd, as [Tya26, Corollary 3.5] requires. Program B finds the same for $k = 1, 2, 3$ on $[0, 98.55]$, $[0, 142.5]$, $[0, 199.1]$. A scan for minima checks the criterion of Theorem 2.4(e) up to degenerate critical points ($S'_k = S''_k = 0$ at one x , a non-generic event at which (R) and (D) fail, by Theorem 2.4(d); none was met). The inequality in the first step of the proof of Proposition 2.5, which uses no hypothesis (a lift-off at x_0 forces $(\log \Xi)''(x_0) \geq (\log L_t)''(x_0)$), was tabulated on the part of $[0, 4(k+2)^2 + 40]$ where $\Xi > 0$ (step 0.02, $u \in \{0, \frac{1}{4}, \frac{1}{2}, \frac{3}{4}, 1\}$): there $-(\log L_u)''$ is at most 0.066 times $-(\log \Xi)''$ at $k = 1$ and at most $5.6 \cdot 10^{-6}$ times at $k = 20$.

The site of the earlier counterexample (program A). The certified non-real zero of Ξ_{27} near $3143.2206824215 + 0.3152587994i$ [Tya26, Theorem 1.1] is the end, at $t = 1$, of the branch of the pencil $k = 26$ that starts at the zero $3130.2619 + 52.4297i$ of Ξ_{26} ; it descends throughout (margin at least 0.970). Its right neighbor, from $3132.1660 + 52.9056i$, lands at $x^* = 3145.2290390$ at $u^* = 3.3 \cdot 10^{-76}$ and gives the real zeros 3144.89466 and 3145.59985 of Ξ_{27} . In the pencil $k = 27$ the first zero lands at $x^* = 3143.2466268840$ at $u^* = 0.41051$ ($t = 0.58949$) and becomes the real zeros 3142.979457 and 3143.567499 of Ξ_{28} ; the lobe scan of [Tya26, Section 5] had put the height of that lobe at 0.411 of the level Q_{27} , which is the same number seen from the real axis. So the failure of Conjecture 1 at $N = 27$ is a branch that had not yet landed when its neighbor on the right had: compatible with (D), as the frozen-level picture of Section 3 says.

Controls. The tests can fail. For $f = \Xi + \lambda$, $\lambda = 5 \cdot 10^{-5}$, which has zeros off the real axis, the pencil $(\Xi_k + \lambda) + t\Phi_{k+1} = f - L_t$ has real-axis quotient $(\Xi_{k+1} + \lambda)/\Phi_{k+1}$. For $k = 1$ it has a local minimum at 24.34012104 with value 0.611995156 (three independent codes), so (R) fails; and the branch near the off-axis zero $28.6324 + 8.5243i$ of f , where $\text{Im } f' > 0$, ascends for $k = 2, \dots, 8$, as Theorem 2.4 and Proposition 4.2 (applied to f) predict.

Far field. For $k = 2$, four zeros of Ξ_2 near $500 + 160i$ were followed through the pencil: they descend at every step, by 8.44–8.47 in the imaginary part, the amount the formula of Section 6(III) predicts to three digits.

6 The true level: heuristics

This section is heuristic, except Lemma 6.1. Write $A(s) := \frac{1}{2}s(s-1)\pi^{-s/2}\Gamma(s/2)$ and $a := \pi(k+1)^2$, the second argument of the incomplete gamma functions in Φ_{k+1} . There are three regimes for a zero $z = x + iy$ of F_t in the first quadrant.

(I) *The frontier:* x from about $4(k+1)^2$ to $4(k+2)^2$, y from 0 to a few units. Here Ξ and L_t have comparable size ($|\Xi(x)| \approx e^{-\pi x/4} x^{7/4}$ against $L_t \approx u\Phi_{k+1}(0) \approx 4ua e^{-a}$), and all landings happen here. For $k \geq 3$ the frontier lies below the height $x \approx 2a$ where Φ_{k+1} changes regime ($4(k+2)^2 < 2\pi(k+1)^2$: by 0.5% at $k = 3$, 8% at $k = 4$, the ratio tending to $2/\pi$); for $k = 1, 2$ the regimes overlap. L_t is then slowly varying: its logarithmic derivative is of size $x/(2a^2)$ (from the second moment of the kernel, about $1/(2a^2)$), against $|\Xi'/\Xi| \approx \frac{1}{2} \log(x/2\pi)$ and $-\text{Im } \Xi'/\Xi = y \sum_\gamma |z - \gamma|^{-2}$. The criterion $\text{Im}(\rho/\Lambda) > 0$ of Section 3 holds with a wide margin in this model; the neglected terms are $O(xy/a^2)$ in the phase of ρ and $O(x/a^2)$ in Λ .

(II) *The middle:* x between the frontier and about $2a$, on the curve where $|\Xi(z)| = |L_t(z)|$, which rises like $(\pi/2)(x - x_{\text{frontier}})/\log(x/2\pi)$, since $\partial_y \log |\Xi| = \frac{1}{2} \log(x/2\pi)$ and $\partial_x \log |\Xi| \approx -\pi/4$ there. Above height $\frac{1}{2}$, $\Xi(z) = A(\frac{1}{2} - iz)\zeta(\frac{1}{2} - iz)$, with $\zeta(\frac{1}{2} - iz)$ within $\zeta(\frac{1}{2} + y) - 1$ of 1 (34% at $y = 2$, 20% at $y = 2.5$), $\Lambda \approx -\pi/4 - (i/2) \log(x/2\pi)$, and the zero descends as long as the phase of ρ stays below $\arctan(2 \log(x/2\pi)/\pi) \approx 1.2$. The phase of $\rho = \Phi_{k+1}/L_t$ is 0 when $u \gg Q_{k+1}/\Phi_{k+1}$,

and otherwise at most the phase of $\Phi_{k+2}(z)/\Phi_{k+1}(z)$, which the leading terms of the incomplete gamma functions put at a few hundredths to a tenth of a radian there.

(III) *Far out:* $x \gg 2a$. All terms $n \leq k+1$ are in their oscillatory regime, the zero condition is $A(\frac{1}{2} - iz)D(z) = (\text{the algebraic part})$ with D close to 1, and the algebraic part is $c(t)/z^2 + O(z^{-4})$ with $c(t) := \lim_{x \rightarrow \infty} x^2 F_t(x)$. The sign and the monotonicity of $c(t)$ are proved:

Lemma 6.1. *For $k \geq 1$ and $0 \leq t \leq 1$, $c(t) = 2\kappa'_t(0) = -2[u\alpha_{k+1} + \sum_{n>k+1} \alpha_n]$, where $\alpha_n := -\tilde{\varphi}'_n(0) = (8X^3 - 30X^2 + 15X)e^{-X} > 0$ for $n \geq 2$, $X = \pi n^2$. Hence $c(t) < 0$, and $|c(t)|$ is strictly decreasing in t .*

Proof. $x^2 \Xi(x) \rightarrow 0$ and $x^2 L_t(x) \rightarrow -2\kappa'_t(0)$ (proof of Theorem 2.4(b)), so $c(t) = 2\kappa'_t(0)$; the values $\tilde{\varphi}'_n(0) = -(8X^3 - 30X^2 + 15X)e^{-X}$ and their signs for $n \geq 2$ are [Tya26, Proposition 3.4 and its proof]. As t increases, $u = 1 - t$ decreases and $\alpha_{k+1} > 0$. $\square$

Heuristically, a far zero then moves by $\delta z \approx (\delta c/c)/(T'/T)$, T the dominant gamma term, and $\text{Im}(1/(T'/T)) > 0$ there, so a decrease of $|c|$ moves it down. The arXiv and journal versions of Haglund’s paper state that the coefficient of $1/x^2$ in $\Xi_N(x)$ “is positive for $k \geq 3$ and negative for $1 \leq k < 3$ ” [Hag09, p. 10]; with that sign the far zeros would have to move up as t increases. The author’s later copy states instead that the coefficient “approaches zero from below as $N \rightarrow \infty$ ” [Hag11w, p. 10], and it is negative for every N [Tya26, Proposition 3.4]: that is the sign consistent with Conjecture 4. The far-field check of Section 5 tests the formula for δz at $k = 2$.

Turning (I)–(III) into proofs would need, among other things, uniform estimates of $\Phi_n(z)$ near $x \approx 2\pi n^2$, where the incomplete gamma function changes regime.

7 Prior work

Haglund’s paper exists in three versions [Hag09, Hag11, Hag11w]; the author’s copy of February 9, 2011 [Hag11w] has the entry 31 at $N = 4$ and the corrected sentence on the sign of the $1/x^2$ coefficient (Sections 5 and 6).

Conjecture 4 for $k = 1$. Baccaro’s preprint [Bac26] states Conjecture 4 for $k = 1$ as its Theorem 2.1 [Bac26, p. 2] and proves it with an interval-arithmetic certificate on a bounded region together with an analytic estimate for the zeros with $|z| \geq 256$ [Bac26, (11), p. 3]; its abstract says: “These results prove Haglund’s Conjecture 4 for $k = 1$. The cases $k \geq 2$ remain open.” It uses $\Phi_2 > 0$ on the real axis, with the convex-kernel reason [Bac26, (12), p. 3] (Lemma 2.1 for $n = 2$), and its criterion for descent is that of Lemma 2.3 for $k = 1$ in the normalization $g = -\Phi_1/\Phi_2 = 1 - S_1$ [Bac26, (7)–(10), p. 2]. The positivity of Φ_n on the real axis for every $n \geq 2$ is also recorded in the repository that accompanies [Bac26]. Ahn’s thesis [Ahn12] computes the zeros of the fixed members $\Phi_1 + t\Phi_2$, $t = 0.1, 0.5, 0.9, 0.95$, among fourteen combinations, in the rectangle $0.01 \leq x \leq 250$, $0 \leq y \leq 50$, and tests the ordering of Conjecture 1 for each; it does not follow zeros in t .

Searches. Among the works citing [Hag09] or [Hag11] that we found (through Semantic Scholar, OpenAlex, Google Scholar and zbMATH Open, and searches of Zenodo, MathOverflow and the web), and in a second, independent web search, only [Bac26] states, computes or proves anything about the zeros of $\Xi_k + t\Phi_{k+1}$ as t varies; we found nothing for any $k \geq 2$.

Section 3. Titchmarsh’s formula for $\text{Im } f'/f$ and the inequality $d(f'/f)/dz < 0$ on the real axis [Tit39, §8.52, p. 266] are stated for real entire functions with real zeros of order less than 2, or of order 2 and genus 1. Csordas and Smith prove $\text{Im } f'/f < 0$ above the strip $|\text{Im } z| \leq A$ containing the zeros, for their class $S_\infty(A)$ [CS00, Definition 1.2, p. 602; (2.5), p. 604]; Ξ belongs

to it with $A = \frac{1}{2}$, so $\text{Im } \Xi' / \Xi < 0$ for $\text{Im } z > \frac{1}{2}$ with no hypothesis, and there the level curves have no horizontal tangent [CS00, (3.3)–(3.4), p. 609].

Code availability. <https://github.com/x67ai/riemann-rh-program>.

References

- [Ahn12] S.-H. (Shirley) Ahn, *On the zeros of component functions of the Riemann Ξ approximates*, M.A. thesis, University of Pennsylvania, 2012.
- [Bac26] M. L. Baccaro, *Haglund’s Zero-Trajectory Conjecture for the First Riemann Ξ Approximant*, preprint, version v1.0.0, Zenodo, August 22, 2026, DOI 10.5281/zenodo.22059236.
- [CS00] G. Csordas and W. Smith, *Level sets and the distribution of zeros of entire functions*, Michigan Math. J. **47** (2000), 601–612.
- [FLINT] The FLINT team, *FLINT: Fast Library for Number Theory*, version 3.0.1, <https://flintlib.org>; used through python-flint 0.6.0, <https://github.com/flintlib/python-flint>.
- [Hag09] J. Haglund, *Some conjectures on the zeros of approximates to the Riemann Ξ -function and incomplete gamma functions*, arXiv:0910.5228v1, 27 October 2009. Page and equation numbers in this paper refer to this version.
- [Hag11] J. Haglund, *Some conjectures on the zeros of approximates to the Riemann Ξ -function and incomplete gamma functions*, Cent. Eur. J. Math. **9** (2011), no. 2, 302–318, DOI 10.2478/s11533-010-0095-3.
- [Hag11w] J. Haglund, *Some conjectures on the zeros of approximates to the Riemann Ξ -function and incomplete gamma functions*, author’s copy dated February 9, 2011, <https://www2.math.upenn.edu/~jhaglund/preprints/rh8.pdf> (read October 3, 2026).
- [Joh17] F. Johansson, *Arb: efficient arbitrary-precision midpoint-radius interval arithmetic*, IEEE Trans. Comput. **66** (2017), no. 8, 1281–1292, DOI 10.1109/TC.2017.2690633.
- [mpm] The mpmath development team, *mpmath: a Python library for arbitrary-precision floating-point arithmetic*, version 1.3.0, <https://mpmath.org>.
- [Tit39] E. C. Titchmarsh, *The Theory of Functions*, 2nd ed., Oxford University Press, Oxford, 1939.
- [Tya26] K. Tyagi, *A counterexample to Haglund’s monotonic-zeros conjecture for the Riemann Ξ approximants*, preprint, Zenodo, October 1, 2026, DOI 10.5281/zenodo.23071930.