

# Exercises and Problems in Linear Algebra

John M. Erdman  
Portland State University

Version July 13, 2014

©2010 John M. Erdman

*E-mail address:* `erdman@pdx.edu`



# Contents

PREFACE	vii
<b>Part 1. MATRICES AND LINEAR EQUATIONS</b>	<b>1</b>
Chapter 1. SYSTEMS OF LINEAR EQUATIONS	3
1.1. Background	3
1.2. Exercises	4
1.3. Problems	7
1.4. Answers to Odd-Numbered Exercises	8
Chapter 2. ARITHMETIC OF MATRICES	9
2.1. Background	9
2.2. Exercises	10
2.3. Problems	12
2.4. Answers to Odd-Numbered Exercises	14
Chapter 3. ELEMENTARY MATRICES; DETERMINANTS	15
3.1. Background	15
3.2. Exercises	17
3.3. Problems	22
3.4. Answers to Odd-Numbered Exercises	23
Chapter 4. VECTOR GEOMETRY IN $\mathbb{R}^n$	25
4.1. Background	25
4.2. Exercises	26
4.3. Problems	28
4.4. Answers to Odd-Numbered Exercises	29
<b>Part 2. VECTOR SPACES</b>	<b>31</b>
Chapter 5. VECTOR SPACES	33
5.1. Background	33
5.2. Exercises	34
5.3. Problems	37
5.4. Answers to Odd-Numbered Exercises	38
Chapter 6. SUBSPACES	39
6.1. Background	39
6.2. Exercises	40
6.3. Problems	44
6.4. Answers to Odd-Numbered Exercises	45
Chapter 7. LINEAR INDEPENDENCE	47
7.1. Background	47
7.2. Exercises	49

7.3. Problems	51
7.4. Answers to Odd-Numbered Exercises	53
<b>Chapter 8. BASIS FOR A VECTOR SPACE</b>	<b>55</b>
8.1. Background	55
8.2. Exercises	56
8.3. Problems	57
8.4. Answers to Odd-Numbered Exercises	58
<b>Part 3. LINEAR MAPS BETWEEN VECTOR SPACES</b>	<b>59</b>
<b>Chapter 9. LINEARITY</b>	<b>61</b>
9.1. Background	61
9.2. Exercises	63
9.3. Problems	67
9.4. Answers to Odd-Numbered Exercises	70
<b>Chapter 10. LINEAR MAPS BETWEEN EUCLIDEAN SPACES</b>	<b>71</b>
10.1. Background	71
10.2. Exercises	72
10.3. Problems	74
10.4. Answers to Odd-Numbered Exercises	75
<b>Chapter 11. PROJECTION OPERATORS</b>	<b>77</b>
11.1. Background	77
11.2. Exercises	78
11.3. Problems	79
11.4. Answers to Odd-Numbered Exercises	80
<b>Part 4. SPECTRAL THEORY OF VECTOR SPACES</b>	<b>81</b>
<b>Chapter 12. EIGENVALUES AND EIGENVECTORS</b>	<b>83</b>
12.1. Background	83
12.2. Exercises	84
12.3. Problems	85
12.4. Answers to Odd-Numbered Exercises	86
<b>Chapter 13. DIAGONALIZATION OF MATRICES</b>	<b>87</b>
13.1. Background	87
13.2. Exercises	89
13.3. Problems	91
13.4. Answers to Odd-Numbered Exercises	92
<b>Chapter 14. SPECTRAL THEOREM FOR VECTOR SPACES</b>	<b>93</b>
14.1. Background	93
14.2. Exercises	94
14.3. Answers to Odd-Numbered Exercises	96
<b>Chapter 15. SOME APPLICATIONS OF THE SPECTRAL THEOREM</b>	<b>97</b>
15.1. Background	97
15.2. Exercises	98
15.3. Problems	102
15.4. Answers to Odd-Numbered Exercises	103
<b>Chapter 16. EVERY OPERATOR IS DIAGONALIZABLE PLUS NILPOTENT</b>	<b>105</b>

16.1.	Background	105
16.2.	Exercises	106
16.3.	Problems	110
16.4.	Answers to Odd-Numbered Exercises	111
<b>Part 5.</b>	<b>THE GEOMETRY OF INNER PRODUCT SPACES</b>	<b>113</b>
Chapter 17.	COMPLEX ARITHMETIC	115
17.1.	Background	115
17.2.	Exercises	116
17.3.	Problems	118
17.4.	Answers to Odd-Numbered Exercises	119
Chapter 18.	REAL AND COMPLEX INNER PRODUCT SPACES	121
18.1.	Background	121
18.2.	Exercises	123
18.3.	Problems	125
18.4.	Answers to Odd-Numbered Exercises	126
Chapter 19.	ORTHONORMAL SETS OF VECTORS	127
19.1.	Background	127
19.2.	Exercises	128
19.3.	Problems	129
19.4.	Answers to Odd-Numbered Exercises	131
Chapter 20.	QUADRATIC FORMS	133
20.1.	Background	133
20.2.	Exercises	134
20.3.	Problems	136
20.4.	Answers to Odd-Numbered Exercises	137
Chapter 21.	OPTIMIZATION	139
21.1.	Background	139
21.2.	Exercises	140
21.3.	Problems	141
21.4.	Answers to Odd-Numbered Exercises	142
<b>Part 6.</b>	<b>ADJOINT OPERATORS</b>	<b>143</b>
Chapter 22.	ADJOINTS AND TRANSPOSES	145
22.1.	Background	145
22.2.	Exercises	146
22.3.	Problems	147
22.4.	Answers to Odd-Numbered Exercises	148
Chapter 23.	THE FOUR FUNDAMENTAL SUBSPACES	149
23.1.	Background	149
23.2.	Exercises	151
23.3.	Problems	155
23.4.	Answers to Odd-Numbered Exercises	157
Chapter 24.	ORTHOGONAL PROJECTIONS	159
24.1.	Background	159
24.2.	Exercises	160

24.3. Problems	163
24.4. Answers to Odd-Numbered Exercises	164
Chapter 25. LEAST SQUARES APPROXIMATION	165
25.1. Background	165
25.2. Exercises	166
25.3. Problems	167
25.4. Answers to Odd-Numbered Exercises	168
<b>Part 7. SPECTRAL THEORY OF INNER PRODUCT SPACES</b>	<b>169</b>
Chapter 26. SPECTRAL THEOREM FOR REAL INNER PRODUCT SPACES	171
26.1. Background	171
26.2. Exercises	172
26.3. Problem	174
26.4. Answers to the Odd-Numbered Exercise	175
Chapter 27. SPECTRAL THEOREM FOR COMPLEX INNER PRODUCT SPACES	177
27.1. Background	177
27.2. Exercises	178
27.3. Problems	181
27.4. Answers to Odd-Numbered Exercises	182
Bibliography	183
Index	185

## PREFACE

This collection of exercises is designed to provide a framework for discussion in a junior level linear algebra class such as the one I have conducted fairly regularly at Portland State University. There is no assigned text. Students are free to choose their own sources of information. Students are encouraged to find books, papers, and web sites whose writing style they find congenial, whose emphasis matches their interests, and whose price fits their budgets. The short introductory *background* section in these exercises, which precede each assignment, are intended only to fix notation and provide “official” definitions and statements of important theorems for the exercises and problems which follow.

There are a number of excellent online texts which are available free of charge. Among the best are *Linear Algebra* [7] by Jim Hefferon,

<http://joshua.smcvt.edu/linearalgebra>

and *A First Course in Linear Algebra* [2] by Robert A. Beezer,

<http://linear.ups.edu/download/fcla-electric-2.00.pdf>

Another very useful online resource is Przemyslaw Bogacki’s *Linear Algebra Toolkit* [3].

<http://www.math.odu.edu/~bogacki/lat>

And, of course, many topics in linear algebra are discussed with varying degrees of thoroughness in the *Wikipedia* [12]

<http://en.wikipedia.org>

and Eric Weisstein’s *Mathworld* [11].

<http://mathworld.wolfram.com>

Among the dozens and dozens of linear algebra books that have appeared, two that were written before “dumbing down” of textbooks became fashionable are especially notable, in my opinion, for the clarity of their authors’ mathematical vision: Paul Halmos’s *Finite-Dimensional Vector Spaces* [6] and Hoffman and Kunze’s *Linear Algebra* [8]. Some students, especially mathematically inclined ones, love these books, but others find them hard to read. If you are trying seriously to learn the subject, give them a look when you have the chance. Another excellent traditional text is *Linear Algebra: An Introductory Approach* [5] by Charles W. Curits. And for those more interested in applications both *Elementary Linear Algebra: Applications Version* [1] by Howard Anton and Chris Rorres and *Linear Algebra and its Applications* [10] by Gilbert Strang are loaded with applications.

If you are a student and find the level at which many of the current beginning linear algebra texts are written depressingly pedestrian and the endless routine computations irritating, you might examine some of the more advanced texts. Two excellent ones are Steven Roman’s *Advanced Linear Algebra* [9] and William C. Brown’s *A Second Course in Linear Algebra* [4].

Concerning the material in these notes, I make no claims of originality. While I have dreamed up many of the items included here, there are many others which are standard linear algebra exercises that can be traced back, in one form or another, through generations of linear algebra texts, making any serious attempt at proper attribution quite futile. If anyone feels slighted, please contact me.

There will surely be errors. I will be delighted to receive corrections, suggestions, or criticism at

erdman@pdx.edu

I have placed the the  $\text{\LaTeX}$  source files on my web page so that those who wish to use these exercises for homework assignments, examinations, or any other noncommercial purpose can download the material and, without having to retype everything, edit it and supplement it as they wish.



**Part 1**

**MATRICES AND LINEAR EQUATIONS**



## CHAPTER 1

# SYSTEMS OF LINEAR EQUATIONS

### 1.1. Background

**Topics:** systems of linear equations; Gaussian elimination (Gauss' method), elementary row operations, leading variables, free variables, echelon form, matrix, augmented matrix, Gauss-Jordan reduction, reduced echelon form.

**1.1.1. Definition.** We will say that an operation (sometimes called *scaling*) which multiplies a row of a matrix (or an equation) by a nonzero constant is a ROW OPERATION OF TYPE I. An operation (sometimes called *swapping*) that interchanges two rows of a matrix (or two equations) is a ROW OPERATION OF TYPE II. And an operation (sometimes called *pivoting*) that adds a multiple of one row of a matrix to another row (or adds a multiple of one equation to another) is a ROW OPERATION OF TYPE III.

**1.2. Exercises**

- (1) Suppose that  $L_1$  and  $L_2$  are lines in the plane, that the  $x$ -intercepts of  $L_1$  and  $L_2$  are 5 and  $-1$ , respectively, and that the respective  $y$ -intercepts are 5 and 1. Then  $L_1$  and  $L_2$  intersect at the point ( \_\_\_\_ , \_\_\_\_ ) .

- (2) Consider the following system of equations.

$$\begin{cases} w + x + y + z = 6 \\ w \quad + y + z = 4 \\ w \quad + y \quad = 2 \end{cases} \quad (*)$$

- (a) List the leading variables \_\_\_\_ .  
 (b) List the free variables \_\_\_\_ .  
 (c) The general solution of (\*) (expressed in terms of the free variables) is  
 (\_\_\_\_ , \_\_\_\_ , \_\_\_\_ , \_\_\_\_ ) .  
 (d) Suppose that a fourth equation  $-2w + y = 5$  is included in the system (\*). What is the solution of the resulting system? Answer: ( \_\_\_\_ , \_\_\_\_ , \_\_\_\_ , \_\_\_\_ ) .  
 (e) Suppose that instead of the equation in part (d), the equation  $-2w - 2y = -3$  is included in the system (\*). Then what can you say about the solution(s) of the resulting system? Answer: \_\_\_\_\_ .

- (3) Consider the following system of equations:

$$\begin{cases} x + y + z = 2 \\ x + 3y + 3z = 0 \\ x + 3y + 6z = 3 \end{cases} \quad (*)$$

- (a) Use Gaussian elimination to put the augmented coefficient matrix into row echelon form. The result will be  $\begin{bmatrix} 1 & 1 & 1 & a \\ 0 & 1 & 1 & b \\ 0 & 0 & 1 & c \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$  ,  $b = \underline{\hspace{1cm}}$  , and  $c = \underline{\hspace{1cm}}$  .  
 (b) Use Gauss-Jordan reduction to put the augmented coefficient matrix in reduced row echelon form. The result will be  $\begin{bmatrix} 1 & 0 & 0 & d \\ 0 & 1 & 0 & e \\ 0 & 0 & 1 & f \end{bmatrix}$  where  $d = \underline{\hspace{1cm}}$  ,  $e = \underline{\hspace{1cm}}$  , and  $f = \underline{\hspace{1cm}}$  .  
 (c) The solutions of (\*) are  $x = \underline{\hspace{1cm}}$  ,  $y = \underline{\hspace{1cm}}$  , and  $z = \underline{\hspace{1cm}}$  .

- (4) Consider the following system of equations.

$$0.003000x + 59.14y = 59.17$$

$$5.291x - 6.130y = 46.78.$$

- (a) Using only row operation III and back substitution find the exact solution of the system. Answer:  $x = \underline{\hspace{1cm}}$  ,  $y = \underline{\hspace{1cm}}$  .  
 (b) Same as (a), but after performing each arithmetic operation round off your answer to four significant figures. Answer:  $x = \underline{\hspace{1cm}}$  ,  $y = \underline{\hspace{1cm}}$  .

(5) Find the values of  $k$  for which the system of equations

$$\begin{cases} x + ky = 1 \\ kx + y = 1 \end{cases}$$

has (a) no solution. Answer: \_\_\_\_\_ .

(b) exactly one solution. Answer: \_\_\_\_\_ .

(c) infinitely many solutions. Answer: \_\_\_\_\_ .

(d) When there is exactly one solution, it is  $x =$  \_\_\_\_\_ and  $y =$  \_\_\_\_\_ .

(6) Consider the following two systems of equations.

$$\begin{cases} x + y + z = 6 \\ x + 2y + 2z = 11 \\ 2x + 3y - 4z = 3 \end{cases} \quad (1)$$

and

$$\begin{cases} x + y + z = 7 \\ x + 2y + 2z = 10 \\ 2x + 3y - 4z = 3 \end{cases} \quad (2)$$

Solve both systems simultaneously by applying Gauss-Jordan reduction to an appropriate  $3 \times 5$  matrix.

(a) The resulting row echelon form of this  $3 \times 5$  matrix is  $\begin{bmatrix} & & & & \\ & & & & \\ & & & & \end{bmatrix}$ .

(b) The resulting reduced row echelon form is  $\begin{bmatrix} & & & & \\ & & & & \\ & & & & \end{bmatrix}$ .

(c) The solution for (1) is (\_\_\_\_, \_\_\_\_, \_\_\_\_) and the solution for (2) is (\_\_\_\_, \_\_\_\_, \_\_\_\_).

(7) Consider the following system of equations:

$$\begin{cases} x - y - 3z = 3 \\ 2x + z = 0 \\ 2y + 7z = c \end{cases}$$

(a) For what values of  $c$  does the system have a solution? Answer:  $c =$  \_\_\_\_\_ .

(b) For the value of  $c$  you found in (a) describe the solution set geometrically as a subset of  $\mathbb{R}^3$ . Answer: \_\_\_\_\_ .

(c) What does part (a) say about the planes  $x - y - 3z = 3$ ,  $2x + z = 0$ , and  $2y + 7z = 4$  in  $\mathbb{R}^3$ ? Answer: \_\_\_\_\_ .



**1.3. Problems**

- (1) Give a geometric description of a single linear equation in three variables.

Then give a geometric description of the solution set of a system of 3 linear equations in 3 variables if the system

- (a) is inconsistent.
- (b) is consistent and has no free variables.
- (c) is consistent and has exactly one free variable.
- (d) is consistent and has two free variables.

- (2) Consider the following system of equations:

$$\begin{cases} -m_1x + y = b_1 \\ -m_2x + y = b_2 \end{cases} \quad (*)$$

- (a) Prove that if  $m_1 \neq m_2$ , then  $(*)$  has exactly one solution. What is it?
- (b) Suppose that  $m_1 = m_2$ . Then under what conditions will  $(*)$  be consistent?
- (c) Restate the results of (a) and (b) in geometrical language.

**1.4. Answers to Odd-Numbered Exercises**

- (1) 2, 3
- (3) (a) 2, -1, 1  
(b) 3, -2, 1  
(c) 3, -2, 1
- (5) (a)  $k = -1$   
(b)  $k \neq -1, 1$   
(c)  $k = 1$   
(d)  $\frac{1}{k+1}, \frac{1}{k+1}$
- (7) (a) -6  
(b) a line  
(c) They have no points in common.
- (9) (a) 4  
(b) 40, -10  
(c) 10, 20



## CHAPTER 2

# ARITHMETIC OF MATRICES

### 2.1. Background

**Topics:** addition, scalar multiplication, and multiplication of matrices, inverse of a nonsingular matrix.

**2.1.1. Definition.** Two square matrices  $A$  and  $B$  of the same size are said to COMMUTE if  $AB = BA$ .

**2.1.2. Definition.** If  $A$  and  $B$  are square matrices of the same size, then the COMMUTATOR (or LIE BRACKET) of  $A$  and  $B$ , denoted by  $[A, B]$ , is defined by

$$[A, B] = AB - BA.$$

**2.1.3. Notation.** If  $A$  is an  $m \times n$  matrix (that is, a matrix with  $m$  rows and  $n$  columns), then the element in the  $i^{\text{th}}$  row and the  $j^{\text{th}}$  column is denoted by  $a_{ij}$ . The matrix  $A$  itself may be denoted by  $[a_{ij}]_{i=1}^m \substack{m \\ n} \substack{n \\ j=1}$  or, more simply, by  $[a_{ij}]$ . In light of this notation it is reasonable to refer to the index  $i$  in the expression  $a_{ij}$  as the ROW INDEX and to call  $j$  the COLUMN INDEX. When we speak of the “value of a matrix  $A$  at  $(i, j)$ ,” we mean the entry in the  $i^{\text{th}}$  row and  $j^{\text{th}}$  column of  $A$ . Thus, for example,

$$A = \begin{bmatrix} 1 & 4 \\ 3 & -2 \\ 7 & 0 \\ 5 & -1 \end{bmatrix}$$

is a  $4 \times 2$  matrix and  $a_{31} = 7$ .

**2.1.4. Definition.** A matrix  $A = [a_{ij}]$  is UPPER TRIANGULAR if  $a_{ij} = 0$  whenever  $i > j$ .

**2.1.5. Definition.** The TRACE of a square matrix  $A$ , denoted by  $\text{tr } A$ , is the sum of the diagonal entries of the matrix. That is, if  $A = [a_{ij}]$  is an  $n \times n$  matrix, then

$$\text{tr } A := \sum_{j=1}^n a_{jj}.$$

**2.1.6. Definition.** The TRANSPOSE of an  $n \times n$  matrix  $A = [a_{ij}]$  is the matrix  $A^t = [a_{ji}]$  obtained by interchanging the rows and columns of  $A$ . The matrix  $A$  is SYMMETRIC if  $A^t = A$ .

**2.1.7. Proposition.** If  $A$  is an  $m \times n$  matrix and  $B$  is an  $n \times p$  matrix, then  $(AB)^t = B^t A^t$ .

## 2.2. Exercises

(1) Let  $A = \begin{bmatrix} 1 & 0 & -1 & 2 \\ 0 & 3 & 1 & -1 \\ 2 & 4 & 0 & 3 \\ -3 & 1 & -1 & 2 \end{bmatrix}$ ,  $B = \begin{bmatrix} 1 & 2 \\ 3 & -1 \\ 0 & -2 \\ 4 & 1 \end{bmatrix}$ , and  $C = \begin{bmatrix} 3 & -2 & 0 & 5 \\ 1 & 0 & -3 & 4 \end{bmatrix}$ .

- (a) Does the matrix  $D = ABC$  exist? \_\_\_\_\_ If so, then  $d_{34} =$  \_\_\_\_\_.  
 (b) Does the matrix  $E = BAC$  exist? \_\_\_\_\_ If so, then  $e_{22} =$  \_\_\_\_\_.  
 (c) Does the matrix  $F = BCA$  exist? \_\_\_\_\_ If so, then  $f_{43} =$  \_\_\_\_\_.  
 (d) Does the matrix  $G = ACB$  exist? \_\_\_\_\_ If so, then  $g_{31} =$  \_\_\_\_\_.  
 (e) Does the matrix  $H = CAB$  exist? \_\_\_\_\_ If so, then  $h_{21} =$  \_\_\_\_\_.  
 (f) Does the matrix  $J = CBA$  exist? \_\_\_\_\_ If so, then  $j_{13} =$  \_\_\_\_\_.

(2) Let  $A = \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{bmatrix}$ ,  $B = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$ , and  $C = AB$ . Evaluate the following.

(a)  $A^{37} = \begin{bmatrix} & \\ & \end{bmatrix}$  (b)  $B^{63} = \begin{bmatrix} & \\ & \end{bmatrix}$   
 (c)  $B^{138} = \begin{bmatrix} & \\ & \end{bmatrix}$  (d)  $C^{42} = \begin{bmatrix} & \\ & \end{bmatrix}$

Note: If  $M$  is a matrix  $M^p$  is the product of  $p$  copies of  $M$ .

(3) Let  $A = \begin{bmatrix} 1 & 1/3 \\ c & d \end{bmatrix}$ . Find numbers  $c$  and  $d$  such that  $A^2 = -I$ .

Answer:  $c =$  \_\_\_\_\_ and  $d =$  \_\_\_\_\_.

(4) Let  $A$  and  $B$  be symmetric  $n \times n$ -matrices. Then  $[A, B] = [B, X]$ , where  $X =$  \_\_\_\_\_.

(5) Let  $A$ ,  $B$ , and  $C$  be  $n \times n$  matrices. Then  $[A, B]C + B[A, C] = [X, Y]$ , where  $X =$  \_\_\_\_\_ and  $Y =$  \_\_\_\_\_.

(6) Let  $A = \begin{bmatrix} 1 & 1/3 \\ c & d \end{bmatrix}$ . Find numbers  $c$  and  $d$  such that  $A^2 = 0$ . Answer:  $c =$  \_\_\_\_\_ and  $d =$  \_\_\_\_\_.

(7) Consider the matrix  $\begin{bmatrix} 1 & 3 & 2 \\ a & 6 & 2 \\ 0 & 9 & 5 \end{bmatrix}$  where  $a$  is a real number.

- (a) For what value of  $a$  will a row interchange be required during Gaussian elimination?

Answer:  $a =$  \_\_\_\_\_.

- (b) For what value of  $a$  is the matrix singular? Answer:  $a =$  \_\_\_\_\_.

(8) Let  $A = \begin{bmatrix} 1 & 0 & -1 & 2 \\ 0 & 3 & 1 & -1 \\ 2 & 4 & 0 & 3 \\ -3 & 1 & -1 & 2 \end{bmatrix}$ ,  $B = \begin{bmatrix} 1 & 2 \\ 3 & -1 \\ 0 & -2 \\ 4 & 1 \end{bmatrix}$ ,  $C = \begin{bmatrix} 3 & -2 & 0 & 5 \\ 1 & 0 & -3 & 4 \end{bmatrix}$ , and

$M = 3A^3 - 5(BC)^2$ . Then  $m_{14} =$  \_\_\_\_\_ and  $m_{41} =$  \_\_\_\_\_.

(9) If  $A$  is an  $n \times n$  matrix and it satisfies the equation  $A^3 - 4A^2 + 3A - 5I_n = 0$ , then  $A$  is nonsingular

and its inverse is \_\_\_\_\_ .

- (10) Let  $A$ ,  $B$ , and  $C$  be  $n \times n$  matrices. Then  $[[A, B], C] + [[B, C], A] + [[C, A], B] = X$ , where

$$X = \begin{bmatrix} & & & \\ & & & \\ & & & \\ & & & \end{bmatrix}.$$

- (11) Let  $A$ ,  $B$ , and  $C$  be  $n \times n$  matrices. Then  $[A, C] + [B, C] = [X, Y]$ , where  $X =$  \_\_\_\_\_ and  $Y =$  \_\_\_\_\_ .

- (12) Find the inverse of  $\begin{bmatrix} 1 & 0 & 0 & 0 \\ \frac{1}{4} & 1 & 0 & 0 \\ \frac{1}{3} & \frac{1}{3} & 1 & 0 \\ \frac{1}{2} & \frac{1}{2} & \frac{1}{2} & 1 \end{bmatrix}$ . Answer:  $\begin{bmatrix} & & & \\ & & & \\ & & & \\ & & & \end{bmatrix}$ .

- (13) The matrix

$$H = \begin{bmatrix} 1 & \frac{1}{2} & \frac{1}{3} & \frac{1}{4} \\ \frac{1}{2} & \frac{1}{3} & \frac{1}{4} & \frac{1}{5} \\ \frac{1}{3} & \frac{1}{4} & \frac{1}{5} & \frac{1}{6} \\ \frac{1}{4} & \frac{1}{5} & \frac{1}{6} & \frac{1}{7} \end{bmatrix}$$

is the  $4 \times 4$  HILBERT MATRIX. Use Gauss-Jordan elimination to compute  $K = H^{-1}$ . Then  $K_{44}$  is (exactly) \_\_\_\_\_. Now, create a new matrix  $H'$  by replacing each entry in  $H$  by its approximation to 3 decimal places. (For example, replace  $\frac{1}{6}$  by 0.167.) Use Gauss-Jordan elimination again to find the inverse  $K'$  of  $H'$ . Then  $K'_{44}$  is \_\_\_\_\_ .

- (14) Suppose that  $A$  and  $B$  are symmetric  $n \times n$  matrices. In this exercise we prove that  $AB$  is symmetric if and only if  $A$  commutes with  $B$ . Below are portions of the proof. Fill in the missing steps and the missing reasons. Choose reasons from the following list.

- (H1) Hypothesis that  $A$  and  $B$  are symmetric.
- (H2) Hypothesis that  $AB$  is symmetric.
- (H3) Hypothesis that  $A$  commutes with  $B$ .
- (D1) Definition of *commutes*.
- (D2) Definition of *symmetric*.
- (T) Proposition 2.1.7.

PROOF. Suppose that  $AB$  is symmetric. Then

$$\begin{aligned} AB &= \text{_____} \quad (\text{reason: (H2) and _____}) \\ &= B^t A^t \quad (\text{reason: _____}) \\ &= \text{_____} \quad (\text{reason: (D2) and _____}) \end{aligned}$$

So  $A$  commutes with  $B$  (reason: \_\_\_\_\_).

Conversely, suppose that  $A$  commutes with  $B$ . Then

$$\begin{aligned} (AB)^t &= \text{_____} \quad (\text{reason: (T)}) \\ &= BA \quad (\text{reason: _____ and _____}) \\ &= \text{_____} \quad (\text{reason: _____ and _____}) \end{aligned}$$

Thus  $AB$  is symmetric (reason: \_\_\_\_\_). □

### 2.3. Problems

- (1) Let  $A$  be a square matrix. Prove that if  $A^2$  is invertible, then so is  $A$ .

*Hint.* Our assumption is that there exists a matrix  $B$  such that

$$A^2B = BA^2 = I.$$

We want to show that there exists a matrix  $C$  such that

$$AC = CA = I.$$

Now to start with, you ought to find it fairly easy to show that there are matrices  $L$  and  $R$  such that

$$LA = AR = I. \quad (*)$$

A matrix  $L$  is a LEFT INVERSE of the matrix  $A$  if  $LA = I$ ; and  $R$  is a RIGHT INVERSE of  $A$  if  $AR = I$ . Thus the problem boils down to determining whether  $A$  can have a left inverse and a right inverse that are *different*. (Clearly, if it turns out that they must be the same, then the  $C$  we are seeking is their common value.) So try to prove that if  $(*)$  holds, then  $L = R$ .

- (2) Anton speaks French and German; Geraldine speaks English, French and Italian; James speaks English, Italian, and Spanish; Lauren speaks all the languages the others speak except French; and no one speaks any other language. Make a matrix  $A = [a_{ij}]$  with rows representing the four people mentioned and columns representing the languages they speak. Put  $a_{ij} = 1$  if person  $i$  speaks language  $j$  and  $a_{ij} = 0$  otherwise. Explain the significance of the matrices  $AA^t$  and  $A^tA$ .

- (3) Portland Fast Foods (PFF), which produces 138 food products all made from 87 basic ingredients, wants to set up a simple data structure from which they can quickly extract answers to the following questions:

- (a) How many ingredients does a given product contain?
- (b) A given pair of ingredients are used together in how many products?
- (c) How many ingredients do two given products have in common?
- (d) In how many products is a given ingredient used?

In particular, PFF wants to set up a single table in such a way that:

- (i) the answer to any of the above questions can be extracted easily and quickly (matrix arithmetic permitted, of course); and
- (ii) if one of the 87 ingredients is added to or deleted from a product, only a single entry in the table needs to be changed.

Is this possible? Explain.

- (4) Prove proposition 2.1.7.

- (5) Let  $A$  and  $B$  be  $2 \times 2$  matrices.

- (a) Prove that if the trace of  $A$  is 0, then  $A^2$  is a scalar multiple of the identity matrix.
- (b) Prove that the square of the commutator of  $A$  and  $B$  commutes with every  $2 \times 2$  matrix  $C$ . *Hint.* What can you say about the trace of  $[A, B]$ ?
- (c) Prove that the commutator of  $A$  and  $B$  can never be a nonzero multiple of the identity matrix.

- (6) The matrices that represent rotations of the  $xy$ -plane are

$$A(\theta) = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}.$$

- Let  $\mathbf{x}$  be the vector  $(-1, 1)$ ,  $\theta = 3\pi/4$ , and  $\mathbf{y}$  be  $A(\theta)$  acting on  $\mathbf{x}$  (that is,  $\mathbf{y} = A(\theta)\mathbf{x}$ ). Make a sketch showing  $\mathbf{x}$ ,  $\mathbf{y}$ , and  $\theta$ .
- Verify that  $A(\theta_1)A(\theta_2) = A(\theta_1 + \theta_2)$ . Discuss what this means geometrically.
- What is the product of  $A(\theta)$  times  $A(-\theta)$ ? Discuss what this means geometrically.
- Two sheets of graph paper are attached at the origin and rotated in such a way that the point  $(1, 0)$  on the upper sheet lies directly over the point  $(-5/13, 12/13)$  on the lower sheet. What point on the lower sheet lies directly below  $(6, 4)$  on the upper one?

- (7) Let

$$A = \begin{bmatrix} 0 & a & a^2 & a^3 & a^4 \\ 0 & 0 & a & a^2 & a^3 \\ 0 & 0 & 0 & a & a^2 \\ 0 & 0 & 0 & 0 & a \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

The goal of this problem is to develop a “calculus” for the matrix  $A$ . To start, recall (or look up) the power series expansion for  $\frac{1}{1-x}$ . Now see if this formula works for the matrix  $A$  by first computing  $(I - A)^{-1}$  directly and then computing the power series expansion substituting  $A$  for  $x$ . (Explain why there are no convergence difficulties for the series when we use this particular matrix  $A$ .) Next try to define  $\ln(I + A)$  and  $e^A$  by means of appropriate series. Do you get what you expect when you compute  $e^{\ln(I+A)}$ ? Do formulas like  $e^A e^A = e^{2A}$  hold? What about other familiar properties of the exponential and logarithmic functions?

Try some trigonometry with  $A$ . Use series to define  $\sin$ ,  $\cos$ ,  $\tan$ ,  $\arctan$ , and so on. Do things like  $\tan(\arctan(A))$  produce the expected results? Check some of the more obvious trigonometric identities. (What do you get for  $\sin^2 A + \cos^2 A - I$ ? Is  $\cos(2A)$  the same as  $\cos^2 A - \sin^2 A$ ?)

A relationship between the exponential and trigonometric functions is given by the famous formula  $e^{ix} = \cos x + i \sin x$ . Does this hold for  $A$ ?

Do you think there are other matrices for which the same results might hold? Which ones?

- (8) (a) Give an example of two symmetric matrices whose product is not symmetric.  
*Hint.* Matrices containing only 0's and 1's will suffice.
- (b) Now suppose that  $A$  and  $B$  are symmetric  $n \times n$  matrices. Prove that  $AB$  is symmetric if and only if  $A$  commutes with  $B$ .

*Hint.* To prove that a statement  $P$  holds “if and only if” a statement  $Q$  holds you must first show that  $P$  implies  $Q$  and then show that  $Q$  implies  $P$ . In the current problem, there are 4 conditions to be considered:

- $A^t = A$  ( $A$  is symmetric),
- $B^t = B$  ( $B$  is symmetric),
- $(AB)^t = AB$  ( $AB$  is symmetric), and
- $AB = BA$  ( $A$  commutes with  $B$ ).

Recall also the fact given in

- (v) theorem 2.1.7.

The first task is to derive (iv) from (i), (ii), (iii), and (v). Then try to derive (iii) from (i), (ii), (iv), and (v).

**2.4. Answers to Odd-Numbered Exercises**

- (1) (a) yes, 142  
(b) no,  $-$   
(c) yes,  $-45$   
(d) no,  $-$   
(e) yes,  $-37$   
(f) no,  $-$
- (3)  $-6, -1$
- (5)  $A, BC$
- (7) (a) 2  
(b)  $-4$
- (9)  $\frac{1}{5}(A^2 - 4A + 3I_n)$
- (11)  $A + B, C$
- (13) 2800,  $-1329.909$

## CHAPTER 3

# ELEMENTARY MATRICES; DETERMINANTS

### 3.1. Background

**Topics:** elementary (reduction) matrices, determinants.

The following definition says that we often regard the effect of multiplying a matrix  $M$  on the left by another matrix  $A$  as the *action of  $A$  on  $M$* .

**3.1.1. Definition.** We say that the matrix  $A$  ACTS ON the matrix  $M$  to produce the matrix  $N$  if  $N = AM$ . For example the matrix  $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$  acts on any  $2 \times 2$  matrix by interchanging (swapping) its rows because  $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} c & d \\ a & b \end{bmatrix}$ .

**3.1.2. Notation.** We adopt the following notation for elementary matrices which implement type I row operations. Let  $A$  be a matrix having  $n$  rows. For any real number  $r \neq 0$  denote by  $M_j(r)$  the  $n \times n$  matrix which acts on  $A$  by multiplying its  $j^{\text{th}}$  row by  $r$ . (See exercise 1.)

**3.1.3. Notation.** We use the following notation for elementary matrices which implement type II row operations. (See definition 1.1.1.) Let  $A$  be a matrix having  $n$  rows. Denote by  $P_{ij}$  the  $n \times n$  matrix which acts on  $A$  by interchanging its  $i^{\text{th}}$  and  $j^{\text{th}}$  rows. (See exercise 2.)

**3.1.4. Notation.** And we use the following notation for elementary matrices which implement type III row operations. (See definition 1.1.1.) Let  $A$  be a matrix having  $n$  rows. For any real number  $r$  denote by  $E_{ij}(r)$  the  $n \times n$  matrix which acts on  $A$  by adding  $r$  times the  $j^{\text{th}}$  row of  $A$  to the  $i^{\text{th}}$  row. (See exercise 3.)

**3.1.5. Definition.** If a matrix  $B$  can be produced from a matrix  $A$  by a sequence of elementary row operations, then  $A$  and  $B$  are ROW EQUIVALENT.

### Some Facts about Determinants

**3.1.6. Proposition.** Let  $n \in \mathbb{N}$  and  $\mathbf{M}_{n \times n}$  be the collection of all  $n \times n$  matrices. There is exactly one function

$$\det: \mathbf{M}_{n \times n} \rightarrow \mathbb{R}: A \mapsto \det A$$

which satisfies

- (a)  $\det I_n = 1$ .
- (b) If  $A \in \mathbf{M}_{n \times n}$  and  $A'$  is the matrix obtained by interchanging two rows of  $A$ , then  $\det A' = -\det A$ .
- (c) If  $A \in \mathbf{M}_{n \times n}$ ,  $c \in \mathbb{R}$ , and  $A'$  is the matrix obtained by multiplying each element in one row of  $A$  by the number  $c$ , then  $\det A' = c \det A$ .
- (d) If  $A \in \mathbf{M}_{n \times n}$ ,  $c \in \mathbb{R}$ , and  $A'$  is the matrix obtained from  $A$  by multiplying one row of  $A$  by  $c$  and adding it to another row of  $A$  (that is, choose  $i$  and  $j$  between 1 and  $n$  with  $i \neq j$  and replace  $a_{jk}$  by  $a_{jk} + ca_{ik}$  for  $1 \leq k \leq n$ ), then  $\det A' = \det A$ .

**3.1.7. Definition.** The unique function  $\det: \mathbf{M}_{n \times n} \rightarrow \mathbb{R}$  described above is the  $n \times n$  DETERMINANT FUNCTION.

**3.1.8. Proposition.** If  $A = [a]$  for  $a \in \mathbb{R}$  (that is, if  $A \in \mathbf{M}_{1 \times 1}$ ), then  $\det A = a$ ; if  $A \in \mathbf{M}_{2 \times 2}$ , then

$$\det A = a_{11}a_{22} - a_{12}a_{21}.$$

**3.1.9. Proposition.** If  $A, B \in \mathbf{M}_{n \times n}$ , then  $\det(AB) = (\det A)(\det B)$ .

**3.1.10. Proposition.** If  $A \in \mathbf{M}_{n \times n}$ , then  $\det A^t = \det A$ . (An obvious corollary of this: in conditions (b), (c), and (d) of proposition 3.1.6 the word “columns” may be substituted for the word “rows”.)

**3.1.11. Definition.** Let  $A$  be an  $n \times n$  matrix. The MINOR of the element  $a_{jk}$ , denoted by  $M_{jk}$ , is the determinant of the  $(n-1) \times (n-1)$  matrix which results from the deletion of the  $j^{\text{th}}$  row and  $k^{\text{th}}$  column of  $A$ . The COFACTOR of the element  $a_{jk}$ , denoted by  $C_{jk}$  is defined by

$$C_{jk} := (-1)^{j+k} M_{jk}.$$

**3.1.12. Proposition.** If  $A \in \mathbf{M}_{n \times n}$  and  $1 \leq j \leq n$ , then

$$\det A = \sum_{k=1}^n a_{jk} C_{jk}.$$

This is the (LAPLACE) EXPANSION of the determinant along the  $j^{\text{th}}$  row.

In light of 3.1.10, it is clear that expansion along columns works as well as expansion along rows. That is,

$$\det A = \sum_{j=1}^n a_{jk} C_{jk}$$

for any  $k$  between 1 and  $n$ . This is the (LAPLACE) EXPANSION of the determinant along the  $k^{\text{th}}$  column.

**3.1.13. Proposition.** An  $n \times n$  matrix  $A$  is invertible if and only if  $\det A \neq 0$ . If  $A$  is invertible, then

$$A^{-1} = (\det A)^{-1} C^t$$

where  $C = [C_{jk}]$  is the matrix of cofactors of elements of  $A$ .



**3.2. Exercises**

- (1) Let  $A$  be a matrix with 4 rows. The matrix  $M_3(4)$  which multiplies the 3<sup>rd</sup> row of  $A$  by 4

is  $\begin{bmatrix} & & & \\ & & & \\ & & & \\ & & & \end{bmatrix}$ . (See 3.1.2.)

- (2) Let  $A$  be a matrix with 4 rows. The matrix  $P_{24}$  which interchanges the 2<sup>nd</sup> and 4<sup>th</sup> rows

of  $A$  is  $\begin{bmatrix} & & & \\ & & & \\ & & & \\ & & & \end{bmatrix}$ . (See 3.1.3.)

- (3) Let  $A$  be a matrix with 4 rows. The matrix  $E_{23}(-2)$  which adds  $-2$  times the 3<sup>rd</sup> row of

$A$  to the 2<sup>nd</sup> row is  $\begin{bmatrix} & & & \\ & & & \\ & & & \\ & & & \end{bmatrix}$ . (See 3.1.4.)

- (4) Let  $A$  be the  $4 \times 4$  elementary matrix  $E_{43}(-6)$ . Then  $A^{11} = \begin{bmatrix} & & & \\ & & & \\ & & & \\ & & & \end{bmatrix}$  and

$A^{-9} = \begin{bmatrix} & & & \\ & & & \\ & & & \\ & & & \end{bmatrix}$ .

- (5) Let  $B$  be the elementary  $4 \times 4$  matrix  $P_{24}$ . Then  $B^{-9} = \begin{bmatrix} & & & \\ & & & \\ & & & \\ & & & \end{bmatrix}$  and

$B^{10} = \begin{bmatrix} & & & \\ & & & \\ & & & \\ & & & \end{bmatrix}$ .

- (6) Let  $C$  be the elementary  $4 \times 4$  matrix  $M_3(-2)$ . Then  $C^4 = \begin{bmatrix} & & & \\ & & & \\ & & & \\ & & & \end{bmatrix}$  and

$C^{-3} = \begin{bmatrix} & & & \\ & & & \\ & & & \\ & & & \end{bmatrix}$ .

- (7) Let  $A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & -1 & 1 \\ -2 & 1 & 0 \\ -1 & 2 & -3 \end{bmatrix}$  and  $B = P_{23}E_{34}(-2)M_3(-2)E_{42}(1)P_{14}A$ . Then  $b_{23} = \underline{\hspace{2cm}}$   
and  $b_{32} = \underline{\hspace{2cm}}$ .

- (8) We apply Gaussian elimination (using type III elementary row operations only) to put a  $4 \times 4$  matrix  $A$  into upper triangular form. The result is

$$E_{43}\left(\frac{5}{2}\right)E_{42}(2)E_{31}(1)E_{21}(-2)A = \begin{bmatrix} 1 & 2 & -2 & 0 \\ 0 & -1 & 0 & 1 \\ 0 & 0 & -2 & 2 \\ 0 & 0 & 0 & 10 \end{bmatrix}.$$

Then the determinant of  $A$  is  $\underline{\hspace{2cm}}$ .

(9) The system of equations:

$$\begin{cases} 2y+3z = 7 \\ x+ y- z = -2 \\ -x+ y-5z = 0 \end{cases}$$

is solved by applying Gauss-Jordan reduction to the augmented coefficient matrix

$$A = \begin{bmatrix} 0 & 2 & 3 & 7 \\ 1 & 1 & -1 & -2 \\ -1 & 1 & -5 & 0 \end{bmatrix}. \text{ Give the names of the elementary } 3 \times 3 \text{ matrices } X_1, \dots, X_8$$

which implement the following reduction.

$$\begin{aligned} A &\xrightarrow{X_1} \begin{bmatrix} 1 & 1 & -1 & -2 \\ 0 & 2 & 3 & 7 \\ -1 & 1 & -5 & 0 \end{bmatrix} \xrightarrow{X_2} \begin{bmatrix} 1 & 1 & -1 & -2 \\ 0 & 2 & 3 & 7 \\ 0 & 2 & -6 & -2 \end{bmatrix} \xrightarrow{X_3} \begin{bmatrix} 1 & 1 & -1 & -2 \\ 0 & 2 & 3 & 7 \\ 0 & 0 & -9 & -9 \end{bmatrix} \\ &\xrightarrow{X_4} \begin{bmatrix} 1 & 1 & -1 & -2 \\ 0 & 2 & 3 & 7 \\ 0 & 0 & 1 & 1 \end{bmatrix} \xrightarrow{X_5} \begin{bmatrix} 1 & 1 & -1 & -2 \\ 0 & 2 & 0 & 4 \\ 0 & 0 & 1 & 1 \end{bmatrix} \xrightarrow{X_6} \begin{bmatrix} 1 & 1 & -1 & -2 \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 1 & 1 \end{bmatrix} \\ &\xrightarrow{X_7} \begin{bmatrix} 1 & 1 & 0 & -1 \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 1 & 1 \end{bmatrix} \xrightarrow{X_8} \begin{bmatrix} 1 & 0 & 0 & -3 \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 1 & 1 \end{bmatrix}. \end{aligned}$$

Answer:  $X_1 = \underline{\hspace{2cm}}$ ,  $X_2 = \underline{\hspace{2cm}}$ ,  $X_3 = \underline{\hspace{2cm}}$ ,  $X_4 = \underline{\hspace{2cm}}$ ,

$X_5 = \underline{\hspace{2cm}}$ ,  $X_6 = \underline{\hspace{2cm}}$ ,  $X_7 = \underline{\hspace{2cm}}$ ,  $X_8 = \underline{\hspace{2cm}}$ .

(10) Solve the following equation for  $x$ :

$$\det \begin{bmatrix} 3 & -4 & 7 & 0 & 6 & -2 \\ 2 & 0 & 1 & 8 & 0 & 0 \\ 3 & 4 & -8 & 3 & 1 & 2 \\ 27 & 6 & 5 & 0 & 0 & 3 \\ 3 & x & 0 & 2 & 1 & -1 \\ 1 & 0 & -1 & 3 & 4 & 0 \end{bmatrix} = 0. \quad \text{Answer: } x = \underline{\hspace{2cm}}.$$

(11) Let  $A = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 2 & 4 \\ 1 & 2 & 3 \end{bmatrix}$ . Find  $A^{-1}$  using the technique of augmenting  $A$  by the identity matrix

$I$  and performing Gauss-Jordan reduction on the augmented matrix. The reduction can be accomplished by the application of five elementary  $3 \times 3$  matrices. Find elementary matrices  $X_1$ ,  $X_2$ , and  $X_3$  such that  $A^{-1} = X_3 E_{13}(-3) X_2 M_2(1/2) X_1 I$ .

(a) The required matrices are  $X_1 = P_{1i}$  where  $i = \underline{\hspace{1cm}}$ ,  $X_2 = E_{jk}(-2)$  where  $j = \underline{\hspace{1cm}}$  and  $k = \underline{\hspace{1cm}}$ , and  $X_3 = E_{12}(r)$  where  $r = \underline{\hspace{1cm}}$ .

(b) And then  $A^{-1} = \begin{bmatrix} \phantom{0} & \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} & \phantom{0} \end{bmatrix}$ .

$$(12) \det \begin{bmatrix} 1 & t & t^2 & t^3 \\ t & 1 & t & t^2 \\ t^2 & t & 1 & t \\ t^3 & t^2 & t & 1 \end{bmatrix} = (1 - a(t))^p \text{ where } a(t) = \underline{\hspace{2cm}} \text{ and } p = \underline{\hspace{2cm}}.$$

(13) Evaluate each of the following determinants.

$$(a) \det \begin{bmatrix} 6 & 9 & 39 & 49 \\ 5 & 7 & 32 & 37 \\ 3 & 4 & 4 & 4 \\ 1 & 1 & 1 & 1 \end{bmatrix} = \underline{\hspace{2cm}} .$$

$$(b) \det \begin{bmatrix} 1 & 0 & 1 & 1 \\ 1 & -1 & 2 & 0 \\ 2 & -1 & 3 & 1 \\ 4 & 17 & 0 & -5 \end{bmatrix} = \underline{\hspace{2cm}} .$$

$$(c) \det \begin{bmatrix} 13 & 3 & -8 & 6 \\ 0 & 0 & -4 & 0 \\ 1 & 0 & 7 & -2 \\ 3 & 0 & 2 & 0 \end{bmatrix} = \underline{\hspace{2cm}} .$$

$$(14) \text{ Let } M \text{ be the matrix } \begin{bmatrix} 5 & 4 & -2 & 3 \\ 5 & 7 & -1 & 8 \\ 5 & 7 & 6 & 10 \\ 5 & 7 & 1 & 9 \end{bmatrix} .$$

(a) The determinant of  $M$  can be expressed as the constant 5 times the determinant of the single  $3 \times 3$  matrix  $\begin{bmatrix} 3 & 1 & 5 \\ 3 & & \\ 3 & & \end{bmatrix}$ .

(b) The determinant of this  $3 \times 3$  matrix can be expressed as the constant 3 times the determinant of the single  $2 \times 2$  matrix  $\begin{bmatrix} 7 & 2 \\ 2 & \end{bmatrix}$ .

(c) The determinant of this  $2 \times 2$  matrix is  $\underline{\hspace{2cm}}$ .

(d) Thus the determinant of  $M$  is  $\underline{\hspace{2cm}}$ .

$$(15) \text{ Find the determinant of the matrix } \begin{bmatrix} 1 & 2 & 5 & 7 & 10 \\ 1 & 2 & 3 & 6 & 7 \\ 1 & 1 & 3 & 5 & 5 \\ 1 & 1 & 2 & 4 & 5 \\ 1 & 1 & 1 & 1 & 1 \end{bmatrix} . \text{ Answer: } \underline{\hspace{2cm}} .$$

(16) Find the determinants of the following matrices.

$$A = \begin{bmatrix} -73 & 78 & 24 \\ 92 & 66 & 25 \\ -80 & 37 & 10 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} -73 & 78 & 24 \\ 92 & 66 & 25 \\ -80 & 37 & 10.01 \end{bmatrix} .$$

*Hint.* Use a calculator (thoughtfully). Answer:  $\det A = \underline{\hspace{2cm}}$  and  $\det B = \underline{\hspace{2cm}}$ .

(17) Find the determinant of the following matrix.

$$\begin{bmatrix} 283 & 5 & \pi & 347.86 \times 10^{15^{83}} \\ 3136 & 56 & 5 & \cos(2.7402) \\ 6776 & 121 & 11 & 5 \\ 2464 & 44 & 4 & 2 \end{bmatrix} .$$

*Hint.* Do not use a calculator. Answer:  $\underline{\hspace{2cm}}$ .

(18) Let  $A = \begin{bmatrix} 0 & -\frac{1}{2} & 0 & \frac{1}{2} \\ 0 & 0 & \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & 0 & -\frac{1}{2} & 0 \\ 1 & 0 & \frac{1}{2} & \frac{1}{2} \end{bmatrix}$ . We find  $A^{-1}$  using elementary row operations to convert the

$4 \times 8$  matrix  $\begin{bmatrix} A & : & I_4 \end{bmatrix}$  to the matrix  $\begin{bmatrix} I_4 & : & A^{-1} \end{bmatrix}$ .

Give the names of the elementary  $4 \times 4$  matrices  $X_1, \dots, X_{11}$  which implement the following Gauss-Jordan reduction and fill in the missing matrix entries.

$$\begin{aligned}
 & \begin{bmatrix} 0 & -\frac{1}{2} & 0 & \frac{1}{2} & : & 1 & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{2} & \frac{1}{2} & : & 0 & 1 & 0 & 0 \\ \frac{1}{2} & 0 & -\frac{1}{2} & 0 & : & 0 & 0 & 1 & 0 \\ 1 & 0 & \frac{1}{2} & \frac{1}{2} & : & 0 & 0 & 0 & 1 \end{bmatrix} \xrightarrow{X_1} \begin{bmatrix} 1 & 0 & \frac{1}{2} & \frac{1}{2} & : & & & & \\ 0 & 0 & \frac{1}{2} & \frac{1}{2} & : & & & & \\ \frac{1}{2} & 0 & -\frac{1}{2} & 0 & : & & & & \\ 0 & -\frac{1}{2} & 0 & \frac{1}{2} & : & & & & \end{bmatrix} \\
 & \xrightarrow{X_2} \begin{bmatrix} 1 & 0 & \frac{1}{2} & \frac{1}{2} & : & & & & \\ 0 & 0 & \frac{1}{2} & \frac{1}{2} & : & & & & \\ 0 & 0 & -\frac{3}{4} & -\frac{1}{4} & : & & & & \\ 0 & -\frac{1}{2} & 0 & \frac{1}{2} & : & & & & \end{bmatrix} \xrightarrow{X_3} \begin{bmatrix} 1 & 0 & \frac{1}{2} & \frac{1}{2} & : & & & & \\ 0 & -\frac{1}{2} & 0 & \frac{1}{2} & : & & & & \\ 0 & 0 & -\frac{3}{4} & -\frac{1}{4} & : & & & & \\ 0 & 0 & \frac{1}{2} & \frac{1}{2} & : & & & & \end{bmatrix} \\
 & \xrightarrow{X_4} \begin{bmatrix} 1 & 0 & \frac{1}{2} & \frac{1}{2} & : & & & & \\ 0 & 1 & 0 & -1 & : & & & & \\ 0 & 0 & -\frac{3}{4} & -\frac{1}{4} & : & & & & \\ 0 & 0 & \frac{1}{2} & \frac{1}{2} & : & & & & \end{bmatrix} \xrightarrow{X_5} \begin{bmatrix} 1 & 0 & \frac{1}{2} & \frac{1}{2} & : & & & & \\ 0 & 1 & 0 & -1 & : & & & & \\ 0 & 0 & 1 & \frac{1}{3} & : & & & & \\ 0 & 0 & \frac{1}{2} & \frac{1}{2} & : & & & & \end{bmatrix} \\
 & \xrightarrow{X_6} \begin{bmatrix} 1 & 0 & \frac{1}{2} & \frac{1}{2} & : & & & & \\ 0 & 1 & 0 & -1 & : & & & & \\ 0 & 0 & 1 & \frac{1}{3} & : & & & & \\ 0 & 0 & 0 & \frac{1}{3} & : & & & & \end{bmatrix} \xrightarrow{X_7} \begin{bmatrix} 1 & 0 & \frac{1}{2} & \frac{1}{2} & : & & & & \\ 0 & 1 & 0 & -1 & : & & & & \\ 0 & 0 & 1 & 0 & : & & & & \\ 0 & 0 & 0 & \frac{1}{3} & : & & & & \end{bmatrix} \\
 & \xrightarrow{X_8} \begin{bmatrix} 1 & 0 & \frac{1}{2} & \frac{1}{2} & : & & & & \\ 0 & 1 & 0 & -1 & : & & & & \\ 0 & 0 & 1 & 0 & : & & & & \\ 0 & 0 & 0 & 1 & : & & & & \end{bmatrix} \xrightarrow{X_9} \begin{bmatrix} 1 & 0 & \frac{1}{2} & \frac{1}{2} & : & & & & \\ 0 & 1 & 0 & 0 & : & & & & \\ 0 & 0 & 1 & 0 & : & & & & \\ 0 & 0 & 0 & 1 & : & & & & \end{bmatrix} \\
 & \xrightarrow{X_{10}} \begin{bmatrix} 1 & 0 & \frac{1}{2} & 0 & : & & & & \\ 0 & 1 & 0 & 0 & : & & & & \\ 0 & 0 & 1 & 0 & : & & & & \\ 0 & 0 & 0 & 1 & : & & & & \end{bmatrix} \xrightarrow{X_{11}} \begin{bmatrix} 1 & 0 & 0 & 0 & : & & & & \\ 0 & 1 & 0 & 0 & : & & & & \\ 0 & 0 & 1 & 0 & : & & & & \\ 0 & 0 & 0 & 1 & : & & & & \end{bmatrix}
 \end{aligned}$$

Answer:  $X_1 =$  \_\_\_\_\_,  $X_2 =$  \_\_\_\_\_,  $X_3 =$  \_\_\_\_\_,  $X_4 =$  \_\_\_\_\_,

$X_5 =$  \_\_\_\_\_,  $X_6 =$  \_\_\_\_\_,  $X_7 =$  \_\_\_\_\_,  $X_8 =$  \_\_\_\_\_.

$X_9 =$  \_\_\_\_\_,  $X_{10} =$  \_\_\_\_\_,  $X_{11} =$  \_\_\_\_\_.

(19) Suppose that  $A$  is a square matrix with determinant 7. Then

(a)  $\det(P_{24}A) = \underline{\hspace{1cm}}$  .

(b)  $\det(E_{23}(-4)A) = \underline{\hspace{1cm}}$  .

(c)  $\det(M_3(2)A) = \underline{\hspace{1cm}}$  .

### 3.3. Problems

- (1) For this problem assume that we know the following: If  $X$  is an  $m \times m$  matrix, if  $Y$  is an  $m \times n$  matrix and if  $\mathbf{0}$  and  $\mathbf{I}$  are zero and identity matrices of appropriate sizes, then

$$\det \begin{bmatrix} X & Y \\ \mathbf{0} & \mathbf{I} \end{bmatrix} = \det X.$$

Let  $A$  be an  $m \times n$  matrix and  $B$  be an  $n \times m$  matrix. Prove carefully that

$$\det \begin{bmatrix} \mathbf{0} & A \\ -B & \mathbf{I} \end{bmatrix} = \det AB.$$

*Hint.* Consider the product  $\begin{bmatrix} \mathbf{0} & A \\ -B & \mathbf{I} \end{bmatrix} \begin{bmatrix} \mathbf{I} & \mathbf{0} \\ B & \mathbf{I} \end{bmatrix}$ .

- (2) Let  $A$  and  $B$  be  $n \times n$ -matrices. Your good friend Fred R. Dimm believes that

$$\det \begin{bmatrix} A & B \\ B & A \end{bmatrix} = \det(A + B) \det(A - B).$$

He offers the following argument to support this claim:

$$\begin{aligned} \det \begin{bmatrix} A & B \\ B & A \end{bmatrix} &= \det(A^2 - B^2) \\ &= \det[(A + B)(A - B)] \\ &= \det(A + B) \det(A - B). \end{aligned}$$

- (a) Comment (helpfully) on his “proof”. In particular, explain carefully why each of the three steps in his “proof” is correct or incorrect. (That is, provide a proof or a counterexample to each step.)
- (b) Is the result he is trying to prove actually true?

*Hint:* Consider the product  $\begin{bmatrix} I & B \\ 0 & A - B \end{bmatrix} \begin{bmatrix} A + B & 0 \\ 0 & I \end{bmatrix}$ .

- (3) Let  $x$  be a fixed real number which is not an integer multiple of  $\pi$ . For each natural number  $n$  let  $A_n = [a_{jk}]$  be the  $n \times n$ -matrix defined by

$$a_{jk} = \begin{cases} 0, & \text{for } |j - k| > 1 \\ 1, & \text{for } |j - k| = 1 \\ 2 \cos x, & \text{for } j = k. \end{cases}$$

Show that  $\det A_n = \frac{\sin(n+1)x}{\sin x}$ . *Hint.* For each integer  $n$  let  $D_n = \det A_n$  and prove that

$$D_{n+2} - 2D_{n+1} \cos x + D_n = 0.$$

(Use mathematical induction.)

**3.4. Answers to Odd-Numbered Exercises**

$$(1) \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 4 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$(3) \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & -2 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$(5) \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$(7) -8, -1$$

$$(9) P_{12}, E_{31}(1), E_{32}(-1), M_3(-\frac{1}{9}), E_{23}(-3), M_2(\frac{1}{2}), E_{13}(1), E_{12}(-1)$$

$$(11) (a) 3, 2, 3, -2$$

$$(b) \begin{bmatrix} 1 & -1 & 1 \\ -2 & \frac{1}{2} & 0 \\ 1 & 0 & 0 \end{bmatrix}$$

$$(13) 100, 0, -72$$

$$(15) -10$$

$$(17) 6$$

$$(19) (a) -7$$

$$(b) 7$$

$$(c) 14$$





## CHAPTER 4

# VECTOR GEOMETRY IN $\mathbb{R}^n$

### 4.1. Background

**Topics:** inner (dot) products, cross products, lines and planes in 3-space, norm of a vector, angle between vectors.

**4.1.1. Notation.** There are many more or less standard notations for the inner product (or dot product) of two vectors  $\mathbf{x}$  and  $\mathbf{y}$ . The two that we will use interchangeably in these exercises are  $\mathbf{x} \cdot \mathbf{y}$  and  $\langle \mathbf{x}, \mathbf{y} \rangle$ .

**4.1.2. Definition.** If  $\mathbf{x}$  is a vector in  $\mathbb{R}^n$ , then the **NORM** (or **LENGTH**) of  $\mathbf{x}$  is defined by

$$\|\mathbf{x}\| = \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle}.$$

**4.1.3. Definition.** Let  $\mathbf{x}$  and  $\mathbf{y}$  be nonzero vectors in  $\mathbb{R}^n$ . Then  $\angle(\mathbf{x}, \mathbf{y})$ , the **ANGLE** between  $\mathbf{x}$  and  $\mathbf{y}$ , is defined by

$$\angle(\mathbf{x}, \mathbf{y}) = \arccos \frac{\langle \mathbf{x}, \mathbf{y} \rangle}{\|\mathbf{x}\| \|\mathbf{y}\|}$$

**4.1.4. Theorem** (Cauchy-Schwarz inequality). *If  $\mathbf{x}$  and  $\mathbf{y}$  are vectors in  $\mathbb{R}^n$ , then*

$$|\langle \mathbf{x}, \mathbf{y} \rangle| \leq \|\mathbf{x}\| \|\mathbf{y}\|.$$

(We will often refer to this just as the *Schwarz inequality*.)

**4.1.5. Definition.** If  $\mathbf{x} = (x_1, x_2, x_3)$  and  $\mathbf{y} = (y_1, y_2, y_3)$  are vectors in  $\mathbb{R}^3$ , then their **CROSS PRODUCT**, denoted by  $\mathbf{x} \times \mathbf{y}$ , is the vector  $(x_2y_3 - x_3y_2, x_3y_1 - x_1y_3, x_1y_2 - x_2y_1)$ .

## 4.2. Exercises

- (1) The angle between the vectors  $(1, 0, -1, 3)$  and  $(1, \sqrt{3}, 3, -3)$  in  $\mathbb{R}^4$  is  $a\pi$  where  $a = \underline{\hspace{2cm}}$ .
- (2) Find the angle  $\theta$  between the vectors  $\mathbf{x} = (3, -1, 1, 0, 2, 1)$  and  $\mathbf{y} = (2, -1, 0, \sqrt{2}, 2, 1)$  in  $\mathbb{R}^6$ . Answer:  $\theta = \underline{\hspace{2cm}}$ .

- (3) If  $a_1, \dots, a_n > 0$ , then

$$\left(\sum_{j=1}^n a_j\right) \left(\sum_{k=1}^n \frac{1}{a_k}\right) \geq n^2.$$

The proof of this is obvious from the *Cauchy-Schwarz inequality* when we choose the vectors  $\mathbf{x}$  and  $\mathbf{y}$  as follows:

$$\mathbf{x} = \underline{\hspace{2cm}} \text{ and } \mathbf{y} = \underline{\hspace{2cm}}.$$

- (4) Find all real numbers  $\alpha$  such that the angle between the vectors  $2\mathbf{i} + 2\mathbf{j} + (\alpha - 2)\mathbf{k}$  and  $2\mathbf{i} + (\alpha - 2)\mathbf{j} + 2\mathbf{k}$  is  $\frac{\pi}{3}$ . Answer:  $\alpha = \underline{\hspace{1cm}}$  and  $\underline{\hspace{1cm}}$ .
- (5) Which of the angles (if any) of triangle  $ABC$ , with  $A = (1, -2, 0)$ ,  $B = (2, 1, -2)$ , and  $C = (6, -1, -3)$ , is a right angle? Answer: the angle at vertex  $\underline{\hspace{1cm}}$ .
- (6) The hydrogen atoms of a methane molecule ( $\text{CH}_4$ ) are located at  $(0, 0, 0)$ ,  $(1, 1, 0)$ ,  $(0, 1, 1)$ , and  $(1, 0, 1)$  while the carbon atom is at  $(\frac{1}{2}, \frac{1}{2}, \frac{1}{2})$ . Find the cosine of the angle  $\theta$  between two rays starting at the carbon atom and going to different hydrogen atoms.

$$\text{Answer: } \cos \theta = \underline{\hspace{1cm}}.$$

- (7) If  $a, b, c, d, e, f \in \mathbb{R}$ , then

$$|ad + be + cf| \leq \sqrt{a^2 + b^2 + c^2} \sqrt{d^2 + e^2 + f^2}.$$

The proof of this inequality is obvious since this is just the *Cauchy-Schwarz inequality* where  $x = (\underline{\hspace{1cm}}, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})$  and  $y = (\underline{\hspace{1cm}}, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})$ .

- (8) The volume of the parallelepiped generated by the three vectors  $\mathbf{i} + 2\mathbf{j} - \mathbf{k}$ ,  $\mathbf{j} + \mathbf{k}$ , and  $3\mathbf{i} - \mathbf{j} + 2\mathbf{k}$  is  $\underline{\hspace{1cm}}$ .
- (9) The equations of the line containing the points  $(3, -1, 4)$  and  $(7, 9, 10)$  are

$$\frac{x-3}{2} = \frac{y-j}{b} = \frac{z-k}{c}$$

$$\text{where } b = \underline{\hspace{1cm}}, c = \underline{\hspace{1cm}}, j = \underline{\hspace{1cm}}, \text{ and } k = \underline{\hspace{1cm}}.$$

- (10) The equations of the line containing the points  $(5, 2, -1)$  and  $(9, -4, 1)$  are

$$\frac{x-h}{a} = \frac{y-2}{-3} = \frac{z-k}{c}$$

$$\text{where } a = \underline{\hspace{1cm}}, c = \underline{\hspace{1cm}}, h = \underline{\hspace{1cm}}, \text{ and } k = \underline{\hspace{1cm}}.$$

- (11) Find the equations of the line containing the point  $(1, 0, -1)$  which is parallel to the line  $\frac{x-4}{2} = \frac{2y-3}{5} = \frac{3z-7}{6}$ .

$$\text{Answer: } \frac{x-h}{a} = \frac{y-j}{b} = \frac{z+1}{4} \text{ where } a = \underline{\hspace{1cm}}, b = \underline{\hspace{1cm}}, h = \underline{\hspace{1cm}}, \text{ and } j = \underline{\hspace{1cm}}.$$

- (12) The equation of the plane containing the points  $(0, -1, 1)$ ,  $(1, 0, 2)$ , and  $(3, 0, 1)$  is  $x + by + cz = d$  where  $b = \underline{\hspace{1cm}}$ ,  $c = \underline{\hspace{1cm}}$ , and  $d = \underline{\hspace{1cm}}$ .
- (13) The equation of the plane which passes through the points  $(0, -1, -1)$ ,  $(5, 0, 1)$ , and  $(4, -1, 0)$  is  $ax + by + cz = 1$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ , and  $c = \underline{\hspace{1cm}}$ .
- (14) The angle between the planes  $4x + 4z - 16 = 0$  and  $-2x + 2y - 13 = 0$  is  $\frac{a}{b}\pi$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .

- (15) Suppose that  $\mathbf{u} \in \mathbb{R}^3$  is a vector which lies in the first quadrant of the  $xy$ -plane and has length 3 and that  $\mathbf{v} \in \mathbb{R}^3$  is a vector that lies along the positive  $z$ -axis and has length 5. Then
- (a)  $\|\mathbf{u} \times \mathbf{v}\| = \underline{\hspace{1cm}}$  ;
  - (b) the  $x$ -coordinate of  $\mathbf{u} \times \mathbf{v}$  is  $\underline{\hspace{1cm}}$  0 (choose  $<$ ,  $>$ , or  $=$ );
  - (c) the  $y$ -coordinate of  $\mathbf{u} \times \mathbf{v}$  is  $\underline{\hspace{1cm}}$  0 (choose  $<$ ,  $>$ , or  $=$ ); and
  - (d) the  $z$ -coordinate of  $\mathbf{u} \times \mathbf{v}$  is  $\underline{\hspace{1cm}}$  0 (choose  $<$ ,  $>$ , or  $=$ ).
- (16) Suppose that  $\mathbf{u}$  and  $\mathbf{v}$  are vectors in  $\mathbb{R}^7$  both of length  $2\sqrt{2}$  and that the length of  $\mathbf{u} - \mathbf{v}$  is also  $2\sqrt{2}$ . Then  $\|\mathbf{u} + \mathbf{v}\| = \underline{\hspace{1cm}}$  and the angle between  $\mathbf{u}$  and  $\mathbf{v}$  is  $\underline{\hspace{1cm}}$  .

## 4.3. Problems

- (1) Show that if  $a, b, c > 0$ , then  $(\frac{1}{2}a + \frac{1}{3}b + \frac{1}{6}c)^2 \leq \frac{1}{2}a^2 + \frac{1}{3}b^2 + \frac{1}{6}c^2$ .
- (2) Show that if  $a_1, \dots, a_n, w_1, \dots, w_n > 0$  and  $\sum_{k=1}^n w_k = 1$ , then

$$\left( \sum_{k=1}^n a_k w_k \right)^2 \leq \sum_{k=1}^n a_k^2 w_k.$$

- (3) Prove that if  $(a_1, a_2, \dots)$  is a sequence of real numbers such that the series  $\sum_{k=1}^{\infty} a_k^2$  converges, then the series  $\sum_{k=1}^{\infty} \frac{1}{k} a_k$  converges absolutely.

You may find the following steps helpful in organizing your solution.

- (i) First of all, make sure that you recall the difference between a sequence of numbers  $(c_1, c_2, \dots)$  and an infinite series  $\sum_{k=1}^{\infty} c_k$ .
- (ii) The key to this problem is an important theorem from third term Calculus:  
*A nondecreasing sequence of real numbers converges if and only if it is bounded.* (\*)  
 (Make sure that you know the meanings of all the terms used here.)
- (iii) The hypothesis of the result we are trying to prove is that the series  $\sum_{k=1}^{\infty} a_k^2$  converges. What, exactly, does this mean?
- (iv) For each natural number  $n$  let  $b_n = \sum_{k=1}^n a_k^2$ . Rephrase (iii) in terms of the sequence  $(b_n)$ .
- (v) Is the sequence  $(b_n)$  nondecreasing?
- (vi) What, then, does (\*) say about the sequence  $(b_n)$ ?
- (vii) For each natural number  $n$  let  $c_n = \sum_{k=1}^n \frac{1}{k^2}$ . What do we know about the sequence  $(c_n)$  from third term Calculus? What does (\*) say about the sequence  $(c_n)$ ?
- (viii) The conclusion we are trying to prove is that the series  $\sum_{k=1}^{\infty} \frac{1}{k} a_k$  converges absolutely. What does this mean?
- (ix) For each natural number  $n$  let  $s_n = \sum_{k=1}^n \frac{1}{k} |a_k|$ . Rephrase (viii) in terms of the sequence  $(s_n)$ .
- (x) Explain how for each  $n$  we may regard the number  $s_n$  as the dot product of two vectors in  $\mathbb{R}^n$ .
- (xi) Apply the *Cauchy-Schwarz inequality* to the dot product in (x). Use (vi) and (vii) to establish that the sequence  $(s_n)$  is bounded above.
- (xii) Use (\*) one last time—keeping in mind what you said in (ix).

**4.4. Answers to Odd-Numbered Exercises**

- (1)  $\frac{3}{4}$
- (3)  $(\sqrt{a_1}, \sqrt{a_2}, \dots, \sqrt{a_n}), (\frac{1}{\sqrt{a_1}}, \frac{1}{\sqrt{a_2}}, \dots, \frac{1}{\sqrt{a_n}})$
- (5)  $B$
- (7)  $a, b, c, d, e, f$
- (9)  $5, 3, -1, 4$
- (11)  $4, 5, 1, 0$
- (13)  $1, 3, -4$
- (15)
  - (a) 15
  - (b)  $>$
  - (c)  $<$
  - (d)  $=$



**Part 2**

# **VECTOR SPACES**





## CHAPTER 5

# VECTOR SPACES

### 5.1. Background

**Topics:** real and complex vector spaces, vectors, scalars.

In the following definition  $\mathbb{F}$  may be taken to be an arbitrary field. For this collection of exercises, however, we will be interested in only two cases,  $\mathbb{F} = \mathbb{R}$  (the field of real numbers) and  $\mathbb{F} = \mathbb{C}$  (the field of complex numbers).

**5.1.1. Definition.** A VECTOR SPACE is a set  $V$  together with operations of addition and scalar multiplication which satisfy the following axioms:

- (1) if  $\mathbf{x}, \mathbf{y} \in V$ , then  $\mathbf{x} + \mathbf{y} \in V$ ;
- (2)  $(\mathbf{x} + \mathbf{y}) + \mathbf{z} = \mathbf{x} + (\mathbf{y} + \mathbf{z})$  for every  $\mathbf{x}, \mathbf{y}, \mathbf{z} \in V$  (associativity);
- (3) there exists  $\mathbf{0} \in V$  such that  $\mathbf{x} + \mathbf{0} = \mathbf{x}$  for every  $\mathbf{x} \in V$  (existence of additive identity);
- (4) for every  $\mathbf{x} \in V$  there exists  $-\mathbf{x} \in V$  such that  $\mathbf{x} + (-\mathbf{x}) = \mathbf{0}$  (existence of additive inverses);
- (5)  $\mathbf{x} + \mathbf{y} = \mathbf{y} + \mathbf{x}$  for every  $\mathbf{x}, \mathbf{y} \in V$  (commutativity);
- (6) if  $\alpha \in \mathbb{F}$  and  $\mathbf{x} \in V$ , then  $\alpha\mathbf{x} \in V$ ;
- (7)  $\alpha(\mathbf{x} + \mathbf{y}) = \alpha\mathbf{x} + \alpha\mathbf{y}$  for every  $\alpha \in \mathbb{F}$  and every  $\mathbf{x}, \mathbf{y} \in V$ ;
- (8)  $(\alpha + \beta)\mathbf{x} = \alpha\mathbf{x} + \beta\mathbf{x}$  for every  $\alpha, \beta \in \mathbb{F}$  and every  $\mathbf{x} \in V$ ;
- (9)  $(\alpha\beta)\mathbf{x} = \alpha(\beta\mathbf{x})$  for every  $\alpha, \beta \in \mathbb{F}$  and every  $\mathbf{x} \in V$ ; and
- (10)  $1\mathbf{x} = \mathbf{x}$  for every  $\mathbf{x} \in V$ .

When  $\mathbb{F} = \mathbb{R}$  we speak of  $V$  as a REAL VECTOR SPACE and when  $\mathbb{F} = \mathbb{C}$  we call it a COMPLEX VECTOR SPACE.

**5.1.2. Definition.** An  $n \times n$ -matrix is NONSINGULAR if its determinant is not zero.

## 5.2. Exercises

- (1) Let  $V$  be the set of all real numbers. Define an operation of “addition” by

$$x \boxplus y = \text{the maximum of } x \text{ and } y$$

for all  $x, y \in V$ . Define an operation of “scalar multiplication” by

$$\alpha \boxtimes x = \alpha x$$

for all  $\alpha \in \mathbb{R}$  and  $x \in V$ .

Under the operations  $\boxplus$  and  $\boxtimes$  the set  $V$  is not a vector space. The vector space axioms (see 5.1.1 (1)–(10)) which *fail* to hold are \_\_\_\_\_, \_\_\_\_\_, \_\_\_\_\_, and \_\_\_\_\_.

- (2) Let  $V$  be the set of all real numbers  $x$  such that  $x \geq 0$ . Define an operation of “addition” by

$$x \boxplus y = xy + 1$$

for all  $x, y \in V$ . Define an operation of “scalar multiplication” by

$$\alpha \boxtimes x = \alpha^2 x$$

for all  $\alpha \in \mathbb{R}$  and  $x \in V$ .

Under the operations  $\boxplus$  and  $\boxtimes$  the set  $V$  \_\_\_\_\_ (is/is not) a vector space. If it is not, list all the vector space axioms (see 5.1.1 (1)–(10)) which *fail* to hold. Answer: The axioms which are *not* satisfied are \_\_\_\_\_.

- (3) Let  $V$  be  $\mathbb{R}^2$ , the set of all ordered pairs  $(x, y)$  of real numbers. Define an operation of “addition” by

$$(u, v) \boxplus (x, y) = (u + x + 1, v + y + 1)$$

for all  $(u, v)$  and  $(x, y)$  in  $V$ . Define an operation of “scalar multiplication” by

$$\alpha \boxtimes (x, y) = (\alpha x, \alpha y)$$

for all  $\alpha \in \mathbb{R}$  and  $(x, y) \in V$ .

Under the operations  $\boxplus$  and  $\boxtimes$  the set  $V$  is not a vector space. The vector space axioms (see 5.1.1 (1)–(10)) which *fail* to hold are \_\_\_\_\_ and \_\_\_\_\_.

- (4) Let  $V$  be  $\mathbb{R}^2$ , the set of all ordered pairs  $(x, y)$  of real numbers. Define an operation of “addition” by

$$(u, v) \boxplus (x, y) = (u + x, 0)$$

for all  $(u, v)$  and  $(x, y)$  in  $V$ . Define an operation of “scalar multiplication” by

$$\alpha \boxtimes (x, y) = (\alpha x, \alpha y)$$

for all  $\alpha \in \mathbb{R}$  and  $(x, y) \in V$ .

Under the operations  $\boxplus$  and  $\boxtimes$  the set  $V$  is not a vector space. The vector space axioms (see 5.1.1 (1)–(10)) which *fail* to hold are \_\_\_\_\_, \_\_\_\_\_, and \_\_\_\_\_.

- (5) Let  $V$  be the set of all  $n \times n$  matrices of real numbers. Define an operation of “addition” by

$$A \boxplus B = \frac{1}{2}(AB + BA)$$

for all  $A, B \in V$ . Define an operation of “scalar multiplication” by

$$\alpha \boxtimes A = \mathbf{0}$$

for all  $\alpha \in \mathbb{R}$  and  $A \in V$ .

Under the operations  $\boxplus$  and  $\boxtimes$  the set  $V$  is not a vector space. The vector space axioms (see 5.1.1 (1)–(10)) which *fail* to hold are \_\_\_\_\_, \_\_\_\_\_, and \_\_\_\_\_.

- (6) Below are portions of proofs of four results about vector spaces which establish the fact that multiplying a vector  $x$  by the scalar  $-1$  produces  $-x$ , the additive inverse of  $x$ . Fill in the missing steps and the missing reasons. Choose reasons from the following list.

(H) Hypothesis

(1)–(10) Vector space axioms, see 5.1.1

(PA) Proposition A

(PB) Proposition B

(PC) Proposition C

(RN) Property of the Real Numbers

**5.2.1. Proposition (A).** *A vector  $x$  in a vector space  $V$  has at most one additive inverse. That is, if  $\mathbf{y}$  and  $\mathbf{z}$  are vectors in  $V$  such that  $\mathbf{x} + \mathbf{y} = \mathbf{0}$  and  $\mathbf{x} + \mathbf{z} = \mathbf{0}$ , then  $\mathbf{y} = \mathbf{z}$ .*

PROOF. Suppose that  $\mathbf{x} + \mathbf{y} = \mathbf{0}$  and  $\mathbf{x} + \mathbf{z} = \mathbf{0}$ . Then

$$\begin{aligned}
 \mathbf{y} &= \underline{\hspace{2cm}} && (\text{reason: } \underline{\hspace{2cm}}) \\
 &= \mathbf{y} + (\mathbf{x} + \mathbf{z}) && (\text{reason: } \underline{\hspace{2cm}}) \\
 &= \underline{\hspace{2cm}} && (\text{reason: (2)}) \\
 &= (\mathbf{x} + \mathbf{y}) + \mathbf{z} && (\text{reason: } \underline{\hspace{2cm}}) \\
 &= \underline{\hspace{2cm}} && (\text{reason: (H)}) \\
 &= \underline{\hspace{2cm}} && (\text{reason: (5)}) \\
 &= \mathbf{z} && (\text{reason: } \underline{\hspace{2cm}}).
 \end{aligned}$$

□

**5.2.2. Proposition (B).** *If  $\mathbf{x} \in V$  where  $V$  is a vector space and  $\mathbf{x} + \mathbf{x} = \mathbf{x}$ , then  $\mathbf{x} = \mathbf{0}$ .*

PROOF. If  $\mathbf{x} \in V$  and  $\mathbf{x} + \mathbf{x} = \mathbf{x}$ , then

$$\begin{aligned}
 \mathbf{x} &= \mathbf{x} + \mathbf{0} && (\text{reason: } \underline{\hspace{2cm}}) \\
 &= \underline{\hspace{2cm}} && (\text{reason: (4)}) \\
 &= (\mathbf{x} + \mathbf{x}) + (-\mathbf{x}) && (\text{reason: } \underline{\hspace{2cm}}) \\
 &= \underline{\hspace{2cm}} && (\text{reason: (H)}) \\
 &= \mathbf{0} && (\text{reason: } \underline{\hspace{2cm}}).
 \end{aligned}$$

□

**5.2.3. Proposition (C).** *If  $x$  is a vector in a vector space  $V$ , then  $0\mathbf{x} = \mathbf{0}$ .*

PROOF. If  $\mathbf{x} \in V$ , then

$$\begin{aligned}
 0\mathbf{x} &= (0 + 0)\mathbf{x} && (\text{reason: } \underline{\hspace{2cm}}) \\
 &= \underline{\hspace{2cm}} && (\text{reason: (8)})
 \end{aligned}$$

Thus  $0\mathbf{x} = \mathbf{0}$  (reason:  $\underline{\hspace{2cm}}$ ).

□

**5.2.4. Proposition (D).** *If  $x$  is a vector in a vector space  $V$ , then  $(-1)\mathbf{x}$  is  $-\mathbf{x}$ , the additive inverse of  $\mathbf{x}$ .*

PROOF. If  $\mathbf{x} \in V$ , then

$$\begin{aligned}\mathbf{x} + (-1) \cdot \mathbf{x} &= \underline{\hspace{2cm}} && \text{(reason: (10))} \\ &= (1 + (-1)) \cdot \mathbf{x} && \text{(reason: } \underline{\hspace{1cm}} \text{)} \\ &= 0 \cdot \mathbf{x} && \text{(reason: } \underline{\hspace{1cm}} \text{)} \\ &= \mathbf{0} && \text{(reason: } \underline{\hspace{1cm}} \text{)}.\end{aligned}$$

It then follows immediately from  $\underline{\hspace{2cm}}$  that  $(-1) \cdot \mathbf{x} = -\mathbf{x}$ . □

- (7) In this exercise we prove that multiplying the zero vector by an arbitrary scalar produces the zero vector. For each step of the proof give the appropriate reason. Choose reasons from the following list.

(1)–(10) Vector space axioms 5.1.1.

(PB) Proposition 5.2.2

(RN) Property of the Real Numbers

**5.2.5. Proposition (E).** *If  $\mathbf{0}$  is the zero vector in a vector space and  $\alpha$  is a scalar, then  $\alpha \cdot \mathbf{0} = \mathbf{0}$ .*

PROOF. Let  $\mathbf{0}$  be the zero vector of some vector space. Then for every scalar  $\alpha$

$$\begin{aligned}\alpha \cdot \mathbf{0} &= \alpha \cdot (\mathbf{0} + \mathbf{0}) && \text{reason: } \underline{\hspace{2cm}} \\ &= \alpha \cdot \mathbf{0} + \alpha \cdot \mathbf{0} && \text{reason: } \underline{\hspace{2cm}}\end{aligned}$$

It then follows immediately from  $\underline{\hspace{2cm}}$  that  $\alpha \cdot \mathbf{0} = \mathbf{0}$ . □

- (8) In this exercise we prove that the product of a scalar and a vector is zero if and only if either the scalar or the vector is zero. After each step of the proof give the appropriate reason. Choose reasons from the following list.

(H) Hypothesis.

(1)–(10) Vector space axioms 5.1.1.

(PC),(PE) Propositions 5.2.3 and 5.2.5, respectively.

(RN) Property of the Real Numbers.

**5.2.6. Proposition.** *Suppose that  $\mathbf{x}$  is a vector and  $\alpha$  is a scalar. Then  $\alpha\mathbf{x} = \mathbf{0}$  if and only if  $\alpha = 0$  or  $\mathbf{x} = \mathbf{0}$ .*

PROOF. We have already shown in  $\underline{\hspace{2cm}}$  and  $\underline{\hspace{2cm}}$  that if  $\alpha = 0$  or  $\mathbf{x} = \mathbf{0}$ , then  $\alpha\mathbf{x} = \mathbf{0}$ .

To prove the converse we suppose that  $\alpha\mathbf{x} = \mathbf{0}$  and that  $\alpha \neq 0$ ; and we prove that  $\mathbf{x} = \mathbf{0}$ . This conclusion results from the following easy calculation:

$$\begin{aligned}\mathbf{x} &= 1 \cdot \mathbf{x} && \text{reason: } \underline{\hspace{2cm}} \\ &= \left(\frac{1}{\alpha} \cdot \alpha\right) \cdot \mathbf{x} && \text{reasons: } \underline{\hspace{2cm}} \text{ and } \underline{\hspace{2cm}} \\ &= \frac{1}{\alpha} \cdot (\alpha \cdot \mathbf{x}) && \text{reason: } \underline{\hspace{2cm}} \\ &= \frac{1}{\alpha} \cdot \mathbf{0} && \text{reason: } \underline{\hspace{2cm}} \\ &= \mathbf{0} && \text{reason: } \underline{\hspace{2cm}}.\end{aligned}$$

□

**5.3. Problems**

- (1) Prove that if  $V$  is a vector space, then its additive identity is unique. That is, show that if  $\mathbf{0}$  and  $\tilde{\mathbf{0}}$  are vectors in  $V$  such that  $\mathbf{x} + \mathbf{0} = \mathbf{x}$  for all  $x \in V$  and  $\mathbf{x} + \tilde{\mathbf{0}} = \mathbf{x}$  for all  $x \in V$ , then  $\mathbf{0} = \tilde{\mathbf{0}}$ .
- (2) Let  $V$  be the set of all real numbers  $x$  such that  $x > 0$ . Define an operation of “addition” by

$$x \boxplus y = xy$$

for all  $x, y \in V$ . Define an operation of “scalar multiplication” by

$$\alpha \boxdot x = x^\alpha$$

for all  $\alpha \in \mathbb{R}$  and  $x \in V$ .

Prove that under the operations  $\boxplus$  and  $\boxdot$  the set  $V$  is a vector space.

- (3) With the usual operations of addition and scalar multiplication the set of all  $n \times n$  matrices of real numbers is a vector space: in particular, all the vector space axioms (see 5.1.1 (1)–(10)) are satisfied. Explain clearly why the set of all nonsingular  $n \times n$  matrices of real numbers is *not* a vector space under these same operations.

**5.4. Answers to Odd-Numbered Exercises**

- (1) 3, 4, 7, 8
- (3) 7, 8
- (5) 2, 4, 10
- (7) 3, 7, PB

## CHAPTER 6

# SUBSPACES

### 6.1. Background

**Topics:** subspaces of a vector space

**6.1.1. Definition.** A nonempty subset of  $M$  of a vector space  $V$  is a **SUBSPACE** of  $V$  if it is closed under addition and scalar multiplication. (That is: if  $\mathbf{x}$  and  $\mathbf{y}$  belong to  $M$ , so does  $\mathbf{x} + \mathbf{y}$ ; and if  $\mathbf{x}$  belongs to  $M$  and  $\alpha \in \mathbb{R}$ , then  $\alpha\mathbf{x}$  belongs to  $M$ .)

**6.1.2. Notation.** We use the notation  $M \preceq V$  to indicate that  $M$  is a subspace of a vector space  $V$ .

**6.1.3. Notation.** Here are some frequently encountered families of functions:

$$\mathcal{F} = \mathcal{F}[a, b] = \{f: f \text{ is a real valued function on the interval } [a, b]\} \quad (6.1.1)$$

$$\mathcal{P} = \mathcal{P}[a, b] = \{p: p \text{ is a polynomial function on } [a, b]\} \quad (6.1.2)$$

$$\mathcal{P}_4 = \mathcal{P}_4[a, b] = \{p \in \mathcal{P}: \text{the degree of } p \text{ is less than } 4\} \quad (6.1.3)$$

$$\mathcal{Q}_4 = \mathcal{Q}_4[a, b] = \{p \in \mathcal{P}: \text{the degree of } p \text{ is equal to } 4\} \quad (6.1.4)$$

$$\mathcal{C} = \mathcal{C}[a, b] = \{f \in \mathcal{F}: f \text{ is continuous}\} \quad (6.1.5)$$

$$\mathcal{D} = \mathcal{D}[a, b] = \{f \in \mathcal{F}: f \text{ is differentiable}\} \quad (6.1.6)$$

$$\mathcal{K} = \mathcal{K}[a, b] = \{f \in \mathcal{F}: f \text{ is a constant function}\} \quad (6.1.7)$$

$$\mathcal{B} = \mathcal{B}[a, b] = \{f \in \mathcal{F}: f \text{ is bounded}\} \quad (6.1.8)$$

$$\mathcal{J} = \mathcal{J}[a, b] = \{f \in \mathcal{F}: f \text{ is integrable}\} \quad (6.1.9)$$

(A function  $f \in \mathcal{F}$  is **BOUNDED** if there exists a number  $M \geq 0$  such that  $|f(x)| \leq M$  for all  $x$  in  $[a, b]$ . It is **(RIEMANN) INTEGRABLE** if it is bounded and  $\int_a^b f(x) dx$  exists.)

**6.1.4. Definition.** If  $A$  and  $B$  are subsets of a vector space then the **SUM** of  $A$  and  $B$ , denoted by  $A + B$ , is defined by

$$A + B := \{\mathbf{a} + \mathbf{b}: \mathbf{a} \in A \text{ and } \mathbf{b} \in B\}.$$

**6.1.5. Definition.** Let  $M$  and  $N$  be subspaces of a vector space  $V$ . If  $M \cap N = \{\mathbf{0}\}$  and  $M + N = V$ , then  $V$  is the **(INTERNAL) DIRECT SUM** of  $M$  and  $N$ . In this case we write

$$V = M \oplus N.$$

In this case the subspaces  $M$  and  $N$  are **COMPLEMENTARY** and each is the **COMPLEMENT** of the other.





- (6) For a fixed interval  $[a, b]$ , which sets of functions in the list 6.1.3 are vector subspaces of which?

Answer:

\_\_\_\_\_  $\preceq$  \_\_\_\_\_  $\preceq$  \_\_\_\_\_  $\preceq$  \_\_\_\_\_  $\preceq$  \_\_\_\_\_  $\preceq$  \_\_\_\_\_  $\preceq$  \_\_\_\_\_  $\preceq$  \_\_\_\_\_ .

- (7) Let  $M$  be the plane  $x + y + z = 0$  and  $N$  be the line  $x = y = z$  in  $\mathbb{R}^3$ . The purpose of this exercise is to confirm that  $\mathbb{R}^3 = M \oplus N$ . This requires establishing three things: (i)  $M$  and  $N$  are subspaces of  $\mathbb{R}^3$  (which is *very* easy and which we omit); (ii)  $\mathbb{R}^3 = M + N$ ; and (iii)  $M \cap N = \{\mathbf{0}\}$ .

- (a) To show that  $\mathbb{R}^3 = M + N$  we need  $\mathbb{R}^3 \subseteq M + N$  and  $M + N \subseteq \mathbb{R}^3$ . Since  $M \subseteq \mathbb{R}^3$  and  $N \subseteq \mathbb{R}^3$ , it is clear that  $M + N \subseteq \mathbb{R}^3$ . So all that is required is to show that  $\mathbb{R}^3 \subseteq M + N$ . That is, given a vector  $\mathbf{x} = (x_1, x_2, x_3)$  in  $\mathbb{R}^3$  we must find vectors  $\mathbf{m} = (m_1, m_2, m_3)$  in  $M$  and  $\mathbf{n} = (n_1, n_2, n_3)$  in  $N$  such that  $\mathbf{x} = \mathbf{m} + \mathbf{n}$ . Find two such vectors.

Answer:  $\mathbf{m} = \frac{1}{3} ( \text{_____} , \text{_____} , \text{_____} )$

and  $\mathbf{n} = \frac{1}{3} ( \text{_____} , \text{_____} , \text{_____} )$ .

- (b) The last thing to verify is that  $M \cap N = \{\mathbf{0}\}$ ; that is, that the only vector  $M$  and  $N$  have in common is the zero vector. Suppose that a vector  $\mathbf{x} = (x_1, x_2, x_3)$  belongs to both  $M$  and  $N$ . Since  $\mathbf{x} \in M$  it must satisfy the equation

$$x_1 + x_2 + x_3 = 0. \quad (1)$$

since  $\mathbf{x} \in N$  it must satisfy the equations

$$x_1 = x_2 \quad \text{and} \quad (2)$$

$$x_2 = x_3. \quad (3)$$

Solve the system of equations (1)–(3).

Answer:  $\mathbf{x} = ( \text{_____} , \text{_____} , \text{_____} )$ .

- (8) Let  $\mathcal{C} = \mathcal{C}[-1, 1]$  be the vector space of all continuous real valued functions on the interval  $[-1, 1]$ . A function  $f$  in  $\mathcal{C}$  is EVEN if  $f(-x) = f(x)$  for all  $x \in [-1, 1]$ ; it is ODD if  $f(-x) = -f(x)$  for all  $x \in [-1, 1]$ . Let  $\mathcal{C}_o = \{f \in \mathcal{C} : f \text{ is odd} \}$  and  $\mathcal{C}_e = \{f \in \mathcal{C} : f \text{ is even} \}$ . To show that  $\mathcal{C} = \mathcal{C}_o \oplus \mathcal{C}_e$  we need to show 3 things.

- (i)  $\mathcal{C}_o$  and  $\mathcal{C}_e$  are subspaces of  $\mathcal{C}$ . This is quite simple: let's do just one part of the proof. We will show that  $\mathcal{C}_o$  is closed under addition. After each step of the following proof indicate the justification for that step. Make your choices from the following list.

(A) Arithmetic of real numbers.

(DA) Definition of addition of functions.

(DE) Definition of "even function."

(DO) Definition of "odd function."

(H) Hypothesis (that is, our assumptions or suppositions).

PROOF. Let  $f, g \in \mathcal{C}_o$ . Then

$$\begin{aligned}(f+g)(-x) &= f(-x) + g(-x) && \text{reason: } \underline{\hspace{2cm}} \\ &= -f(x) + (-g(x)) && \text{reason: } \underline{\hspace{2cm}} \text{ and } \underline{\hspace{2cm}} \\ &= -(f(x) + g(x)) && \text{reason: } \underline{\hspace{2cm}} \\ &= -(f+g)(x). && \text{reason: } \underline{\hspace{2cm}}\end{aligned}$$

Thus  $f+g \in \mathcal{C}_o$ . reason  $\underline{\hspace{2cm}}$ . □

- (ii)  $\mathcal{C}_o \cap \mathcal{C}_e = \{\mathbf{0}\}$  (where  $\mathbf{0}$  is the constant function on  $[-1, 1]$  whose value is zero). Again choose from the reasons listed in part (i) to justify the given proof.

PROOF. Suppose  $f \in \mathcal{C}_o \cap \mathcal{C}_e$ . Then for each  $x \in [-1, 1]$

$$\begin{aligned}f(x) &= f(-x) && \text{reason: } \underline{\hspace{2cm}} \\ &= -f(x). && \text{reason: } \underline{\hspace{2cm}}\end{aligned}$$

Thus  $f(x) = 0$  for every  $x \in [-1, 1]$ ; that is,  $f = \mathbf{0}$ . reason:  $\underline{\hspace{2cm}}$ . □

- (iii)  $\mathcal{C} = \mathcal{C}_o + \mathcal{C}_e$ . To verify this we must show that every continuous function  $f$  on  $[-1, 1]$  can be written as the sum of an odd function  $j$  and an even function  $k$ . It turns out that the functions  $j$  and  $k$  can be written as linear combinations of the given function  $f$  and the function  $g$  defined by  $g(x) = f(-x)$  for  $-1 \leq x \leq 1$ . What are the appropriate coefficients?

$$\begin{aligned}\text{Answer: } \quad j &= \underline{\hspace{2cm}} f + \underline{\hspace{2cm}} g \\ k &= \underline{\hspace{2cm}} f + \underline{\hspace{2cm}} g.\end{aligned}$$

- (9) Let  $M$  be the line  $x = y = z$  and  $N$  be the line  $x = \frac{1}{2}y = \frac{1}{3}z$  in  $\mathbb{R}^3$ .

(a) The line  $M$  is the set of all scalar multiples of the vector  $(1, \underline{\hspace{2cm}}, \underline{\hspace{2cm}})$ .

(b) The line  $N$  is the set of all scalar multiples of the vector  $(1, \underline{\hspace{2cm}}, \underline{\hspace{2cm}})$ .

(c) The set  $M + N$  is (geometrically speaking) a  $\underline{\hspace{2cm}}$  in  $\mathbb{R}^3$ ; its equation is  $ax + by + z = 0$  where  $a = \underline{\hspace{2cm}}$  and  $b = \underline{\hspace{2cm}}$ .

- (10) Let  $M$  be the plane  $x - y + z = 0$  and  $N$  be the plane  $x + 2y - z = 0$  in  $\mathbb{R}^3$ . State in one short sentence how you know that  $\mathbb{R}^3$  is *not* the direct sum of  $M$  and  $N$ .

Answer:  $\underline{\hspace{10cm}}$ .

- (11) Let  $M$  be the plane  $2x - 3y + 4z + 1 = 0$  and  $N$  be the line  $\frac{x}{4} = \frac{y}{2} = \frac{z}{3}$  in  $\mathbb{R}^3$ . State in one short sentence how you know that  $\mathbb{R}^3$  is *not* the direct sum of  $M$  and  $N$ .

Answer:  $\underline{\hspace{10cm}}$ .

- (12) Let  $M$  be the plane  $x + y + z = 0$  and  $N$  be the line  $x - 1 = \frac{1}{2}y = z + 2$  in  $\mathbb{R}^3$ . State in one short sentence how you know that  $\mathbb{R}^3$  is *not* the direct sum of  $M$  and  $N$ .

Answer:  $\underline{\hspace{10cm}}$ .

- (13) Let  $M$  be the line  $x = y = z$  and  $N$  be the line  $\frac{x}{4} = \frac{y}{2} = \frac{z}{3}$  in  $\mathbb{R}^3$ . State in one short sentence how you know that  $\mathbb{R}^3$  is *not* the direct sum of  $M$  and  $N$ .

Answer:  $\underline{\hspace{10cm}}$ .

(14) Let  $M$  be the plane  $x + y + z = 0$  and  $N$  be the line  $x = -\frac{3}{4}y = 3z$ . The purpose of this exercise is to see (in two different ways) that  $\mathbb{R}^3$  is *not* the direct sum of  $M$  and  $N$ .

(a) If  $\mathbb{R}^3$  were equal to  $M \oplus N$ , then  $M \cap N$  would contain only the zero vector. Show that this is not the case by finding a nonzero vector  $\mathbf{x}$  in  $\mathbb{R}^3$  which belongs to  $M \cap N$ .

Answer:  $\mathbf{x} = (\text{____}, \text{____}, 1)$ .

(b) If  $\mathbb{R}^3$  were equal to  $M \oplus N$ , then, in particular, we would have  $\mathbb{R}^3 = M + N$ . Since both  $M$  and  $N$  are subsets of  $\mathbb{R}^3$ , it is clear that  $M + N \subseteq \mathbb{R}^3$ . Show that the reverse inclusion  $\mathbb{R}^3 \subseteq M + N$  is *not* correct by finding a vector  $\mathbf{x} \in \mathbb{R}^3$  which cannot be written in the form  $\mathbf{m} + \mathbf{n}$  where  $\mathbf{m} \in M$  and  $\mathbf{n} \in N$ .

Answer:  $\mathbf{x} = (-6, 8, a)$  is such a vector provided that  $a \neq \text{____}$ .

(c) We have seen in part (b) that  $M + N \neq \mathbb{R}^3$ . Then what *is*  $M + N$ ?

Answer:  $M + N = \text{____}$ .

### 6.3. Problems

- (1) Let  $M$  and  $N$  be subspaces of a vector space  $V$ . Consider the following subsets of  $V$ .
  - (a)  $M \cap N$ . (A vector  $\mathbf{v}$  belongs to  $M \cap N$  if it belongs to both  $M$  and  $N$ .)
  - (b)  $M \cup N$ . (A vector  $\mathbf{v}$  belongs to  $M \cup N$  if it belongs to either  $M$  or  $N$ .)
  - (c)  $M + N$ . (A vector  $\mathbf{v}$  belongs to  $M + N$  if there are vectors  $\mathbf{m} \in M$  and  $\mathbf{n} \in N$  such that  $\mathbf{v} = \mathbf{m} + \mathbf{n}$ .)
  - (d)  $M - N$ . (A vector  $\mathbf{v}$  belongs to  $M - N$  if there are vectors  $\mathbf{m} \in M$  and  $\mathbf{n} \in N$  such that  $\mathbf{v} = \mathbf{m} - \mathbf{n}$ .)

For each of the sets (a)–(d) above, either prove that it *is* a subspace of  $V$  or give a counterexample to show that it *need not* be a subspace of  $V$ .

- (2) Let  $\mathcal{C} = \mathcal{C}[0, 1]$  be the family of continuous real valued functions on the interval  $[0, 1]$ . Define

$$f_1(t) = t \quad \text{and} \quad f_2(t) = t^4$$

for  $0 \leq t \leq 1$ . Let  $M$  be the set of all functions of the form  $\alpha f_1 + \beta f_2$  where  $\alpha, \beta \in \mathbb{R}$ . And let  $N$  be the set of all functions  $g$  in  $\mathcal{C}$  which satisfy

$$\int_0^1 t g(t) dt = 0 \quad \text{and} \quad \int_0^1 t^4 g(t) dt = 0.$$

Is  $\mathcal{C}$  the direct sum of  $M$  and  $N$ ? (Give a careful proof of your claim and illustrate it with an example. What does your result say, for instance, about the function  $h$  defined by  $h(t) = t^2$  for  $0 \leq t \leq 1$ .)

- (3) Let  $V$  be a vector space.
  - (a) Let  $\mathcal{M}$  be a family of subspaces of  $V$ . Prove that the intersection  $\bigcap \mathcal{M}$  of this family is itself a subspace of  $V$ .
  - (b) Let  $A$  be a set of vectors in  $V$ . Explain carefully why it makes sense to say that the intersection of the family of all subspaces containing  $A$  is “the smallest subspace of  $V$  which contains  $A$ ”.
  - (c) Prove that the smallest subspace of  $V$  which contains  $A$  is in fact the span of  $A$ .
- (4) In  $\mathbb{R}^3$  let  $M$  be the line  $x = y = z$ ,  $N$  be the line  $x = \frac{1}{2}y = \frac{1}{3}z$ , and  $L = M + N$ . Give a careful proof that  $L = M \oplus N$ .
- (5) Let  $V$  be a vector space and suppose that  $V = M \oplus N$ . Show that for every  $\mathbf{v} \in V$  there exist unique vectors  $\mathbf{m} \in M$  and  $\mathbf{n} \in N$  such that  $\mathbf{v} = \mathbf{m} + \mathbf{n}$ . *Hint.* It should be clear that the only thing you have to establish is the *uniqueness* of the vectors  $\mathbf{m}$  and  $\mathbf{n}$ . To this end, suppose that a vector  $\mathbf{v}$  in  $V$  can be written as  $\mathbf{m}_1 + \mathbf{n}_1$  and it can also be written as  $\mathbf{m}_2 + \mathbf{n}_2$  where  $\mathbf{m}_1, \mathbf{m}_2 \in M$  and  $\mathbf{n}_1, \mathbf{n}_2 \in N$ . Prove that  $\mathbf{m}_1 = \mathbf{m}_2$  and  $\mathbf{n}_1 = \mathbf{n}_2$ .

**6.4. Answers to Odd-Numbered Exercises**

- (1) (d)
- (3)  $0, -2$
- (5) (a), (c), and (d)
- (7) (a)  $2x_1 - x_2 - x_3, -x_1 + 2x_2 - x_3, -x_1 - x_2 + 2x_3, x_1 + x_2 + x_3, x_1 + x_2 + x_3, x_1 + x_2 + x_3,$   
(b)  $0, 0, 0$
- (9) (a)  $1, 1$   
(b)  $2, 3$   
(c) plane,  $1, -2$
- (11)  $M$  is not a subspace of  $\mathbb{R}^3$ .
- (13)  $M + N$  is a plane, not all of  $\mathbb{R}^3$ .



## CHAPTER 7

# LINEAR INDEPENDENCE

### 7.1. Background

**Topics:** linear combinations, span, linear dependence and independence.

**7.1.1. Remark.** Some authors of linear algebra texts make it appear as if the terms *linear dependence* and *linear independence*, *span*, and *basis* pertain only to *finite* sets of vectors. This is extremely misleading. The expressions should make sense for *arbitrary* sets. In particular, do not be misled into believing that a basis for a vector space must be a finite set of vectors (or a sequence of vectors). While it is true that in most elementary linear algebra courses the emphasis is on the study of *finite dimensional* vector spaces, bases for vector spaces may be *very* large indeed. I recommend the following definitions.

**7.1.2. Definition.** Recall that a vector  $\mathbf{y}$  is a LINEAR COMBINATION of distinct vectors  $\mathbf{x}_1, \dots, \mathbf{x}_n$  if there exist scalars  $\alpha_1, \dots, \alpha_n$  such that  $\mathbf{y} = \sum_{k=1}^n \alpha_k \mathbf{x}_k$ . *Note:* linear combinations *are* finite sums. The linear combination  $\sum_{k=1}^n \alpha_k \mathbf{x}_k$  is TRIVIAL if all the coefficients  $\alpha_1, \dots, \alpha_n$  are zero. If at least one  $\alpha_k$  is different from zero, the linear combination is NONTRIVIAL.

**7.1.3. Example.** In  $\mathbb{R}^2$  the vector  $(8, 2)$  is a linear combination of the vectors  $(1, 1)$  and  $(1, -1)$  because  $(8, 2) = 5(1, 1) + 3(1, -1)$ .

**7.1.4. Example.** In  $\mathbb{R}^3$  the vector  $(1, 2, 3)$  is *not* a linear combination of the vectors  $(1, 1, 0)$  and  $(1, -1, 0)$ .

**7.1.5. Definition.** Suppose that  $A$  is a subset (finite or not) of a vector space  $V$ . The SPAN of  $A$  is the set of all linear combinations of elements of  $A$ . Another way of saying the same thing: the SPAN of  $A$  is the smallest subspace of  $V$  which contains  $A$ . (That these characterizations are equivalent is not completely obvious. Proof is required. See problem 3 in chapter 6. We denote the span of  $A$  by  $\text{span } A$ . If  $U = \text{span } A$ , we say that  $A$  SPANS  $U$  or that  $U$  is SPANNED BY  $A$ .

**7.1.6. Example.** For each  $n = 0, 1, 2, \dots$  define a function  $\mathbf{p}_n$  on  $\mathbb{R}$  by  $\mathbf{p}_n(x) = x^n$ . Let  $\mathcal{P}$  be the set of polynomial functions on  $\mathbb{R}$ . It is a subspace of the vector space of continuous functions on  $\mathbb{R}$ . Then  $\mathcal{P} = \text{span}\{\mathbf{p}_0, \mathbf{p}_1, \mathbf{p}_2, \dots\}$ . The exponential function  $\exp$ , whose value at  $x$  is  $e^x$ , is not in the span of the set  $\{\mathbf{p}_0, \mathbf{p}_1, \mathbf{p}_2, \dots\}$ .

**7.1.7. Definition.** A subset  $A$  (finite or not) of a vector space is LINEARLY DEPENDENT if the zero vector  $\mathbf{0}$  can be written as a nontrivial linear combination of elements of  $A$ ; that is, if there exist vectors  $\mathbf{x}_1, \dots, \mathbf{x}_n \in A$  and scalars  $\alpha_1, \dots, \alpha_n$ , **not all zero**, such that  $\sum_{k=1}^n \alpha_k \mathbf{x}_k = \mathbf{0}$ . A subset of a vector space is LINEARLY INDEPENDENT if it is not linearly dependent.

Technically, it is a *set* of vectors that is linearly dependent or independent. Nevertheless, these terms are frequently used as if they were properties of the vectors themselves. For instance, if  $S = \{\mathbf{x}_1, \dots, \mathbf{x}_n\}$  is a finite set of vectors in a vector space, you may see the assertions “the set  $S$  is linearly independent” and “the vectors  $\mathbf{x}_1, \dots, \mathbf{x}_n$  are linearly independent” used interchangeably.

**7.1.8. Example.** The (vectors going from the origin to) points on the unit circle in  $\mathbb{R}^2$  are linearly dependent. Reason: If  $\mathbf{x} = (1, 0)$ ,  $\mathbf{y} = (-\frac{1}{2}, \frac{\sqrt{3}}{2})$ , and  $\mathbf{z} = (\frac{1}{2}, \frac{\sqrt{3}}{2})$ , then  $\mathbf{x} + \mathbf{y} + (-1)\mathbf{z} = \mathbf{0}$ .

**7.1.9. Example.** For each  $n = 0, 1, 2, \dots$  define a function  $\mathbf{p}_n$  on  $\mathbb{R}$  by  $\mathbf{p}_n(x) = x^n$ . Then the set  $\{\mathbf{p}_0, \mathbf{p}_1, \mathbf{p}_2, \dots\}$  is a linearly independent subset of the vector space of continuous functions on  $\mathbb{R}$ .



## 7.2. Exercises

- (1) Show that in the space  $\mathbb{R}^3$  the vectors  $\mathbf{x} = (1, 1, 0)$ ,  $\mathbf{y} = (0, 1, 2)$ , and  $\mathbf{z} = (3, 1, -4)$  are linearly dependent by finding scalars  $\alpha$  and  $\beta$  such that  $\alpha\mathbf{x} + \beta\mathbf{y} + \mathbf{z} = \mathbf{0}$ .

Answer:  $\alpha = \underline{\hspace{1cm}}$ ,  $\beta = \underline{\hspace{1cm}}$ .

- (2) Let  $\mathbf{w} = (1, 1, 0, 0)$ ,  $\mathbf{x} = (1, 0, 1, 0)$ ,  $\mathbf{y} = (0, 0, 1, 1)$ , and  $\mathbf{z} = (0, 1, 0, 1)$ .

(a) We can show that  $\{\mathbf{w}, \mathbf{x}, \mathbf{y}, \mathbf{z}\}$  is not a spanning set for  $\mathbb{R}^4$  by finding a vector  $\mathbf{u}$  in  $\mathbb{R}^4$  such that  $\mathbf{u} \notin \text{span}\{\mathbf{w}, \mathbf{x}, \mathbf{y}, \mathbf{z}\}$ . One such vector is  $\mathbf{u} = (1, 2, 3, a)$  where  $a$  is any number *except*  $\underline{\hspace{1cm}}$ .

(b) Show that  $\{\mathbf{w}, \mathbf{x}, \mathbf{y}, \mathbf{z}\}$  is a linearly dependent set of vectors by finding scalars  $\alpha$ ,  $\gamma$ , and  $\delta$  such that  $\alpha\mathbf{w} + \mathbf{x} + \gamma\mathbf{y} + \delta\mathbf{z} = \mathbf{0}$ .

Answer:  $\alpha = \underline{\hspace{1cm}}$ ,  $\gamma = \underline{\hspace{1cm}}$ ,  $\delta = \underline{\hspace{1cm}}$ .

(c) Show that  $\{\mathbf{w}, \mathbf{x}, \mathbf{y}, \mathbf{z}\}$  is a linearly dependent set by writing  $\mathbf{z}$  as a linear combination of  $\mathbf{w}$ ,  $\mathbf{x}$ , and  $\mathbf{y}$ . Answer:  $\mathbf{z} = \underline{\hspace{1cm}} \mathbf{w} + \underline{\hspace{1cm}} \mathbf{x} + \underline{\hspace{1cm}} \mathbf{y}$ .

- (3) Let  $p(x) = x^2 + 2x - 3$ ,  $q(x) = 2x^2 - 3x + 4$ , and  $r(x) = ax^2 - 1$ . The set  $\{p, q, r\}$  is linearly dependent if  $a = \underline{\hspace{1cm}}$ .

- (4) Show that in the vector space  $\mathbb{R}^3$  the vectors  $\mathbf{x} = (1, 2, -1)$ ,  $\mathbf{y} = (3, 1, 1)$ , and  $\mathbf{z} = (5, -5, 7)$  are linearly dependent by finding scalars  $\alpha$  and  $\beta$  such that  $\alpha\mathbf{x} + \beta\mathbf{y} + \mathbf{z} = \mathbf{0}$ .

Answer:  $\alpha = \underline{\hspace{1cm}}$ ,  $\beta = \underline{\hspace{1cm}}$ .

- (5) Let  $f_1(x) = \sin x$ ,  $f_2(x) = \cos(x + \pi/6)$ , and  $f_3(x) = \sin(x - \pi/4)$  for  $0 \leq x \leq 2\pi$ . Show that  $\{f_1, f_2, f_3\}$  is linearly dependent by finding constants  $\alpha$  and  $\beta$  such that  $\alpha f_1 - 2f_2 - \beta f_3 = 0$ .

Answer:  $\alpha = \underline{\hspace{1cm}}$  and  $\beta = \underline{\hspace{1cm}}$ .

- (6) In the space  $\mathcal{C}[0, \pi]$  let  $f$ ,  $g$ ,  $h$ , and  $j$  be the vectors defined by

$$f(x) = 1$$

$$g(x) = x$$

$$h(x) = \cos x$$

$$j(x) = \cos^2 \frac{x}{2}$$

for  $0 \leq x \leq \pi$ . Show that  $f$ ,  $g$ ,  $h$ , and  $j$  are linearly dependent by writing  $j$  as a linear combination of  $f$ ,  $g$ , and  $h$ .

Answer:  $j = \underline{\hspace{1cm}} f + \underline{\hspace{1cm}} g + \underline{\hspace{1cm}} h$ .

- (7) Let  $\mathbf{u} = (\lambda, 1, 0)$ ,  $\mathbf{v} = (1, \lambda, 1)$ , and  $\mathbf{w} = (0, 1, \lambda)$ . Find **all** values of  $\lambda$  which make  $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$  a linearly dependent subset of  $\mathbb{R}^3$ . Answer:  $\underline{\hspace{2cm}}$

- (8) Let  $\mathbf{u} = (1, 0, -2)$ ,  $\mathbf{v} = (1, 2, \lambda)$ , and  $\mathbf{w} = (2, 1, -1)$ . Find **all** values of  $\lambda$  which make  $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$  a linearly dependent subset of  $\mathbb{R}^3$ . Answer:  $\underline{\hspace{2cm}}$

- (9) Let  $p(x) = x^3 - x^2 + 2x + 3$ ,  $q(x) = 3x^3 + x^2 - x - 1$ ,  $r(x) = x^3 + 2x + 2$ , and  $s(x) = 7x^3 + ax^2 + 5$ . The set  $\{p, q, r, s\}$  is linearly dependent if  $a = \underline{\hspace{1cm}}$ .

(10) In the space  $\mathcal{C}[0, \pi]$  define the vectors  $f$ ,  $g$ , and  $h$  by

$$f(x) = x$$

$$g(x) = \sin x$$

$$h(x) = \cos x$$

for  $0 \leq x \leq \pi$ . We show that  $f$ ,  $g$ , and  $h$  are linearly independent. This is accomplished by showing that if  $\alpha f + \beta g + \gamma h = \mathbf{0}$ , then  $\alpha = \beta = \gamma = 0$ . So we start by supposing that  $\alpha f + \beta g + \gamma h = \mathbf{0}$ ; that is,

$$\alpha x + \beta \sin x + \gamma \cos x = 0 \tag{1}$$

for all  $x \in [0, \pi]$ .

(a) We see that  $\gamma$  must be zero by setting  $x = \underline{\hspace{1cm}}$  in equation (1).

Now differentiate (1) to obtain

$$\alpha + \beta \cos x = 0 \tag{2}$$

for all  $x \in [0, \pi]$ .

(b) We see that  $\alpha$  must be zero by setting  $x = \underline{\hspace{1cm}}$  in equation (2).

Differentiate (2) to obtain

$$-\beta \sin x = 0 \tag{3}$$

for all  $x \in [0, \pi]$ .

(c) We conclude that  $\beta = 0$  by setting  $x = \underline{\hspace{1cm}}$  in (3).

## 7.3. Problems

- (1) In the space  $\mathcal{C}[0, 1]$  define the vectors  $f$ ,  $g$ , and  $h$  by

$$\begin{aligned}f(x) &= x \\g(x) &= e^x \\h(x) &= e^{-x}\end{aligned}$$

for  $0 \leq x \leq 1$ . Use the **definition** of *linear independence* to show that the functions  $f$ ,  $g$ , and  $h$  are linearly independent.

- (2) Let  $a$ ,  $b$ , and  $c$  be distinct real numbers. Use the **definition** of *linear independence* to give a careful proof that the vectors  $(1, 1, 1)$ ,  $(a, b, c)$ , and  $(a^2, b^2, c^2)$  form a linearly independent subset of  $\mathbb{R}^3$ .
- (3) Let  $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$  be a linearly independent set in a vector space  $V$ . Use the **definition** of *linear independence* to give a careful proof that the set  $\{\mathbf{u} + \mathbf{v}, \mathbf{u} + \mathbf{w}, \mathbf{v} + \mathbf{w}\}$  is linearly independent in  $V$ .
- (4) You are the leader of an engineering group in the company you work for and have a routine computation that has to be done repeatedly. At your disposal is an intern, Kim, a beginning high school student, who is bright but has had no advanced mathematics. In particular, Kim knows nothing about vectors or matrices.

Here is the computation that is needed. Three vectors,  $\mathbf{a}$ ,  $\mathbf{b}$ , and  $\mathbf{c}$  are specified in  $\mathbb{R}^5$ . (Denote their span by  $M$ .) Also specified is a (sometimes long) list of other vectors  $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$  in  $\mathbb{R}^5$ . The problem is to

- (1) determine which of the vectors in  $S$  belong to  $M$ , and
- (2) for each vector  $\mathbf{v}_k \in S$  which *does* belong to  $M$

find constants  $\alpha$ ,  $\beta$ , and  $\gamma$  such that  $\mathbf{v}_k = \alpha\mathbf{a} + \beta\mathbf{b} + \gamma\mathbf{c}$ .

Kim has access to Computer Algebra System (Maple, or a similar program) with a Linear Algebra package. Write a simple and efficient algorithm (that is, a set of instructions) which will allow Kim to carry out the desired computation repeatedly. The algorithm should be simple in the sense that it uses only the most basic linear algebra commands (for example, *Matrix*, *Vector*, *Transpose*, *RowReducedEchelonForm*, etc. in Maple). Remember, you must tell Kim everything: how to set up the appropriate matrices, what operations to perform on them, and how to interpret the results. The algorithm should be as efficient as you can make it. For example, it would certainly not be efficient for Kim to retype the coordinates of  $\mathbf{a}$ ,  $\mathbf{b}$ , and  $\mathbf{c}$  for each new  $\mathbf{v}_k$ .

Include in your write-up an actual printout showing how your algorithm works in some special case of your own invention. (For this example, the set  $S$  need contain only 5 or 6 vectors, some in  $U$ , some not.)

- (5) The point of this problem is not just to get a correct answer to (a)–(c) below using tools you may have learned elsewhere, but *to give a careful explanation of how to apply the linear algebra techniques you have already encountered to solve this problem in a systematic fashion*. For background you may wish to read a bit about networks and *Kirchhoff's laws* (see, for example, [7] *Topic: Analyzing Networks*, pages 72–77 or [1] *Electrical Networks*, pages 538–542).

Consider an electrical network having four nodes **A**, **B**, **C**, and **D** connected by six branches **1**, **...**, **6**. Branch **1** connects **A** and **B**; branch **2** connects **B** and **D**; branch **3** connects **C** and **B**; branch **4** connects **C** and **D**; branch **5** connects **A** and **C**; and branch **6** connects **A** and **D**.

The current in branch **k** is  $I_k$ , where  $k = 1, \dots, 6$ . There is a 17 volt battery in branch **1** producing the current  $I_1$  which flows from **A** to **B**. In branches **2**, **4**, and **5** there are 0.5 ohm resistors; and in branches **1**, **3**, and **6** there are 1 ohm resistors.

- (a) Find the current in each branch. (Explain any minus signs which occur in your answer.)
- (b) Find the voltage drop across each branch.
- (c) Let  $p_{\mathbf{n}}$  be the potential at node  $\mathbf{n} = \mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}$ . The voltage drop across the branch connecting node **j** to node **k** is the difference in the potentials at nodes **j** and **k**. Suppose the network is grounded at **D** (so that  $p_{\mathbf{D}} = 0$ ). Find the potential at the other nodes.

**7.4. Answers to Odd-Numbered Exercises**

(1)  $-3, 2$

(3)  $7$

(5)  $\sqrt{3} - 1, \sqrt{6}$

(7)  $-\sqrt{2}, 0, \sqrt{2}$

(9)  $-3$



## CHAPTER 8

# BASIS FOR A VECTOR SPACE

### 8.1. Background

**Topics:** basis, dimension.

**8.1.1. Definition.** A set  $B$  (finite or not) of vectors in a vector space  $V$  is a **BASIS** for  $V$  if it is linearly independent and spans  $V$ .

**8.1.2. Example.** The vectors  $\mathbf{e}^1 = (1, 0, 0)$ ,  $\mathbf{e}^2 = (0, 1, 0)$ , and  $\mathbf{e}^3 = (0, 0, 1)$  constitute a basis for the vector space  $\mathbb{R}^3$ .

**8.1.3. Example.** More generally, consider the vector space  $\mathbb{R}^n$  of all  $n$ -tuples of real numbers. For each natural number  $k$  between 1 and  $n$  let  $\mathbf{e}^k$  be the vector which is 1 in the  $k^{\text{th}}$ -coordinate and 0 in all the others. Then the set  $\{\mathbf{e}^1, \mathbf{e}^2, \dots, \mathbf{e}^n\}$  is a basis for  $\mathbb{R}^n$ . It is called the **STANDARD BASIS** for  $\mathbb{R}^n$ .

**8.1.4. Example.** For each  $n = 0, 1, 2, \dots$  define a function  $\mathbf{p}_n$  on  $\mathbb{R}$  by  $\mathbf{p}_n(x) = x^n$ . Then the set  $\{\mathbf{p}_0, \mathbf{p}_1, \mathbf{p}_2, \dots\}$  is a basis for the vector space  $\mathcal{P}$  of polynomial functions on  $\mathbb{R}$ .

Two important facts of linear algebra are that regardless of the size of the space *every* vector space has a basis and that every subspace has a complement.

**8.1.5. Theorem.** *Let  $B$  be a linearly independent set of vectors in a vector space  $V$ . Then there exists a set  $C$  of vectors in  $V$  such that  $B \cup C$  is a basis for  $V$ .*

**8.1.6. Corollary.** *Every vector space has a basis.*

**8.1.7. Corollary.** *Let  $V$  be a vector space. If  $M \preceq V$ , then there exists  $N \preceq V$  such that  $M \oplus N = V$ .*

The next theorem says that any two bases for a vector space are the same size.

**8.1.8. Theorem.** *If  $B$  and  $C$  are bases for the same vector space, then there is a one-to-one correspondence from  $B$  onto  $C$ .*

**8.1.9. Definition.** A vector space  $V$  is **FINITE DIMENSIONAL** if it has a finite basis. Its **DIMENSION** (denoted by  $\dim V$ ) is the number of elements in the basis. If  $V$  does not have a finite basis it is **INFINITE DIMENSIONAL**.

**8.1.10. Theorem.** *If  $M$  and  $N$  are subspaces of a finite dimensional vector space, then*  
$$\dim(M + N) = \dim M + \dim N - \dim(M \cap N).$$

**8.2. Exercises**

- (1) Let  $\mathbf{u} = (2, 0, -1)$ ,  $\mathbf{v} = (3, 1, 0)$ , and  $\mathbf{w} = (1, -1, c)$  where  $c \in \mathbb{R}$ . The set  $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$  is a basis for  $\mathbb{R}^3$  provided that  $c$  is *not* equal to \_\_\_\_ .
- (2) Let  $\mathbf{u} = (1, -1, 3)$ ,  $\mathbf{v} = (1, 0, 1)$ , and  $\mathbf{w} = (1, 2, c)$  where  $c \in \mathbb{R}$ . The set  $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$  is a basis for  $\mathbb{R}^3$  provided that  $c$  is *not* equal to \_\_\_\_ .
- (3) The dimension of  $\mathfrak{M}_{2 \times 2}$ , the vector space of all  $2 \times 2$  matrices of real numbers is \_\_\_\_ .
- (4) The dimension of  $\mathfrak{T}_2$ , the vector space of all  $2 \times 2$  matrices of real numbers with zero trace is \_\_\_\_ .
- (5) The dimension of the vector space of all real valued polynomial functions on  $\mathbb{R}$  of degree 4 or less is \_\_\_\_ .
- (6) In  $\mathbb{R}^4$  let  $M$  be the subspace spanned by the vectors  $(1, 1, 1, 0)$  and  $(0, -4, 1, 5)$  and let  $N$  be the subspace spanned by  $(0, -2, 1, 2)$  and  $(1, -1, 1, 3)$ . One vector which belongs to both  $M$  and  $N$  is  $(1, \text{____}, \text{____}, \text{____})$ . The dimension of  $M \cap N$  is \_\_\_\_ and the dimension of  $M + N$  is \_\_\_\_ .



**8.3. Problems**

- (1) Exhibit a basis for  $\mathfrak{M}_{2 \times 2}$ , the vector space of all  $2 \times 2$  matrices of real numbers.
- (2) Exhibit a basis for  $\mathfrak{T}_2$ , the vector space of all  $2 \times 2$  matrices of real numbers with zero trace.
- (3) Exhibit a basis for  $\mathfrak{S}_3$ , the vector space of all symmetric  $3 \times 3$  matrices of real numbers.
- (4) Let  $\mathfrak{U}$  be the set of all matrices of real numbers of the form  $\begin{bmatrix} u & -u-x \\ 0 & x \end{bmatrix}$  and  $\mathfrak{V}$  be the set of all real matrices of the form  $\begin{bmatrix} v & 0 \\ w & -v \end{bmatrix}$ . Exhibit a basis for  $\mathfrak{U}$ , for  $\mathfrak{V}$ , for  $\mathfrak{U} + \mathfrak{V}$ , and for  $\mathfrak{U} \cap \mathfrak{V}$ .
- (5) Prove that the vectors  $(1, 1, 0)$ ,  $(1, 2, 3)$ , and  $(2, -1, 5)$  form a basis for  $\mathbb{R}^3$ .
- (6) Let  $V$  be a vector space and  $A$  be a linearly independent subset of  $V$ . Prove that  $A$  is a basis for  $V$  if and only if it is a maximal linearly independent subset of  $V$ . (If  $A$  is a linearly independent subset of  $V$  we say that it is a MAXIMAL linearly independent set if the addition of any vector at all to  $A$  will result in a set which is *not* linearly independent.)
- (7) Let  $V$  be a vector space and  $A$  a subset of  $V$  which spans  $V$ . Prove that  $A$  is a basis for  $V$  if and only if it is a minimal spanning set. (If  $A$  is a set which spans  $V$  we say that it is a MINIMAL spanning set if the removal of any vector at all from  $A$  will result in a set which does *not* span  $V$ .)

**8.4. Answers to Odd-Numbered Exercises**

(1)  $-2$

(3)  $4$

(5)  $5$

**Part 3**

**LINEAR MAPS BETWEEN VECTOR  
SPACES**



## CHAPTER 9

# LINEARITY

### 9.1. Background

**Topics:** linear maps between vector spaces, kernel, nullspace, nullity, range, rank, isomorphism.

**9.1.1. Definition.** A function  $f: A \rightarrow B$  is ONE-TO-ONE (or INJECTIVE) if  $u = v$  in  $A$  whenever  $f(u) = f(v)$  in  $B$ .

**9.1.2. Definition.** A function  $f: A \rightarrow B$  is ONTO (or SURJECTIVE) if for every  $b \in B$  there exists  $a \in A$  such that  $b = f(a)$ .

**9.1.3. Definition.** A function  $f: A \rightarrow B$  is a ONE-TO-ONE CORRESPONDENCE (or BIJECTIVE) if it is both injective and surjective (one-to-one and onto).

**9.1.4. Definition.** A map  $T: V \rightarrow W$  between vector spaces is LINEAR if

$$T(\mathbf{x} + \mathbf{y}) = T\mathbf{x} + T\mathbf{y} \quad \text{for all } \mathbf{x}, \mathbf{y} \in V \quad (9.1.1)$$

and

$$T(\alpha\mathbf{x}) = \alpha T\mathbf{x} \quad \text{for all } \mathbf{x} \in V \text{ and } \alpha \in \mathbb{F}. \quad (9.1.2)$$

Here  $\mathbb{F} = \mathbb{R}$  if  $V$  and  $W$  are real vector spaces and  $\mathbb{F} = \mathbb{C}$  if they are complex vector spaces.

A scalar valued linear map on a vector space  $V$  is a LINEAR FUNCTIONAL.

A linear map is frequently called a LINEAR TRANSFORMATION, and, in case the domain and codomain are the same, it is often called a (LINEAR) OPERATOR. The family of all linear transformations from  $V$  into  $W$  is denoted by  $\mathfrak{L}(V, W)$ . We shorten  $\mathfrak{L}(V, V)$  to  $\mathfrak{L}(V)$ .

Two oddities of notation concerning linear transformations deserve comment. First, the value of  $T$  at  $\mathbf{x}$  is usually written  $T\mathbf{x}$  rather than  $T(\mathbf{x})$ . Naturally the parentheses are used whenever their omission would create ambiguity. For example, in (9.1.1) above  $T\mathbf{x} + \mathbf{y}$  is not an acceptable substitute for  $T(\mathbf{x} + \mathbf{y})$ . Second, the symbol for composition of two linear transformations is ordinarily omitted. If  $S \in \mathfrak{L}(U, V)$  and  $T \in \mathfrak{L}(V, W)$ , then the composite of  $T$  and  $S$  is denoted by  $TS$  (rather than by  $T \circ S$ ). As a consequence of this convention when  $T \in \mathfrak{L}(V)$  the linear operator  $T \circ T$  is written as  $T^2$ ,  $T \circ T \circ T$  as  $T^3$ , and so on.

For future reference here are two obvious properties of a linear map.

**9.1.5. Proposition.** *If  $T: V \rightarrow W$  is a linear map between vector spaces, then  $T(\mathbf{0}) = \mathbf{0}$ .*

**9.1.6. Proposition.** *If  $T: V \rightarrow W$  is a linear map between vector spaces, then  $T(-\mathbf{x}) = -T\mathbf{x}$  for every  $\mathbf{x} \in V$ .*

You should prove these propositions if (and only if) it is not immediately obvious to you how to do so.

**9.1.7. Definition.** Let  $T: V \rightarrow W$  be a linear transformation between vector spaces. Then  $\ker T$ , the KERNEL of  $T$ , is defined to be the set of all  $\mathbf{x}$  in  $V$  such that  $T\mathbf{x} = \mathbf{0}$ . The kernel of  $T$  is also called the *nullspace* of  $T$ . If  $V$  is finite dimensional, the dimension of the kernel of  $T$  is the NULLITY of  $T$ .

Also,  $\text{ran } T$ , the RANGE of  $T$ , is the set of all  $\mathbf{y}$  in  $W$  such that  $\mathbf{y} = T\mathbf{x}$  for some  $\mathbf{x}$  in  $V$ . If the range of  $T$  is finite dimensional, its dimension is the RANK of  $T$ .

**9.1.8. Notation.** Let  $V$  be a vector space. We denote the IDENTITY MAP on  $V$  (that is, the map  $\mathbf{x} \mapsto \mathbf{x}$  from  $V$  into itself) by  $I_V$ , or just  $I$ .

The following fundamental result is proved in most linear algebra texts.

**9.1.9. Theorem.** *If  $T: V \rightarrow W$  is a linear map between finite dimensional vector spaces, then*

$$\text{rank}(T) + \text{nullity}(T) = \dim V.$$

**9.1.10. Definition.** Let  $T: V \rightarrow W$  and  $S: W \rightarrow V$  be linear maps. If  $ST = I_V$ , then  $T$  is a RIGHT INVERSE for  $S$  and  $S$  is a LEFT INVERSE for  $T$ . The mapping  $T$  is INVERTIBLE (or is an ISOMORPHISM) if there exists a linear transformation, which we denote by  $T^{-1}: W \rightarrow V$ , such that

$$TT^{-1} = I_W \quad \text{and} \quad T^{-1}T = I_V.$$

The vector spaces  $V$  and  $W$  are ISOMORPHIC if there exists an isomorphism  $T$  from  $V$  to  $W$ .

**9.1.11. Notation.** Let  $V$  and  $W$  be vector spaces. We denote by  $\mathfrak{L}(V, W)$  the set of all linear maps from  $V$  into  $W$  and by  $\mathfrak{L}(V)$  the set of all linear operators  $T: V \rightarrow V$ .

## 9.2. Exercises

- (1) Define
- $T: \mathbb{R}^3 \rightarrow \mathbb{R}^4$
- by

$$T\mathbf{x} = (x_1 - x_3, x_1 + x_2, x_3 - x_2, x_1 - 2x_2)$$

for all  $\mathbf{x} = (x_1, x_2, x_3)$  in  $\mathbb{R}^3$ .

- (a) Then
- $T(1, -2, 3) = ( \quad , \quad , \quad , \quad )$
- .

- (b) Find a vector
- $\mathbf{x} \in \mathbb{R}^3$
- such that
- $T\mathbf{x} = (8, 9, -5, 0)$
- .

Answer:  $\mathbf{x} = ( \quad , \quad , \quad )$ .

- (2) Define
- $T: \mathbb{R}^4 \rightarrow \mathbb{R}^3$
- by

$$T\mathbf{x} = (2x_1 + x_3 + x_4, x_1 - 2x_2 - x_3, x_2 - x_3 + x_4)$$

for all  $\mathbf{x} = (x_1, x_2, x_3, x_4)$  in  $\mathbb{R}^4$ .

- (a) Then
- $T(2, 1, -1, 3) = ( \quad , \quad , \quad )$
- .

- (b) Find a vector
- $\mathbf{x} \in \mathbb{R}^4$
- such that
- $T\mathbf{x} = (3, -1, -3)$
- .

Answer:  $\mathbf{x} = ( \quad , \quad , \quad , \quad )$ .

- (3) Let
- $T$
- be the linear map from
- $\mathbb{R}^3$
- to
- $\mathbb{R}^3$
- defined by

$$T(x, y, z) = (x + 2y - z, 2x + 3y + z, 4x + 7y - z).$$

The kernel of  $T$  is (geometrically) a \_\_\_\_\_ whose equation(s) is(are) \_\_\_\_\_;  
and the range of  $T$  is geometrically a \_\_\_\_\_ whose equation(s) is(are) \_\_\_\_\_.

- (4) Let
- $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$
- be the linear transformation whose action on the standard basis vectors of
- $\mathbb{R}^3$
- is

$$T(1, 0, 0) = (1, -\frac{3}{2}, 2)$$

$$T(0, 1, 0) = (-3, \frac{9}{2}, -6)$$

$$T(0, 0, 1) = (2, -3, 4).$$

Then  $T(5, 1, -1) = ( \quad , \quad , \quad )$ . The kernel of  $T$  is the \_\_\_\_\_ whose equation is  $x + ay + bz = 0$  where  $a = \frac{y}{2} = \frac{z}{c} = \frac{z}{d}$  where  $c = \frac{y}{2} = \frac{z}{d}$  and where  $d = \frac{y}{2} = \frac{z}{d}$ .

- (5) Let
- $\mathcal{P}$
- be the vector space of all polynomial functions on
- $\mathbb{R}$
- with real coefficients. Define linear transformations
- $T, D: \mathcal{P} \rightarrow \mathcal{P}$
- by

$$(Dp)(x) = p'(x)$$

and

$$(Tp)(x) = xp(x)$$

for all  $x \in \mathbb{R}$ .

- (a) Let
- $p(x) = x^3 - 7x^2 + 5x + 6$
- for all
- $x \in \mathbb{R}$
- . Then
- $((D+T)(p))(x) = x^4 - ax^3 + bx^2 - cx + d$
- where
- $a = \frac{y}{2} = \frac{z}{c} = \frac{z}{d}$
- and
- $d = \frac{y}{2} = \frac{z}{d}$
- .

- (b) Let
- $p$
- be as in (a). Then
- $(DTp)(x) = ax^3 - bx^2 + cx + d$
- where
- $a = \frac{y}{2} = \frac{z}{c} = \frac{z}{d}$
- and
- $d = \frac{y}{2} = \frac{z}{d}$
- .

- (c) Let
- $p$
- be as in (a). Then
- $(TDp)(x) = ax^3 - bx^2 + cx + d$
- where
- $a = \frac{y}{2} = \frac{z}{c} = \frac{z}{d}$
- and
- $d = \frac{y}{2} = \frac{z}{d}$
- .

- (d) Evaluate (and simplify) the commutator
- $[D, T] := DT - TD$
- .

Answer:  $[D, T] = \frac{y}{2} = \frac{z}{c} = \frac{z}{d}$ .

- (e) Find a number
- $p$
- such that
- $(TD)^p = T^p D^p + TD$
- . Answer:
- $p = \frac{y}{2} = \frac{z}{c} = \frac{z}{d}$
- .

- (6) Let  $\mathcal{C} = \mathcal{C}[a, b]$  be the vector space of all continuous real valued functions on the interval  $[a, b]$  and  $\mathcal{C}^1 = \mathcal{C}^1[a, b]$  be the vector space of all continuously differentiable real valued functions on  $[a, b]$ . (Recall that a function is CONTINUOUSLY DIFFERENTIABLE if it has a derivative and the derivative is continuous.) Let  $D: \mathcal{C}^1 \rightarrow \mathcal{C}$  be the linear transformation defined by

$$Df = f'$$

and let  $T: \mathcal{C} \rightarrow \mathcal{C}^1$  be the linear transformation defined by

$$(Tf)(x) = \int_a^x f(t) dt$$

for all  $f \in \mathcal{C}$  and  $x \in [a, b]$ .

- (a) Compute (and simplify)  $(DTf)(x)$ . Answer: \_\_\_\_\_ .
- (b) Compute (and simplify)  $(TDf)(x)$ . Answer: \_\_\_\_\_ .
- (c) The kernel of  $T$  is \_\_\_\_\_ .
- (d) The range of  $T$  is  $\{g \in \mathcal{C}^1: \text{_____}\}$
- (7) In this exercise we prove that a linear transformation  $T: V \rightarrow W$  between two vector spaces is one-to-one if and only if its kernel contains only the zero vector. After each step of the proof give the appropriate reason. Choose reasons from the following list.
- (DK) Definition of “kernel”.
- (DL) Definition of “linear”.
- (DO) Definition of “one-to-one”.
- (H) Hypothesis.
- (Pa) Proposition 9.1.5.
- (Pb) Proposition 9.1.6.
- (VA) Vector space arithmetic (consequences of vector space axioms, definition of subtraction of vectors, etc.)

PROOF. Suppose that  $T$  is one-to-one. We show that  $\ker T = \{\mathbf{0}_V\}$ . Since  $\mathbf{0}_V \in \ker T$  (reason: \_\_\_\_\_ and \_\_\_\_\_), we need only show that  $\ker T \subseteq \{\mathbf{0}_V\}$ ; that is, we show that if  $\mathbf{x} \in \ker T$ , then  $\mathbf{x} = \mathbf{0}_V$ . So let  $\mathbf{x} \in \ker T$ . Then  $T\mathbf{x} = \mathbf{0}_W$  (reason: \_\_\_\_\_ and \_\_\_\_\_) and  $T\mathbf{0}_V = \mathbf{0}_W$  (reason: \_\_\_\_\_). From this we conclude that  $\mathbf{x} = \mathbf{0}_V$  (reason: \_\_\_\_\_ and \_\_\_\_\_).

Now we prove the converse. Suppose that  $\ker T = \{\mathbf{0}_V\}$ . We wish to show that  $T$  is one-to-one. Let  $\mathbf{x}, \mathbf{y} \in V$  and suppose that  $T\mathbf{x} = T\mathbf{y}$ . Then

$$\begin{aligned} T(\mathbf{x} - \mathbf{y}) &= T(\mathbf{x} + (-\mathbf{y})) && \text{reason: } \_\_\_\_\_\_ \\ &= T\mathbf{x} + T(-\mathbf{y}) && \text{reason: } \_\_\_\_\_\_ \\ &= T\mathbf{x} + (-T\mathbf{y}) && \text{reason: } \_\_\_\_\_\_ \\ &= T\mathbf{x} - T\mathbf{y} && \text{reason: } \_\_\_\_\_\_ \\ &= \mathbf{0}_W && \text{reason: } \_\_\_\_\_\_ \text{ and } \_\_\_\_\_\_ \end{aligned}$$

Then  $\mathbf{x} - \mathbf{y} \in \ker T$  (reason: \_\_\_\_\_). So  $\mathbf{x} - \mathbf{y} = \mathbf{0}_V$  (reason: \_\_\_\_\_); that is,  $\mathbf{x} = \mathbf{y}$  (reason: \_\_\_\_\_). Thus  $T$  is one-to-one (reason: \_\_\_\_\_ and \_\_\_\_\_).  $\square$

- (8) Let  $\mathcal{C}^1(\mathbb{R})$  be the vector space of all functions defined on the real line  $\mathbb{R}$  which have continuous derivatives at each point of  $\mathbb{R}$  and  $\mathcal{C}(\mathbb{R})$  be the vector space of continuous functions on  $\mathbb{R}$ . Define the function  $T: \mathcal{C}^1(\mathbb{R}) \rightarrow \mathcal{C}(\mathbb{R})$  by

$$(Tf)(t) = f'(t) + 3f(t)$$



for every  $t \in \mathbb{R}$ . (Notice that  $T$  is a linear map.) The kernel of  $T$  is the set of all scalar multiples of the function  $g$  where  $g(t) = \underline{\hspace{2cm}}$  for each  $t$ . Thus the kernel of the linear map  $T$  is the solution space of the differential equation  $\underline{\hspace{2cm}}$ .

- (9) Let  $\mathcal{C}^2(\mathbb{R})$  be the vector space of all functions defined on the real line  $\mathbb{R}$  which have continuous second derivatives at each point of  $\mathbb{R}$  and  $\mathcal{C}(\mathbb{R})$  be the vector space of continuous functions on  $\mathbb{R}$ . Define the function  $T: \mathcal{C}^2(\mathbb{R}) \rightarrow \mathcal{C}(\mathbb{R})$  by

$$(Tf)(t) = f''(t) + f(t)$$

for every  $t \in \mathbb{R}$ . (Notice that  $T$  is a linear map.) Assume that the kernel of  $T$  is two dimensional. Then  $\ker T = \text{span}\{g, h\}$  where  $g(t) = \underline{\hspace{2cm}}$  and  $h(t) = \underline{\hspace{2cm}}$  for all  $t$ . Thus the kernel of the linear map  $T$  is the solution space of the differential equation  $\underline{\hspace{2cm}}$ .

- (10) Define a function  $k$  on the unit square  $[0, 1] \times [0, 1]$  by

$$k(x, y) = \begin{cases} x, & \text{for } 0 \leq x \leq y \leq 1 \\ y, & \text{for } 0 \leq y < x \leq 1 \end{cases}.$$

Define an integral operator  $K$  on the vector space  $\mathcal{C}[0, 1]$  of continuous real valued functions on  $[0, 1]$  by

$$(Kf)(x) = \int_0^1 k(x, y)f(y) dy$$

for  $0 \leq x \leq 1$ . Find the function  $Kf$  when  $f$  is the function defined by  $f(x) = x^2$  for  $0 \leq x \leq 1$ .

Answer:  $(Kf)(x) = \underline{\hspace{2cm}}$ .

- (11) Let  $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3: \mathbf{x} \mapsto (x_1 + 3x_2 - 2x_3, x_1 - 4x_3, x_1 + 6x_2)$ .

(a) The kernel of  $T$  is a  $\underline{\hspace{2cm}}$  in  $\mathbb{R}^3$  given by the equation(s)

$\underline{\hspace{2cm}}$ .

(b) The range of  $T$  is a  $\underline{\hspace{2cm}}$  in  $\mathbb{R}^3$  given by the equation(s)

$\underline{\hspace{2cm}}$ .

- (12) Let  $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3: (x, y) \mapsto (2x - 3y, x + 2y + 1, 5x - 2y)$ . State in one short sentence how you know that  $T$  is *not* a linear transformation.

Answer:  $\underline{\hspace{2cm}}$ .

- (13) Let  $\mathbf{a} = (1, 0, 0, 0)$ ,  $\mathbf{b} = (1, 1, 0, 0)$ ,  $\mathbf{c} = (1, 1, 1, 0)$ , and  $\mathbf{d} = (1, 1, 1, 1)$ . Suppose that  $T: \mathbb{R}^4 \rightarrow \mathbb{R}^7$  is a mapping such that  $T(\mathbf{a}) = T(\mathbf{b}) = T(\mathbf{c}) = T(\mathbf{d}) = \mathbf{0}$  and that  $T(3, -19, 7, -8) = (1, 1, 1, -3, 6, 2, 5)$ . State in a short sentence or two how you know that  $T$  is *not* a linear transformation.

Answer:  $\underline{\hspace{2cm}}$ .

- (14) Suppose that  $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$  is a mapping (not identically zero) whose range is contained in the paraboloid  $z = x^2 + y^2$ . State in a short sentence or two how you know that  $T$  is *not* a linear transformation.

Answer:  $\underline{\hspace{2cm}}$ .

- (15) Let  $T: \mathbb{R}^2 \rightarrow \mathbb{R}^4: (x, y) \mapsto (2x - 3y, x - 7y, x + 2y + 1, 5x - 2y)$ . State in one short sentence how you know that  $T$  is *not* a linear transformation.

Answer:  $\underline{\hspace{2cm}}$ .

- (16) Let  $\mathbf{a} = (1, 1, 0)$  and  $\mathbf{b} = (0, 1, 1)$ , and  $\mathbf{c} = (1, 2, 1)$ . Suppose that  $T: \mathbb{R}^3 \rightarrow \mathbb{R}^5$  is a mapping such that  $T(\mathbf{a}) = T(\mathbf{b}) = \mathbf{0}$  and that  $T(\mathbf{c}) = (1, -3, 6, 2, 5)$ . State in a short sentence or two how you know that  $T$  is *not* a linear transformation.

Answer: \_\_\_\_\_.

- (17) Suppose that  $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  is a mapping (not identically zero) such that  $T(1, 1) = (3, -6)$  and  $T(-2, -2) = (-6, 3)$ . State in a short sentence or two how you know that  $T$  is *not* a linear transformation.

Answer: \_\_\_\_\_.

## 9.3. Problems

- (1) Let  $T: V \rightarrow W$  be a linear transformation between vector spaces and let  $N$  be a subspace of  $W$ . Define  $T^{\leftarrow}(N) := \{\mathbf{v} \in V: T\mathbf{v} \in N\}$ . Prove that  $T^{\leftarrow}(N)$  is a subspace of  $V$ .
- (2) Prove that a linear transformation  $T: \mathbb{R}^3 \rightarrow \mathbb{R}^2$  cannot be one-to-one and that a linear transformation  $S: \mathbb{R}^2 \rightarrow \mathbb{R}^3$  cannot be onto. Generalize these assertions.
- (3) Prove that one-to-one linear transformations preserve linear independence. That is: Let  $T: V \rightarrow W$  be a one-to-one linear transformation between vector spaces and  $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n\}$  be a linearly independent subset of  $V$ . Prove that  $\{T\mathbf{x}_1, T\mathbf{x}_2, \dots, T\mathbf{x}_n\}$  is a linearly independent subset of  $W$ . *Hint.* To prove that the vectors  $T\mathbf{x}_1, T\mathbf{x}_2, \dots, T\mathbf{x}_n$  are linearly independent, it must be shown that the only linear combination of these vectors which equals zero is the trivial linear combination. So *suppose* that  $\sum_{k=1}^n \alpha_k T\mathbf{x}_k = \mathbf{0}$  and *prove* that every  $\alpha_k$  must be zero. Use the result proved in exercise 7.
- (4) The goal of this problem is to understand and write up an introduction to *invertible* linear transformations. Your write-up should explain with spectacular clarity the basic facts about invertible linear transformations. Include answers to the following questions—giving complete proofs or counterexamples as required. (But don't number things in your report to correspond with the items that follow.)
  - (a) If a linear transformation has a right inverse must it have a left inverse?
  - (b) If a linear transformation has a left inverse must it have a right inverse?
  - (c) If a linear transformation has both a left and a right inverse, must it be invertible? (That is, must the left and right inverse be the same?)
  - (d) If a linear transformation  $T$  has a unique right inverse is  $T$  necessarily invertible? *Hint.* Consider  $ST + S - I$ , where  $S$  is a unique right inverse for  $T$ .
  - (e) What is the relationship between a linear transformation being one-to-one and onto and being invertible?
  - (f) Let  $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$  be a linearly independent set of vectors in  $V$ . What condition should a linear transformation  $T: V \rightarrow W$  satisfy so that  $\{T\mathbf{v}_1, \dots, T\mathbf{v}_n\}$  is a linearly independent subset of  $W$ ?
  - (g) Let  $\{\mathbf{u}_1, \dots, \mathbf{u}_n\}$  be a basis for a subspace  $U$  of  $V$ . What conditions should a linear transformation  $T: V \rightarrow W$  satisfy so that  $\{T\mathbf{u}_1, \dots, T\mathbf{u}_n\}$  is a basis for the subspace  $T(U)$ ?
  - (h) Suppose the vectors  $\mathbf{v}_1, \dots, \mathbf{v}_n$  span the vector space  $V$  and  $T: V \rightarrow W$  is a linear transformation. If  $\{T\mathbf{v}_1, \dots, T\mathbf{v}_n\}$  is a basis for  $W$  what can you conclude about the vectors  $\mathbf{v}_1, \dots, \mathbf{v}_n$ ? What can you conclude about the linear transformation  $T$ ?
  - (i) When are two finite dimensional vector spaces isomorphic? (Give a simple—but nontrivial—necessary and sufficient condition *on the spaces*.)
  - (j) Suppose  $S: V \rightarrow V$  is linear and  $V$  has finite dimension. What is the relationship between the following properties of  $S$ ?
    - (1)  $S$  is one-to-one.
    - (2)  $S$  is onto.
    - (3)  $S$  is an isomorphism.
- (5) A sequence of vector spaces and linear maps

$$\cdots \longrightarrow V_{n-1} \xrightarrow{j_n} V_n \xrightarrow{j_{n+1}} V_{n+1} \longrightarrow \cdots$$

is said to be EXACT AT  $V_n$  if  $\text{ran } j_n = \ker j_{n+1}$ . A sequence is EXACT if it is exact at each of its constituent vector spaces. A sequence of vector spaces and linear maps of the form

$$\mathbf{0} \longrightarrow U \xrightarrow{j} V \xrightarrow{k} W \longrightarrow \mathbf{0} \tag{1}$$

is a SHORT EXACT SEQUENCE. (Here  $\mathbf{0}$  denotes the trivial 0-dimensional vector space, and the unlabeled arrows are the obvious linear maps.)

- (a) The sequence (1) is exact at  $U$  if and only if  $j$  is injective.  
 (b) The sequence (1) is exact at  $W$  if and only if  $k$  is surjective.  
 (c) Let  $U$  and  $V$  be vector spaces. Then the following sequence is short exact:

$$\mathbf{0} \longrightarrow U \xrightarrow{\iota_1} U \times V \xrightarrow{\pi_2} V \longrightarrow \mathbf{0}.$$

The indicated linear maps are defined by

$$\iota_1: U \rightarrow U \times V: a \mapsto (a, 0)$$

and

$$\pi_2: U \times V \rightarrow V: (a, b) \mapsto b.$$

- (d) Suppose  $a < b$ . Let  $\mathcal{K}$  be the family of constant functions on the interval  $[a, b]$ ,  $\mathcal{C}^1$  be the family of all continuously differentiable functions on  $[a, b]$ , and  $\mathcal{C}$  be the family of all continuous functions on  $[a, b]$ . Specify linear maps  $j$  and  $k$  so that the following sequence is short exact:

$$\mathbf{0} \longrightarrow \mathcal{K} \xrightarrow{j} \mathcal{C}^1 \xrightarrow{k} \mathcal{C} \longrightarrow \mathbf{0}.$$

- (e) Let  $\mathcal{C}$  be the family of all continuous functions on the interval  $[0, 2]$ . Let  $E_1$  be the mapping from  $\mathcal{C}$  into  $\mathbb{R}$  defined by  $E_1(f) = f(1)$ . (The functional  $E_1$  is called “evaluation at 1”.)

Find a subspace  $\mathcal{F}$  of  $\mathcal{C}$  such that the following sequence is short exact.

$$\mathbf{0} \longrightarrow \mathcal{F} \xrightarrow{\iota} \mathcal{C} \xrightarrow{E_1} \mathbb{R} \longrightarrow \mathbf{0}.$$

- (f) Suppose that the following sequence of finite dimensional vector spaces and linear maps is exact.

$$\mathbf{0} \longrightarrow V_n \xrightarrow{f_n} V_{n-1} \xrightarrow{f_{n-1}} \cdots \xrightarrow{f_2} V_1 \xrightarrow{f_1} V_0 \longrightarrow \mathbf{0}$$

Show that

$$\sum_{k=0}^n (-1)^k \dim(V_k) = 0.$$

**9.3.1. Definition.** It is frequently useful to think of functions as arrows in diagrams. For example, the situation  $j: R \rightarrow U$ ,  $f: R \rightarrow S$ ,  $k: S \rightarrow T$ ,  $h: U \rightarrow T$  may be represented by the following diagram.

$$\begin{array}{ccc} U & \xrightarrow{f} & X \\ \downarrow h & & \downarrow g \\ W & \xrightarrow{j} & V \end{array}$$

The diagram is said to COMMUTE (or to be a COMMUTATIVE DIAGRAM) if  $j \circ h = g \circ f$ .

- (g) Suppose that in the following diagram of vector spaces and linear maps

$$\begin{array}{ccccccc} \mathbf{0} & \longrightarrow & U & \xrightarrow{j} & V & \xrightarrow{k} & W \longrightarrow \mathbf{0} \\ & & \downarrow f & & \downarrow g & & \downarrow h \\ \mathbf{0} & \longrightarrow & U' & \xrightarrow{j'} & V' & \xrightarrow{k'} & W' \longrightarrow \mathbf{0} \end{array}$$

the rows are exact and the left square commutes. Then there exists a unique linear map  $h: W \rightarrow W'$  which makes the right square commute.

In parts (h)–(k) consider the diagram

$$\begin{array}{ccccccc}
 \mathbf{0} & \longrightarrow & U & \xrightarrow{j} & V & \xrightarrow{k} & W \longrightarrow \mathbf{0} \\
 & & \downarrow f & & \downarrow g & & \downarrow h \\
 \mathbf{0} & \longrightarrow & U' & \xrightarrow{j'} & V' & \xrightarrow{k'} & W' \longrightarrow \mathbf{0}
 \end{array}$$

where the rows are exact and the squares commute.

- (h) If  $g$  is surjective, so is  $h$ .
  - (i) If  $f$  is surjective and  $g$  is injective, then  $h$  is injective.
  - (j) If  $f$  and  $h$  are surjective, so is  $g$ .
  - (k) If  $f$  and  $h$  are injective, so is  $g$ .
- (6) Let  $V$  and  $W$  be vector spaces. Prove that (under the usual pointwise operations of addition and scalar multiplication)  $\mathfrak{L}(V, W)$  is a vector space.

**9.4. Answers to Odd-Numbered Exercises**

- (1) (a)  $-2, -1, 5, 5$   
(b)  $6, 3, -2$
- (3) line,  $-\frac{x}{5} = \frac{y}{3} = z$ , plane,  $2x + y - z = 0$
- (5) (a)  $7, 8, 5$   
(b)  $4, 21, 10$   
(c)  $3, 14, 5$   
(d)  $I$   
(e)  $2$
- (7) Pa, DK, H, DK, Pa, H, DO, VA, DL, Pb, VA, H, VA, DK, H, VA, DO, H
- (9)  $\sin t, \cos t, y'' + y = 0$
- (11) (a) line,  $\frac{x}{4} = -\frac{3y}{2} = z$   
(b) plane,  $2x - y - z = 0$
- (13)  $\mathbb{R}^4$  is the span of  $a, b, c$ , and  $d$ , all of which  $T$  takes to  $\mathbf{0}$ ; so were  $T$  linear, its range would contain only  $\mathbf{0}$ .
- (15)  $T$  does not map  $\mathbf{0}$  to  $\mathbf{0}$ .
- (17) If  $T$  were linear, then  $T(-2, -2)$  would be  $-2T(1, 1) = -2(3, -6) = (-6, 12)$ .

## CHAPTER 10

# LINEAR MAPS BETWEEN EUCLIDEAN SPACES

### 10.1. Background

**Topics:** linear mappings between finite dimensional spaces, a matrix as a linear map, the representation of a linear map as a matrix.

**10.1.1. Proposition.** *Let  $T \in \mathfrak{L}(V, W)$  where  $V$  is an  $n$ -dimensional vector space and  $W$  is an  $m$ -dimensional vector space and let  $\{\mathbf{e}^1, \mathbf{e}^2, \dots, \mathbf{e}^n\}$  be a basis for  $V$ . Define an  $m \times n$ -matrix  $[T]$  whose  $k^{\text{th}}$  column ( $1 \leq k \leq n$ ) is the column vector  $Te^k$ . Then for each  $x \in V$  we have*

$$Tx = [T]x.$$

The displayed equation above requires a little interpretation. The left side is  $T$  evaluated at  $x$ ; the right side is an  $m \times n$  matrix multiplied by an  $n \times 1$  matrix (that is, a column vector). Then the asserted equality can be thought of as identifying

- (1) two vectors in  $\mathbb{R}^m$ ,
- (2) two  $m$ -tuples of real numbers, or
- (3) two column vectors of length  $m$  (that is,  $m \times 1$  matrices).

If we wished to distinguish rigorously between column vectors and row vectors and also wished to identify  $m$ -tuples with row vectors, then the equation in the preceding proposition would have to read

$$Tx = ([T](x^t))^t.$$

To avoid the extra notation in these notes we will not make this distinction. In an equation interpret a vector as a row vector or as a column vector in any way that makes sense.

**10.1.2. Definition.** If  $V$  and  $W$  are finite dimensional vector spaces with bases and  $T \in \mathfrak{L}(V, W)$ , then the matrix  $[T]$  in the preceding proposition is the MATRIX REPRESENTATION of  $T$ . It is also called the STANDARD MATRIX for  $T$ .

## 10.2. Exercises

- (1) Let
- $T: \mathbb{R}^4 \rightarrow \mathbb{R}^3$
- be defined by

$$T\mathbf{x} = (x_1 - 3x_3 + x_4, 2x_1 + x_2 + x_3 + x_4, 3x_2 - 4x_3 + 7x_4)$$

for every  $\mathbf{x} = (x_1, x_2, x_3, x_4) \in \mathbb{R}^4$ . (The map  $T$  is linear, but you need not prove this.)

(a) Find  $[T]$ . Answer:  $\begin{bmatrix} \phantom{0} & \phantom{0} & \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} & \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} & \phantom{0} & \phantom{0} \end{bmatrix}$ .

(b) Find  $T(1, -2, 1, 3)$ . Answer: \_\_\_\_\_.

(c) Find  $([T]((1, -2, 1, 3)^t))^t$ . Answer: \_\_\_\_\_.

(d) Find  $\ker T$ . Answer:  $\ker T = \text{span}\{\text{_____}\}$ .

(e) Find  $\text{ran } T$ . Answer:  $\text{ran } T = \text{_____}$ .

- (2) Let
- $T: \mathbb{R}^3 \rightarrow \mathbb{R}^4$
- be defined by

$$T\mathbf{x} = (x_1 - 3x_3, x_1 + x_2 - 6x_3, x_2 - 3x_3, x_1 - 3x_3)$$

for every  $\mathbf{x} = (x_1, x_2, x_3) \in \mathbb{R}^3$ . (The map  $T$  is linear, but you need not prove this.) Then

(a)  $[T] = \begin{bmatrix} \phantom{0} & \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} & \phantom{0} \end{bmatrix}$ .

(b)  $T(3, -2, 4) = \text{_____}$ .

(c)  $\ker T = \text{span}\{\text{_____}\}$ .

(d)  $\text{ran } T = \text{span}\{\text{_____}\}$ .

- (3) Let  $\mathcal{P}_n$  be the vector space of all polynomial functions on  $\mathbb{R}$  with degree strictly less than  $n$ . The usual basis for  $\mathcal{P}_n$  is the set of polynomials  $1, t, t^2, t^3, \dots, t^{n-1}$ . Define  $T: \mathcal{P}_3 \rightarrow \mathcal{P}_5$  by

$$Tf(x) = \int_0^x \int_0^u p(t) dt$$

for all  $x, u \in \mathbb{R}$ .

- (a) Then the matrix representation of the linear map  $T$  (with respect to the usual

bases for  $\mathcal{P}_3$  and  $\mathcal{P}_5$  is  $\begin{bmatrix} \phantom{0} & \phantom{0} & \phantom{0} & \phantom{0} & \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} & \phantom{0} & \phantom{0} & \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} & \phantom{0} & \phantom{0} & \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} & \phantom{0} & \phantom{0} & \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} & \phantom{0} & \phantom{0} & \phantom{0} & \phantom{0} \end{bmatrix}$ .

(b) The kernel of  $T$  is \_\_\_\_\_.

(c) The range of  $T$  is  $\text{span}\{\text{_____}\}$ .



- (4) Let  $\mathcal{P}_4$  be the vector space of polynomials of degree strictly less than 4. Consider the linear transformation  $D^2: \mathcal{P}_4 \rightarrow \mathcal{P}_4: f \mapsto f''$ .

- (a) Then the matrix representation  $[D^2]$  of  $D^2$  (with respect to the usual basis  $\{1, t, t^2, t^3\}$ )

for  $\mathcal{P}_4$ ) is given by  $[D^2] = \begin{bmatrix} \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \end{bmatrix}$ .

- (b)  $\ker D^2 = \text{span}\{\underline{\hspace{2cm}}\}$  .

- (c)  $\text{ran } D^2 = \text{span}\{\underline{\hspace{2cm}}\}$  .

- (5) Let  $\mathcal{P}_4$  be the vector space of polynomials of degree strictly less than 4 and  $T: \mathcal{P}_4 \rightarrow \mathcal{P}_5$  be the linear transformation defined by  $(Tp)(t) = (2 + 3t)p(t)$  for every  $p \in \mathcal{P}_4$  and  $t \in \mathbb{R}$ . Then the matrix representation of  $T$  (with respect to the usual basis  $\{1, t, t^2, t^3\}$  for  $\mathcal{P}_4$ ) is given by

$$[T] = \begin{bmatrix} & & \\ & & \\ & & \end{bmatrix}.$$

- (6) Let  $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be the linear transformation whose standard matrix is  $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & -1 \end{bmatrix}$ . We

know that  $T$  is not onto because the only vectors  $(u, v, w)$  that are in the range of  $T$  are those that satisfy the relation  $u + av + bw = 0$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .

- (7) Let  $T$  be the linear map from  $\mathbb{R}^3$  to  $\mathbb{R}^3$  defined by

$$T(x, y, z) = (3x + 2y + z, x + 3z, -y + 4z).$$

- (a) The matrix representation of  $T$  is given by

$$[T] = \begin{bmatrix} & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \end{bmatrix}.$$

- (b) The range of  $T$  is (geometrically speaking) a \_\_\_\_\_ whose equation is

---

## 10.3. Problems

- (1) Define
- $T: \mathbb{R}^3 \rightarrow \mathbb{R}^2$
- by

$$T\mathbf{x} = (x_1 + 2x_2 - x_3, x_2 + x_3)$$

for all  $\mathbf{x} = (x_1, x_2, x_3)$  in  $\mathbb{R}^3$ .

- (a) Explain how to find  $[T]$ , the matrix representation for  $T$ .
- (b) Explain how to use  $[T]$  to find  $T(\mathbf{x})$  when  $\mathbf{x} = (-1, 2, -1)$ .
- (c) Explain how to use  $[T]$  to find a vector  $\mathbf{x}$  in  $\mathbb{R}^3$  such that  $T\mathbf{x} = (0, 1)$ .

Carry out the computations you describe.

- (2) Let
- $\mathcal{P}$
- be the vector space of all polynomial functions on
- $\mathbb{R}$
- with real coefficients. Define linear transformations
- $T, D: \mathcal{P} \rightarrow \mathcal{P}$
- by

$$(Dp)(x) = p'(x)$$

and

$$(Tp)(x) = x^2 p(x)$$

for all  $x \in \mathbb{R}$ .

Explain carefully how to find matrix representations for the linear transformations  $D+T$ ,  $DT$ , and  $TD$  (with respect to the usual basis  $\{1, t, t^2\}$  for the space of polynomials of degree two or less). Carry out the computations you describe. Use the resulting matrices to find  $((D+T)(p))(x)$ ,  $(DTp)(x)$ , and  $(TDp)(x)$  where  $p(x) = 3x^2 + 4x - 3$  for all  $x \in \mathbb{R}$ .

- (3) Define
- $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$
- by

$$T\mathbf{x} = (x_1 - x_2, x_2 - x_1, x_1 + x_2)$$

for all  $\mathbf{x} = (x_1, x_2)$  in  $\mathbb{R}^2$ .

- (a) Explain carefully how to find  $[T]$ , the matrix representation for  $T$ .
  - (b) How do we use  $[T]$  to find  $T(1, -2)$ ?
  - (c) Are there any nonzero vectors  $\mathbf{x}$  in  $\mathbb{R}^2$  such that  $T\mathbf{x} = \mathbf{0}$ ? Explain.
  - (d) Under what conditions is a vector  $(u, v, w)$  in the range of  $T$ ? Explain.
- (4) Let  $\mathcal{C}^1([0, 1])$  be the vector space of all functions defined on the interval  $[0, 1]$  which have continuous derivatives at each point and  $\mathcal{C}([0, 1])$  be the vector space of continuous functions on  $[0, 1]$ . Define a function  $T: \mathcal{C}^1([0, 1]) \rightarrow \mathcal{C}([0, 1])$  by

$$(Tf)(x) = \int_0^x f(t) dt + f'(x)$$

for every  $x \in [0, 1]$ .

- (a) Prove that the function  $T$  is linear.
  - (b) Let  $f(x) = \sin x$  and  $g(x) = \cos x$  for all  $x \in [0, 1]$ . Explain why one of these functions belongs to the kernel of  $T$  while the other does not.
- (5) Let  $\mathcal{P}_4$  be the vector space of polynomials of degree strictly less than 4. Consider the linear transformation  $D^2: \mathcal{P}_4 \rightarrow \mathcal{P}_4: f \mapsto f''$ . Explain carefully how to find  $[T]$ , the matrix representation of  $D^2$  (with respect to the usual basis  $\{1, t, t^2, t^3\}$  for  $\mathcal{P}_4$ ). Then explain how to use  $[T]$  to find  $\ker D^2$  and  $\text{ran } D^2$ .

**10.4. Answers to Odd-Numbered Exercises**

- (1) (a)  $\begin{bmatrix} 1 & 0 & -3 & 1 \\ 2 & 1 & 1 & 1 \\ 0 & 3 & -4 & 7 \end{bmatrix}$   
 (b)  $(1, 4, 11)$   
 (c)  $(1, 4, 11)$  (or  $[1 \ 4 \ 11]$ )  
 (d)  $(1, -9, 2, 5)$   
 (e)  $\mathbb{R}^3$

- (3) (a)  $\begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{6} & 0 \\ 0 & 0 & \frac{1}{12} \end{bmatrix}$

- (b)  $\{0\}$   
 (c)  $x^2, x^3, x^4$

- (5)  $\begin{bmatrix} 2 & 0 & 0 & 0 \\ 3 & 2 & 0 & 0 \\ 0 & 3 & 2 & 0 \\ 0 & 0 & 3 & 2 \\ 0 & 0 & 0 & 3 \end{bmatrix}$

- (7) (a)  $\begin{bmatrix} 3 & 2 & 1 \\ 1 & 0 & 3 \\ 0 & -1 & 4 \end{bmatrix}$   
 (b) plane,  $u - 3v + 2w = 0$



## CHAPTER 11

# PROJECTION OPERATORS

### 11.1. Background

**Topics:** projections along one subspace onto another.

**11.1.1. Definition.** Let  $V$  be a vector space and suppose that  $V = M \oplus N$ . We know that for each  $\mathbf{v} \in V$  there exist unique vectors  $\mathbf{m} \in M$  and  $\mathbf{n} \in N$  such that  $\mathbf{v} = \mathbf{m} + \mathbf{n}$  (see problem 5 in chapter 6). Define a function  $E_{MN}: V \rightarrow V$  by  $E_{MN}v = n$ . The function  $E_{MN}$  is the PROJECTION OF  $V$  ALONG  $M$  ONTO  $N$ . (Frequently we write  $E$  for  $E_{MN}$ . But keep in mind that  $E$  depends on both  $M$  and  $N$ .)

**11.1.2. Theorem.** Let  $V$  be a vector space and suppose that  $V = M \oplus N$ . If  $E$  is the projection of  $V$  along  $M$  onto  $N$ , then

- (i)  $E$  is linear;
- (ii)  $E^2 = E$  (that is,  $E$  is IDEMPOTENT);
- (iii)  $\text{ran } E = N$ ; and
- (iv)  $\ker E = M$ .

**11.1.3. Theorem.** Let  $V$  be a vector space and suppose that  $E: V \rightarrow V$  is a function which satisfies

- (i)  $E$  is linear, and
- (ii)  $E^2 = E$ .

Then

$$V = \ker E \oplus \text{ran } E$$

and  $E$  is the projection of  $V$  along  $\ker E$  onto  $\text{ran } E$ .

**11.1.4. Theorem.** Let  $V$  be a vector space and suppose that  $V = M \oplus N$ . If  $E$  is the projection of  $V$  along  $M$  onto  $N$ , then  $I - E$  is the projection of  $V$  along  $N$  onto  $M$ .

## 11.2. Exercises

- (1) Let  $M$  be the line  $y = 2x$  and  $N$  be the  $y$ -axis in  $\mathbb{R}^2$ . Then

$$[E_{MN}] = \begin{bmatrix} a & a \\ -b & c \end{bmatrix} \quad \text{and} \quad [E_{NM}] = \begin{bmatrix} c & a \\ b & a \end{bmatrix}$$

where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ , and  $c = \underline{\hspace{1cm}}$ .

- (2) Let  $P$  be the plane in  $\mathbb{R}^3$  whose equation is  $x - z = 0$  and  $L$  be the line whose equations are  $y = 0$  and  $x = -z$ . Let  $E$  be the projection of  $\mathbb{R}^3$  along  $L$  onto  $P$  and  $F$  be the projection of  $\mathbb{R}^3$  along  $P$  onto  $L$ . Then

$$[E] = \begin{bmatrix} a & b & a \\ b & c & b \\ a & b & a \end{bmatrix} \quad \text{and} \quad [F] = \begin{bmatrix} a & b & -a \\ b & b & b \\ -a & b & a \end{bmatrix}$$

where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ , and  $c = \underline{\hspace{1cm}}$ .

- (3) Let  $P$  be the plane in  $\mathbb{R}^3$  whose equation is  $x + 2y - z = 0$  and  $L$  be the line whose equations are  $\frac{x}{3} = y = \frac{z}{2}$ . Let  $E$  be the projection of  $\mathbb{R}^3$  along  $L$  onto  $P$  and  $F$  be the projection of  $\mathbb{R}^3$  along  $P$  onto  $L$ . Then

$$[E] = \frac{1}{3} \begin{bmatrix} a & -b & c \\ -d & d & d \\ a - 2d & -b + 2d & c + 2d \end{bmatrix} \quad \text{and} \quad [F] = \frac{1}{3} \begin{bmatrix} 3d & 3e & -3d \\ d & e & -d \\ 2d & 2e & -2d \end{bmatrix}$$

where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ ,  $c = \underline{\hspace{1cm}}$ ,  $d = \underline{\hspace{1cm}}$ , and  $e = \underline{\hspace{1cm}}$ .

- (4) Let  $P$  be the plane in  $\mathbb{R}^3$  whose equation is  $x - y - 2z = 0$  and  $L$  be the line whose equations are  $x = 0$  and  $y = -z$ . Let  $E$  be the projection of  $\mathbb{R}^3$  along  $L$  onto  $P$  and  $F$  be the projection of  $\mathbb{R}^3$  along  $P$  onto  $L$ . Then

$$[E] = \begin{bmatrix} a & b & b \\ -a & c & c \\ a & -a & -a \end{bmatrix} \quad \text{and} \quad [F] = \begin{bmatrix} b & b & b \\ a & -a & -c \\ -a & a & c \end{bmatrix}$$

where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ , and  $c = \underline{\hspace{1cm}}$ .

- (5) Let  $E$  be the projection of  $\mathbb{R}^3$  along the  $z$ -axis onto the plane  $3x - y + 2z = 0$  and let  $F$  be the projection of  $\mathbb{R}^3$  along the plane  $3x - y + 2z = 0$  onto the  $z$ -axis.

(a) Then  $[E] = \begin{bmatrix} & & \\ & & \\ & & \end{bmatrix}$ .

(b) Where does  $F$  take the point  $(4, 5, 1)$ ? Answer:  $(\underline{\hspace{1cm}}, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})$ .

- (6) Let  $M$  be the  $y$ -axis and  $N$  be the plane  $x + y - 2z = 0$  in  $\mathbb{R}^3$ .

(a) Then the projection  $E_{MN}$  of  $\mathbb{R}^3$  along  $M$  onto  $N$  is  $\begin{bmatrix} & & \\ & & \\ & & \end{bmatrix}$ .

(b) The projection  $E_{NM}$  takes the vector  $(3, 2, 1)$  to  $(\underline{\hspace{1cm}}, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})$ .

**11.3. Problems**

- (1) Let  $E$  be a projection on a vector space. Show that a vector  $\mathbf{x}$  belongs to the range of  $E$  if and only if  $E\mathbf{x} = \mathbf{x}$ . *Hint.* Recall (from Theorems 11.1.2 and 11.1.3) that a projection is a linear map  $E$  such that  $E^2 = E$ .
- (2) Prove Theorem 11.1.2.
- (3) Prove Theorem 11.1.3.
- (4) Prove Theorem 11.1.4. *Hint.* Use Theorem 11.1.3.
- (5) Let  $P$  be the plane in  $\mathbb{R}^3$  whose equation is  $x - z = 0$  and  $L$  be the line whose equations are  $y = 0$  and  $x = \frac{1}{2}z$ . Explain carefully how to find the matrix representation of the operator  $E_{LP}$ , that is, the projection of  $\mathbb{R}^3$  along  $L$  onto  $P$ . Carry out the computation you describe.
- (6) Let  $L$  be the line in  $\mathbb{R}^3$  whose equations are  $x = y$  and  $z = 0$ , and let  $P$  be the plane whose equation is  $x - z = 0$ . Explain carefully how to find the matrix representation of the operator  $E_{LP}$ , that is, the projection of  $\mathbb{R}^3$  along  $L$  onto  $P$ . Carry out the computation you describe.
- (7) Let  $P$  be the plane in  $\mathbb{R}^3$  whose equation is  $x - 3y + z = 0$  and  $L$  be the line whose equations are  $x = -2y = -z$ . Explain carefully how to find the matrix representation of the operator  $E_{LP}$  of  $\mathbb{R}^3$  along  $L$  onto  $P$  and the projection  $E_{PL}$  of  $\mathbb{R}^3$  along  $P$  onto  $L$ .
- (8) Prove that a linear transformation between vector spaces has a left inverse if and only if it is injective.
- (9) Prove that a linear transformation between vector spaces has a right inverse if and only if it is surjective.

**11.4. Answers to Odd-Numbered Exercises**

(1) 0, 2, 1

(3) 0, 6, 3, 1, 2

(5) (a)  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -\frac{3}{2} & \frac{1}{2} & 0 \end{bmatrix}$

(b) 0, 0,  $\frac{9}{2}$



**Part 4**

**SPECTRAL THEORY OF VECTOR  
SPACES**



## EIGENVALUES AND EIGENVECTORS

### 12.1. Background

**Topics:** characteristic polynomial, eigenvalues, eigenvectors, eigenspaces, algebraic multiplicity and geometric multiplicity of an eigenvalue.

**12.1.1. Definition.** A number  $\lambda$  is an **EIGENVALUE** of an operator  $T$  on a vector space  $V$  if  $\ker(\lambda I_V - T)$  contains a nonzero vector. Any such vector is an **EIGENVECTOR** of  $T$  associated with  $\lambda$  and  $\ker(\lambda I_V - T)$  is the **EIGENSPACE** of  $T$  associated with  $\lambda$ . The set of all eigenvalues of the operator  $T$  is its **(POINT) SPECTRUM** and is denoted by  $\sigma(T)$ .

If  $M$  is an  $n \times n$  matrix, then  $\det(\lambda I_n - M)$  (where  $I_n$  is the  $n \times n$  identity matrix) is a polynomial in  $\lambda$  of degree  $n$ . This is the **CHARACTERISTIC POLYNOMIAL** of  $M$ . A standard way of computing the eigenvalues of an operator  $T$  on a finite dimensional vector space is to find the zeros of the characteristic polynomial of its matrix representation. It is an easy consequence of the multiplicative property of the determinant function (see proposition 3.1.9) that the characteristic polynomial of an operator  $T$  on a vector space  $V$  is independent of the basis chosen for  $V$  and hence of the particular matrix representation of  $T$  that is used.

**12.1.2. Theorem** (Spectral Mapping Theorem). *If  $T$  is an operator on a finite dimensional vector space and  $p$  is a polynomial, then*

$$\sigma(p(T)) = p(\sigma(T)).$$

*That is, if  $\sigma(T) = \{\lambda_1, \dots, \lambda_k\}$ , then  $\sigma(p(T)) = \{p(\lambda_1), \dots, p(\lambda_k)\}$ .*

## 12.2. Exercises

- (1) If  $A = \begin{bmatrix} 1 & -1 & 4 \\ 3 & 2 & -1 \\ 2 & 1 & -1 \end{bmatrix}$ , then the eigenvalue \_\_\_\_\_ has corresponding eigenvector ( \_\_\_\_\_ , 1 , 1 ), the eigenvalue \_\_\_\_\_ has corresponding eigenvector ( \_\_\_\_\_ , 4 , 1 ), and the eigenvalue \_\_\_\_\_ has corresponding eigenvector ( \_\_\_\_\_ , 2 , 1 ).
- (2) Let  $A = \begin{bmatrix} 0 & 0 & 2 \\ 0 & 2 & 0 \\ 2 & 0 & 0 \end{bmatrix}$ .
- The eigenvalues of  $A$  are \_\_\_\_\_ , \_\_\_\_\_ , and \_\_\_\_\_ .
  - The matrix  $A$  has a one-dimensional eigenspace.  
It is the span of the vector ( 1 , \_\_\_\_\_ , \_\_\_\_\_ ).
  - The matrix  $A$  has a two-dimensional eigenspace.  
It is the span of the vectors ( 1 , 0 , \_\_\_\_\_ ) and ( 0 , 1 , \_\_\_\_\_ ).
- (3) Choose  $a$ ,  $b$  and  $c$  in the matrix  $A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ a & b & c \end{bmatrix}$  so that the characteristic polynomial of  $A$  is  $-\lambda^3 + 4\lambda^2 + 5\lambda + 6$ .  
Answer:  $a =$  \_\_\_\_\_ ;  $b =$  \_\_\_\_\_ ; and  $c =$  \_\_\_\_\_ .
- (4) Suppose that it is known that the matrix  $A = \begin{bmatrix} 1 & 0 & -1 \\ \sqrt{3} & a & 17 \\ 2 & 0 & b \end{bmatrix}$  has eigenvalues 2 and 3 and that the eigenvalue 2 has algebraic multiplicity 2. Then  $a =$  \_\_\_\_\_ and  $b =$  \_\_\_\_\_ .
- (5) The matrices  $A = \begin{bmatrix} a & 1 \\ -2 & d \end{bmatrix}$  and  $B = \frac{1}{25} \begin{bmatrix} 114 & 48 \\ 48 & 86 \end{bmatrix}$  have the same eigenvalues.  
Then  $a =$  \_\_\_\_\_ and  $d =$  \_\_\_\_\_ .
- (6) Let  $A = \begin{bmatrix} 3 & 4 & 2 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}$ .
- The eigenvalues of  $A$  are \_\_\_\_\_ , \_\_\_\_\_ , and \_\_\_\_\_ .
  - The matrix  $A$  has three one-dimensional eigenspaces. They are spanned by the vectors ( \_\_\_\_\_ , \_\_\_\_\_ , \_\_\_\_\_ ), ( \_\_\_\_\_ , \_\_\_\_\_ , \_\_\_\_\_ ), and ( \_\_\_\_\_ , \_\_\_\_\_ , \_\_\_\_\_ ), respectively.
- (7) Let  $A = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix}$ .
- The eigenvalues of  $A - I$  are \_\_\_\_\_ (which has algebraic multiplicity \_\_\_\_\_ ) and \_\_\_\_\_ (which has algebraic multiplicity \_\_\_\_\_ ).
  - The determinant of  $A - I$  is \_\_\_\_\_ .
- (8) Let  $T$  be the operator on  $\mathbb{R}^3$  whose matrix representation is  $\begin{bmatrix} 1 & -1 & 0 \\ 0 & 0 & 0 \\ -2 & 2 & 2 \end{bmatrix}$ . Then the eigenvalues of the operator  $T^5 - 3T^4 + T^3 - T^2 + T - 3I$  are \_\_\_\_\_ , \_\_\_\_\_ , and \_\_\_\_\_ .

**12.3. Problems**

- (1) Suppose that  $A$  and  $B$  are  $n \times n$  matrices. Prove that  $\sigma(AB) = \sigma(BA)$ . *Hint.* Show that if  $\lambda$  is an eigenvalue of  $AB$ , then it is also an eigenvalue of  $BA$ . Deal with the cases  $\lambda = 0$  and  $\lambda \neq 0$  separately.
- (2) Let  $c \in \mathbb{R}$ . Suppose that  $A$  is an  $n \times n$  matrix and that the sum of the entries in each column of  $A$  is  $c$ . Prove that  $c$  is an eigenvalue of  $A$ . *Hint.* Consider the sum of the row vectors of the matrix  $A - cI$ .
- (3) This is a problem in cryptography. Read about Hill ciphers, then decode the following Hill 3-cipher given that the first two words of the plaintext are known to be “My candle”. (See for example [1], section 11.16.)

OGWCGWGKK.EWVD.XZJOHZWLNYH USTFAIOS.A.KBN  
JRCENYQZV,IE LTGCGWGKC YYBLSDWWODLBVFFOS.H

In many discussions of Hill ciphers letters of the alphabet are assigned numbers from 0 to 25 and arithmetic is done modulo 26. The encoding here is done slightly differently. Here each letter is assigned its numerical position in the alphabet (including Z which is assigned 26). Furthermore, a space between words is assigned 27, a comma is assigned 28, and a period is assigned zero. Thus, for this code, all arithmetic should be done modulo 29. (One reason for this is that some computer algebra systems have problems calculating inverses mod 26.) Note: the ciphertext contains exactly three spaces.

**12.4. Answers to Odd-Numbered Exercises**

(1)  $-2, -1, 1, -1, 3, 1$

(3)  $6, 5, 4$

(5)  $2, 6$  (or  $6, 2$ )

(7) (a)  $-1, 3, 3, 1$

(b)  $-3$

## DIAGONALIZATION OF MATRICES

### 13.1. Background

**Topics:** similarity of matrices, triangular and diagonal matrices, diagonalization, annihilating and minimal polynomials, algebraic and geometric multiplicity of an eigenvalue, the *Cayley-Hamilton theorem*.

**13.1.1. Definition.** Two operators  $R$  and  $T$  on a vector space  $V$  are **SIMILAR** if there exists an invertible operator  $S$  on  $V$  such that  $R = S^{-1}TS$ .

**13.1.2. Proposition.** If  $V$  is a vector space, then similarity is an equivalence relation on  $\mathfrak{L}(V)$ .

**13.1.3. Definition.** Let  $V$  be a finite dimensional vector space and  $B = \{e^1, \dots, e^n\}$  be a basis for  $V$ . An operator  $T$  on  $V$  is **DIAGONAL** if there exist scalars  $\alpha_1, \dots, \alpha_n$  such that  $Te^k = \alpha_k e^k$  for each  $k \in \mathbb{N}_n$ . Equivalently,  $T$  is diagonal if its matrix representation  $[T] = [t_{ij}]$  has the property that  $t_{ij} = 0$  whenever  $i \neq j$ .

Asking whether a particular operator on some finite dimensional vector space is diagonal is, strictly speaking, nonsense. As defined the operator property of being diagonal is definitely *not* a vector space concept. It makes sense only for a vector space *for which a basis has been specified*. This important, if obvious, fact seems to go unnoticed in many beginning linear algebra texts, due, I suppose, to a rather obsessive fixation on  $\mathbb{R}^n$  in such courses. Here is the relevant *vector space* property.

**13.1.4. Definition.** An operator  $T$  on a finite dimensional vector space  $V$  is **DIAGONALIZABLE** if there exists a basis for  $V$  with respect to which  $T$  is diagonal. Equivalently, an operator on a finite dimensional vector space *with basis* is diagonalizable if it is similar to a diagonal operator. If a matrix  $D$  is diagonalizable and  $\Lambda = S^{-1}DS$  is diagonal, we say that the matrix  $S$  **DIAGONALIZES**  $D$ .

**13.1.5. Theorem.** Let  $A$  be an  $n \times n$  matrix with  $n$  linear independent eigenvectors. If  $S$  is a matrix with these eigenvectors as columns, then  $S$  diagonalizes  $A$ . The entries along the diagonal of the resulting diagonal matrix are all eigenvalues of  $A$ .

**13.1.6. Definition.** A polynomial is **MONIC** if its leading coefficient is 1. Thus a polynomial of degree  $n$  is monic if it takes the form  $x^n + a_{n-1}x^{n-1} + \dots + a_1x + a_0$ .

**13.1.7. Definition.** Let  $p$  be a polynomial of degree at least one and  $T$  be an operator on some vector space. We say that  $p$  is an **ANNIHILATING POLYNOMIAL** for  $T$  (or that  $p$  **ANNIHILATES**  $T$ ) if  $p(T) = 0$ . For example, if  $T^3 - 4T^2 + T - 7I = 0$ , then the polynomial  $p$  defined by  $p(x) = x^3 - 4x^2 + x - 7$  is an annihilating polynomial for  $T$ .

**13.1.8. Definition.** Let  $T$  be an operator on a finite dimensional vector space. The **MINIMAL POLYNOMIAL** of  $T$  is the unique monic polynomial of smallest degree which annihilates  $T$ . (It is left as a problem to verify the existence and the uniqueness of such a polynomial: see problem 8.)

**13.1.9. Theorem** (Cayley-Hamilton Theorem). On a finite dimensional vector space the characteristic polynomial of an operator  $T$  annihilates  $T$ .

**Paraphrase:** Every matrix satisfies its characteristic equation.

**13.1.10. Definition.** A square matrix  $A = [a_{ij}]$  is UPPER TRIANGULAR if  $a_{ij} = 0$  whenever  $i > j$ . A matrix is TRIANGULABLE (or TRIANGULIZABLE) if it is similar to an upper triangular matrix.

**13.1.11. Theorem.** Let  $T$  be an operator on a finite dimensional vector space and let  $\{\lambda_1, \dots, \lambda_k\}$  be its distinct eigenvalues. Then:

- (1)  $T$  is triangulable if and only if its minimal polynomial can be factored into a product of linear factors. That is, if and only if there are positive integers  $r_1, \dots, r_k$  such that

$$m_T(x) = (x - \lambda_1)^{r_1} \dots (x - \lambda_k)^{r_k}.$$

- (2)  $T$  is diagonalizable if and only if its minimal polynomial has the form

$$m_T(x) = (x - \lambda_1) \dots (x - \lambda_k).$$

**13.1.12. Corollary.** Every operator on a complex finite dimensional vector space is triangulable.

**13.1.13. Definition.** An operator is NILPOTENT if some power of the operator is  $\mathbf{0}$ .



## 13.2. Exercises

- (1) Let  $A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}$ . The characteristic polynomial of  $A$  is  $\lambda^p(\lambda - 3)^q$  where  $p = \underline{\hspace{2cm}}$  and  $q = \underline{\hspace{2cm}}$ .

The minimal polynomial of  $A$  is  $\lambda^r(\lambda - 3)^s$  where  $r = \underline{\hspace{2cm}}$  and  $s = \underline{\hspace{2cm}}$ .

- (2) Let  $T$  be the operator on  $\mathbb{R}^4$  whose matrix representation is  $\begin{bmatrix} 0 & 1 & 0 & -1 \\ -2 & 3 & 0 & -1 \\ -2 & 1 & 2 & -1 \\ 2 & -1 & 0 & 3 \end{bmatrix}$ .

The characteristic polynomial of  $T$  is  $(\lambda - 2)^p$  where  $p = \underline{\hspace{2cm}}$ .

The minimal polynomial of  $T$  is  $(\lambda - 2)^r$  where  $r = \underline{\hspace{2cm}}$ .

- (3) Let  $T$  be the operator on  $\mathbb{R}^3$  whose matrix representation is  $\begin{bmatrix} 3 & 1 & -1 \\ 2 & 2 & -1 \\ 2 & 2 & 0 \end{bmatrix}$ .

- (a) Find the characteristic polynomial of  $T$ .

Answer:  $c_T(\lambda) = (\lambda - 1)^p(\lambda - 2)^q$  where  $p = \underline{\hspace{2cm}}$  and  $q = \underline{\hspace{2cm}}$ .

- (b) Find the minimal polynomial of  $T$ .

Answer:  $m_T(\lambda) = (\lambda - 1)^r(\lambda - 2)^s$  where  $r = \underline{\hspace{2cm}}$  and  $s = \underline{\hspace{2cm}}$ .

- (c) Find the eigenspaces  $M_1$  and  $M_2$  of  $T$ .

Answer:  $M_1 = \text{span} \{ \underline{\hspace{4cm}} \}$  and  $M_2 = \text{span} \{ \underline{\hspace{4cm}} \}$ .

- (4) Let  $T$  be the operator on  $\mathbb{R}^5$  whose matrix representation is  $\begin{bmatrix} 1 & 0 & 0 & 1 & -1 \\ 0 & 1 & -2 & 3 & -3 \\ 0 & 0 & -1 & 2 & -2 \\ 1 & -1 & 1 & 0 & 1 \\ 1 & -1 & 1 & -1 & 2 \end{bmatrix}$ .

- (a) Find the characteristic polynomial of  $T$ .

Answer:  $c_T(\lambda) = (\lambda + 1)^p(\lambda - 1)^q$  where  $p = \underline{\hspace{2cm}}$  and  $q = \underline{\hspace{2cm}}$ .

- (b) Find the minimal polynomial of  $T$ .

Answer:  $m_T(\lambda) = (\lambda + 1)^r(\lambda - 1)^s$  where  $r = \underline{\hspace{2cm}}$  and  $s = \underline{\hspace{2cm}}$ .

- (c) Find the eigenspaces  $V_1$  and  $V_2$  of  $T$ .

Answer:  $V_1 = \text{span} \{ \underline{\hspace{4cm}} \}$  and

$V_2 = \text{span} \{ \underline{\hspace{4cm}} \}$ .

- (5) Let  $T$  be an operator whose matrix representation is  $\begin{bmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 3 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & -1 & -1 \end{bmatrix}$ .

- (a) Regarding  $T$  as an operator on  $\mathbb{R}^5$  find its characteristic polynomial and minimal polynomial.

Answer:  $c_T(\lambda) = \lambda^p(\lambda^2 + 1)^q$  where  $p = \underline{\hspace{2cm}}$  and  $q = \underline{\hspace{2cm}}$ .

and  $m_T(\lambda) = \lambda^r(\lambda^2 + 1)^s$  where  $r = \underline{\hspace{2cm}}$  and  $s = \underline{\hspace{2cm}}$ .

- (b) Regarded as an operator on  $\mathbb{R}^5$  is  $T$  diagonalizable?  $\underline{\hspace{2cm}}$ . Is it triangulable?  $\underline{\hspace{2cm}}$ .

(c) Regarded as an operator on  $\mathbb{C}^5$  is  $T$  diagonalizable? \_\_\_\_\_. Is it triangulable? \_\_\_\_\_.

(6) Let  $T$  be the operator on  $\mathbb{R}^3$  whose matrix representation is  $\begin{bmatrix} 2 & 0 & 0 \\ -1 & 3 & 2 \\ 1 & -1 & 0 \end{bmatrix}$ .

(a) Find the characteristic and minimal polynomials of  $T$ .

Answer:  $c_T(\lambda) = (\lambda - 1)^p(\lambda - 2)^q$  where  $p = \underline{\hspace{1cm}}$  and  $q = \underline{\hspace{1cm}}$ .

and  $m_T(\lambda) = (\lambda - 1)^r(\lambda - 2)^s$  where  $r = \underline{\hspace{1cm}}$  and  $s = \underline{\hspace{1cm}}$ .

(b) What can be concluded from the form of the minimal polynomial?

Answer: \_\_\_\_\_

(c) Find a matrix  $S$  (if one exists) that diagonalizes  $[T]$ . What is the diagonal form  $\Lambda$  of

$[T]$  produced by this matrix? Answer:  $S = \begin{bmatrix} a & b & a \\ b & b & -c \\ -b & a & b \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ ,

and  $c = \underline{\hspace{1cm}}$ ; and  $\Lambda = \begin{bmatrix} \lambda & 0 & 0 \\ 0 & \mu & 0 \\ 0 & 0 & \mu \end{bmatrix}$  where  $\lambda = \underline{\hspace{1cm}}$  and  $\mu = \underline{\hspace{1cm}}$ .

(7) Let  $T$  be the operator on  $\mathbb{R}^3$  whose matrix representation is  $\begin{bmatrix} 8 & -6 & 12 \\ -18 & 11 & 18 \\ -6 & -3 & 26 \end{bmatrix}$ .

(a) Find the characteristic and minimal polynomials of  $T$ .

Answer:  $c_T(\lambda) = (\lambda - 5)^p(\lambda - 20)^q$  where  $p = \underline{\hspace{1cm}}$  and  $q = \underline{\hspace{1cm}}$ .

and  $m_T(\lambda) = (\lambda - 5)^r(\lambda - 20)^s$  where  $r = \underline{\hspace{1cm}}$  and  $s = \underline{\hspace{1cm}}$ .

(b) What can be concluded from the form of the minimal polynomial?

Answer: \_\_\_\_\_

(c) Find a matrix  $S$  (if one exists) that diagonalizes  $[T]$ . What is the diagonal form  $\Lambda$  of

$[T]$  produced by this matrix? Answer:  $S = \begin{bmatrix} a & b & c \\ d & -a & a \\ b & c & b \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ ,

$c = \underline{\hspace{1cm}}$ , and  $d = \underline{\hspace{1cm}}$ ; and  $\Lambda = \begin{bmatrix} \lambda & 0 & 0 \\ 0 & \mu & 0 \\ 0 & 0 & \mu \end{bmatrix}$  where  $\lambda = \underline{\hspace{1cm}}$  and  $\mu = \underline{\hspace{1cm}}$ .

(8) Let  $\mathcal{P}_n$  be the space of polynomials of degree strictly less than  $n$  and  $D$  be the differentiation operator on  $\mathcal{P}_n$ . Then

(a) the only eigenvalue of  $D$  is  $\lambda = \underline{\hspace{1cm}}$ ;

(b) the corresponding eigenspace is the span of \_\_\_\_\_;

(c) the algebraic multiplicity of  $\lambda$  is \_\_\_\_\_; and

(d) the geometric multiplicity of  $\lambda$  is \_\_\_\_\_.

## 13.3. Problems

- (1) Prove that the trace function is a similarity invariant on the family of  $n \times n$  matrices; that is, prove that if  $A$  and  $B$  are similar  $n \times n$  matrices, then  $\operatorname{tr} A = \operatorname{tr} B$ . *Hint.* Prove first that if  $M$  and  $N$  are  $n \times n$  matrices, then  $MN$  and  $NM$  have the same trace.
- (2) Prove that the determinant function is a similarity invariant on the family of  $n \times n$  matrices; that is, prove that if  $A$  and  $B$  are similar  $n \times n$  matrices, then  $\det A = \det B$ .
- (3) Prove that if two matrices are diagonalized by the same matrix, then they commute.
- (4) Prove that if a matrix  $A$  is diagonalizable, then so is every matrix similar to  $A$ .
- (5) Show that if  $A$  is an  $n \times n$  matrix of real (or complex) numbers, then  $\operatorname{tr} A$  is the sum of the eigenvalues of  $A$  and  $\det A$  is their product.
- (6) Suppose that  $T$  is an operator on a finite dimensional complex vector space and that  $\sigma(T) = \{0\}$ . Show that  $T$  is nilpotent.
- (7) Let  $T$  be an operator on a finite dimensional vector space.

- (a) Show that if  $p$  is an annihilating polynomial for  $T$ , then the minimal polynomial  $m_T$  divides  $p$ . *Hint.* Suppose that  $p$  annihilates  $T$  (so that the degree of  $p$  is at least as large as the degree of  $m_T$ ). Divide  $p$  by  $m_T$ . Then there exist polynomials  $q$  (the quotient) and  $r$  (the remainder) such that

$$p = m_T q + r \quad \text{and} \quad \text{degree of } r < \text{degree of } m_T.$$

Conclude that  $r = 0$ .

- (b) Show that the minimal polynomial  $m_T$  and the characteristic polynomial  $c_T$  have exactly the same roots (although not necessarily the same multiplicities). *Hint.* To show that every root of  $m_T$  is also a root of  $c_T$ , it is enough to know that  $m_T$  divides  $c_T$ . Why is that true?

To obtain the converse, suppose that  $\lambda$  is a root of  $c_T$ : that is, suppose that  $\lambda$  is an eigenvalue of  $T$ . Use the *spectral mapping theorem* 12.1.2 to show that  $m_T(\lambda) = 0$ .

- (8) Let  $T$  be an operator on a finite dimensional vector space  $V$ . Show that there is a unique monic polynomial of smallest degree which annihilates  $T$ . *Hint.* This asks for a proof of the existence and the uniqueness of the *minimal polynomial* for the operator  $T$ . The existence part is easy: If there are any polynomials at all which annihilate  $T$ , surely there is one of smallest degree. (How do we know that there is at least one polynomial that annihilates  $T$ ?) We want the annihilating polynomial of smallest degree to be monic—is this a problem?

To verify the uniqueness of the minimal polynomial, consider the case of degree one separately. That is, suppose that  $p$  and  $q$  are monic annihilating polynomials of degree one. How do we know that  $p = q$ ? Then consider polynomials of higher degree. Suppose that  $p$  and  $q$  are monic annihilating polynomials of smallest degree  $k$  where  $k > 1$ . What can you say about  $p - q$ ?

**13.4. Answers to Odd-Numbered Exercises**

- (1) 2, 1, 1, 1
- (3) (a) 1, 2  
(b) 1, 2  
(c) (1, 0, 2), (1, 1, 2)
- (5) (a) 3, 1, 2, 1  
(b) no, no  
(c) no, yes
- (7) (a) 1, 2, 1, 1  
(b)  $T$  is diagonalizable  
(c) 2, 1, 0, 3, 5, 20

## CHAPTER 14

# SPECTRAL THEOREM FOR VECTOR SPACES

### 14.1. Background

**Topics:** the *spectral theorem for finite dimensional vector spaces* (writing a diagonalizable operator as a linear combination of projections).

The central fact asserted by the finite dimensional vector space version of the *spectral theorem* is that every diagonalizable operator on such a space can be written as a linear combination of projection operators where the coefficients of the linear combination are the eigenvalues of the operator and the ranges of the projections are the corresponding eigenspaces. Here is a formal statement of the theorem.

**14.1.1. Theorem** (Spectral theorem: vector space version). *Let  $T$  be a diagonalizable operator on a finite dimensional vector space  $V$ , and  $\lambda_1, \dots, \lambda_k$  be the (distinct) eigenvalues of  $T$ . For each  $j$  let  $M_j$  be the eigenspace associated with  $\lambda_j$  and  $E_j$  be the projection of  $V$  onto  $M_j$  along  $M_1 + \dots + M_{j-1} + M_{j+1} + \dots + M_k$ . Then*

- (i)  $T = \lambda_1 E_1 + \dots + \lambda_k E_k$ ,
- (ii)  $I = E_1 + \dots + E_k$ , and
- (iii)  $E_i E_j = 0$  when  $i \neq j$ .

For a proof of this result see, for example, [8], page 215, theorem 11.

## 14.2. Exercises

- (1) Let  $T$  be the operator on  $\mathbb{R}^2$  whose matrix representation is  $\begin{bmatrix} -7 & 8 \\ -16 & 17 \end{bmatrix}$ .

- (a) Find the characteristic polynomial and minimal polynomial for  $T$ .

Answer:  $c_T(\lambda) = \underline{\hspace{2cm}}$  and  $m_T(\lambda) = \underline{\hspace{2cm}}$ .

- (b) The eigenspace  $M_1$  associated with the smaller eigenvalue  $\lambda_1$  is the span of  $(1, \underline{\hspace{1cm}})$ .

- (c) The eigenspace  $M_2$  associated with the larger eigenvalue  $\lambda_2$  is the span of  $(1, \underline{\hspace{1cm}})$ .

- (d) We wish to write  $T$  as a linear combination of projection operators. Find the (matrix representations of the) appropriate projections  $E_1$  and  $E_2$  onto the eigenspaces  $M_1$  and  $M_2$ , respectively.

Answer:  $E_1 = \begin{bmatrix} a & b \\ a & b \end{bmatrix}$ , where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ , and  $E_2 = \begin{bmatrix} -c & c \\ -d & d \end{bmatrix}$ , where  $c = \underline{\hspace{1cm}}$  and  $d = \underline{\hspace{1cm}}$ .

- (e) Compute the sum and product of  $E_1$  and  $E_2$ .

Answer:  $E_1 + E_2 = \begin{bmatrix} \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} \end{bmatrix}$ ; and  $E_1 E_2 = \begin{bmatrix} \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} \end{bmatrix}$ .

- (f) Write  $T$  as a linear combination of the projections found in (d).

Answer:  $[T] = \underline{\hspace{1cm}} E_1 + \underline{\hspace{1cm}} E_2$ .

- (g) Find a matrix  $S$  which diagonalizes  $[T]$ . What is the associated diagonal form  $\Lambda$  of  $[T]$ ?

Answer:  $S = \begin{bmatrix} 1 & 1 \\ a & b \end{bmatrix}$ , where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ , and  $\Lambda = \begin{bmatrix} \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} \end{bmatrix}$ .

- (2) Let  $T$  be the operator on  $\mathbb{R}^3$  whose matrix representation is

$$\begin{bmatrix} -2 & -1 & -6 \\ -6 & -1 & -12 \\ 2 & 1 & 6 \end{bmatrix}.$$

- (a) Find the characteristic polynomial and minimal polynomial for  $T$ .

Answer:  $c_T(\lambda) = \underline{\hspace{2cm}}$  and  $m_T(\lambda) = \underline{\hspace{2cm}}$ .

- (b) The eigenspace  $M_1$  associated with the smallest eigenvalue  $\lambda_1$  is the span of  $(3, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})$ .

- (c) The eigenspace  $M_2$  associated with the middle eigenvalue  $\lambda_2$  is the span of  $(1, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})$ .

- (d) The eigenspace  $M_3$  associated with the largest eigenvalue  $\lambda_3$  is the span of  $(1, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})$ .

- (e) We wish to write  $T$  as a linear combination of projection operators. Find the (matrix representations of the) appropriate projections  $E_1$ ,  $E_2$ , and  $E_3$  onto the eigenspaces  $M_1$ ,  $M_2$ , and  $M_3$ , respectively.

Answer:  $E_1 = \begin{bmatrix} a & c & a \\ 2a & c & 2a \\ -b & c & -b \end{bmatrix}$ ;  $E_2 = \begin{bmatrix} -b & d & c \\ -2a & a & c \\ b & -d & c \end{bmatrix}$ ; and  $E_3 = \begin{bmatrix} c & -d & -a \\ c & -b & -2a \\ c & d & a \end{bmatrix}$ , where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ ,  $c = \underline{\hspace{1cm}}$ , and  $d = \underline{\hspace{1cm}}$ .

- (f) Write  $T$  as a linear combination of the projections found in (e).

Answer:  $[T] = \underline{\hspace{1cm}} E_1 + \underline{\hspace{1cm}} E_2 + \underline{\hspace{1cm}} E_3$ .

- (g) Find a matrix  $S$  which diagonalizes  $[T]$ . What is the associated diagonal form  $\Lambda$  of  $[T]$ ?

Answer:  $S = \begin{bmatrix} a & b & b \\ 2a & a & c \\ -c & -b & -b \end{bmatrix}$ , where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ , and  $c = \underline{\hspace{1cm}}$ , and

$$\Lambda = \begin{bmatrix} & & \\ & & \\ & & \end{bmatrix}.$$

- (3) Find a matrix  $A$  whose eigenvalues are 1 and 4, and whose eigenvectors are  $(3, 1)$  and  $(2, 1)$ , respectively.

Answer:  $A = \begin{bmatrix} & \\ & \end{bmatrix}.$

**14.3. Answers to Odd-Numbered Exercises**

(1) (a)  $\lambda^2 - 10\lambda + 9$ ,  $\lambda^2 - 10\lambda + 9$

(b) 1

(c) 2

(d) 2, -1, 1, 2

(e)  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ ,  $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$

(f) 1, 9

(g) 1, 2,  $\begin{bmatrix} 1 & 0 \\ 0 & 9 \end{bmatrix}$

(3)  $\begin{bmatrix} -5 & 18 \\ -3 & 10 \end{bmatrix}$



## CHAPTER 15

# SOME APPLICATIONS OF THE SPECTRAL THEOREM

### 15.1. Background

**Topics:** systems of linear differential equations, initial conditions, steady state solutions, the functional calculus for operators on finite dimensional vector spaces, Markov processes.

**15.1.1. Theorem.** *Let*

$$\frac{d\mathbf{u}}{dt} = A\mathbf{u} \tag{15.1.1}$$

*be a vector differential equation (that is, a system of ordinary linear differential equations) where  $A$  is an  $n \times n$  matrix and suppose that  $\mathbf{u}_0 = \mathbf{u}(0)$  is an initial value of the system. If  $A$  is a diagonalizable matrix (so that  $A = S\Lambda S^{-1}$  for some diagonal matrix  $\Lambda$  and some invertible matrix  $S$ ), then the equation (15.1.1) has the solution*

$$\mathbf{u}(t) = e^{At}\mathbf{u}_0 = Se^{\Lambda t}S^{-1}\mathbf{u}_0.$$

**15.1.2. Definition.** A MARKOV MATRIX is a square matrix with nonnegative entries and with each column adding to 1.

**15.1.3. Proposition** (Facts about Markov matrices.). *Let  $A$  be a Markov matrix. Then*

- (i)  $\lambda_1 = 1$  is an eigenvalue.
- (ii) The eigenvector  $\mathbf{e}_1$  corresponding to  $\lambda_1$  is nonnegative and it is a steady state.
- (iii) The other eigenvalues satisfy  $|\lambda_k| \leq 1$ .
- (iv) If any power of  $A$  has all entries strictly positive, then  $|\lambda_k| < 1$  for all  $k \neq 1$  and  $A^k\mathbf{u}_0 \rightarrow \mathbf{u}_\infty$  where the steady state  $\mathbf{u}_\infty$  is a multiple of  $\mathbf{e}_1$ .

## 15.2. Exercises

(1) Let  $A = \begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix}$ .

(a) The eigenvalues of  $A$  are  $\lambda_1 = \underline{\hspace{1cm}}$  and  $\lambda_2 = \underline{\hspace{1cm}}$ .

(b) The corresponding eigenvectors are  $\mathbf{e}_1 = (1, a)$  and  $\mathbf{e}_2 = (a, -a)$  where  $a = \underline{\hspace{1cm}}$ .

(c) Then  $e^{At} = a \begin{bmatrix} 1 + e^{-bt} & 1 - e^{-bt} \\ 1 - e^{-bt} & 1 + e^{-bt} \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .

(d) Let  $\mathbf{u}(t) = (x(t), y(t))$ . The general solution to the system of equations  $\frac{d\mathbf{u}}{dt} = A\mathbf{u}$  with initial conditions  $x_0 = 3$  and  $y_0 = 1$  is  $x(t) = a + be^{-ct}$  and  $y(t) = a - be^{-ct}$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ , and  $c = \underline{\hspace{1cm}}$ .

(e) Find the steady state solution to the system  $\frac{d\mathbf{u}}{dt} = A\mathbf{u}$  under the initial conditions given in (d). That is, find  $\mathbf{u}_\infty = \begin{bmatrix} x_\infty \\ y_\infty \end{bmatrix}$  where  $x_\infty = \lim_{t \rightarrow \infty} x(t)$  and  $y_\infty = \lim_{t \rightarrow \infty} y(t)$ .  
 Answer:  $\mathbf{u}_\infty = \begin{bmatrix} \phantom{0} \\ \phantom{0} \end{bmatrix}$ .

(2) Suppose that at time  $t$  the population  $y(t)$  of a predator and the population  $x(t)$  of its prey are governed by the equations

$$\begin{aligned} \frac{dx}{dt} &= 4x - 2y \\ \frac{dy}{dt} &= x + y. \end{aligned}$$

If at time  $t = 0$  the populations are  $x = 300$  and  $y = 200$ , then the populations at all future times  $t$  are  $x(t) = ae^{bt} + 200e^{ct}$  and  $y(t) = de^{bt} + ae^{ct}$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ ,  $c = \underline{\hspace{1cm}}$ , and  $d = \underline{\hspace{1cm}}$ . The long run ratio of populations of prey to predator approaches  $\underline{\hspace{1cm}}$  to  $\underline{\hspace{1cm}}$ .

(3) Use matrix methods to solve the initial value problem consisting of the system of differential equations

$$\begin{aligned} \frac{du}{dt} &= 4u - v - w \\ \frac{dv}{dt} &= u + 2v - w \\ \frac{dw}{dt} &= u - v + 2w \end{aligned}$$

and the initial conditions

$$u(0) = 2 \quad v(0) = -2 \quad w(0) = 7.$$

Answer:  $u(t) = ae^{bt} - e^{at}$ ;  $v(t) = ae^{bt} - ce^{at}$ ; and  $w(t) = ae^{bt} + de^{at}$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ ,  $c = \underline{\hspace{1cm}}$ , and  $d = \underline{\hspace{1cm}}$ .

(4) Consider the initial value problem:  $y'' - y' - 2y = 0$  with the initial conditions  $y_0 = 3$  and  $y'_0 = 3$ .

(a) Express the differential equation in the form  $\frac{d\mathbf{u}}{dt} = A\mathbf{u}$  where  $\mathbf{u} = (y, z)$  and  $z = y'$ .

Then  $A$  is the matrix  $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ , and  $c = \underline{\hspace{1cm}}$ .

- (b) The smaller eigenvalue of  $A$  is \_\_\_\_\_ and the larger is \_\_\_\_\_. The corresponding eigenvectors are  $(1, a)$  and  $(1, b)$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .
- (c) The diagonal form of  $A$  is  $\Lambda = \begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .
- (d) Find the diagonalizing matrix  $S$  for  $A$ . That is, find  $S$  so that  $\Lambda = S^{-1}AS$ . Answer:  $S = \begin{bmatrix} 1 & 1 \\ a & b \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .
- (e) The matrix  $e^{At}$  is  $\frac{1}{a} \begin{bmatrix} be^{ct} + e^{dt} & -e^{ct} + e^{dt} \\ -be^{ct} + be^{dt} & e^{ct} + be^{dt} \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ ,  $c = \underline{\hspace{1cm}}$ , and  $d = \underline{\hspace{1cm}}$ .
- (f) The solution to the initial value problem is  $y(t) = \underline{\hspace{2cm}}$ .
- (5) Use *the spectral theorem* to solve the initial value problem

$$y''' - 3y'' + 2y' = 0$$

where  $y(0) = 2$ ,  $y'(0) = 0$ , and  $y''(0) = 3$ .

Answer:  $y(t) = a + be^t + ce^{dt}$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ ,  $c = \underline{\hspace{1cm}}$ , and  $d = \underline{\hspace{1cm}}$ .

- (6) Let  $G_0 = 0$  and  $G_1 = \frac{1}{2}$ . For each  $k \geq 0$  let  $G_{k+2}$  be the average of  $G_k$  and  $G_{k+1}$ .
- (a) Find the transition matrix  $A$  which takes the vector  $(G_{k+1}, G_k)$  to the vector  $(G_{k+2}, G_{k+1})$ .

Answer:  $A = \begin{bmatrix} \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} \end{bmatrix}$ .

- (b) Find a diagonal matrix  $\Lambda$  which is similar to  $A$ .

Answer:  $\Lambda = \begin{bmatrix} \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} \end{bmatrix}$ .

- (c) Find a matrix  $S$  such that  $A = SAS^{-1}$ .

Answer:  $S = \begin{bmatrix} \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} \end{bmatrix}$ .

- (d) Determine the long run behavior of the numbers  $G_k$ .

Answer:  $G_\infty := \lim_{k \rightarrow \infty} G_k = \underline{\hspace{1cm}}$ .

- (7) Let  $T$  be the operator on  $\mathbb{R}^2$  whose matrix representation is  $\begin{bmatrix} -7 & 8 \\ -16 & 17 \end{bmatrix}$ . Use the *spectral theorem* to find  $\sqrt{T}$ . (A *square root* of  $T$  is an operator whose square is  $T$ .)

Answer:  $\sqrt{T} = \begin{bmatrix} -1 & a \\ b & c \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ , and  $c = \underline{\hspace{1cm}}$ .

- (8) Let  $A = \begin{bmatrix} 4 & 3 \\ 1 & 2 \end{bmatrix}$ . Find  $A^{100}$ . (Write an *exact* answer—not a decimal approximation.)

Answer:  $A^{100} = \frac{1}{4} \begin{bmatrix} \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} \end{bmatrix}$ .

- (9) Let  $T$  be the operator on  $\mathbb{R}^3$  whose matrix representation is  $\begin{bmatrix} 2 & -2 & 1 \\ -1 & 1 & 1 \\ -1 & 2 & 0 \end{bmatrix}$ .

(a) Write  $T$  as a linear combination of projections.

Answer:  $T = c_1 E_1 + c_2 E_2 + c_3 E_3$  where  $c_1 = \underline{\hspace{1cm}}$ ,  $c_2 = \underline{\hspace{1cm}}$ ,  $c_3 = \underline{\hspace{1cm}}$ ,  
 $E_1 = \begin{bmatrix} a & b & -b \\ a & b & -b \\ a & -b & b \end{bmatrix}$ ,  $E_2 = \begin{bmatrix} b & a & b \\ b & a & b \\ b & a & b \end{bmatrix}$ , and  $E_3 = \begin{bmatrix} b & -b & a \\ -b & b & a \\ -b & b & a \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$  and  
 $b = \underline{\hspace{1cm}}$ .

(b) Calculate the following:  $E_1 E_2 = \begin{bmatrix} & & \\ & & \\ & & \end{bmatrix}$ ;  $E_1 E_3 = \begin{bmatrix} & & \\ & & \\ & & \end{bmatrix}$ ;

$$E_2 E_3 = \begin{bmatrix} & & \\ & & \\ & & \end{bmatrix}.$$

(c)  $E_1 + E_2 + E_3 = \begin{bmatrix} & & \\ & & \\ & & \end{bmatrix}$ .

(d) Write  $T^3$  as a linear combination of  $E_1$ ,  $E_2$ , and  $E_3$ .

Answer:  $T^3 = \underline{\hspace{1cm}} E_1 + \underline{\hspace{1cm}} E_2 + \underline{\hspace{1cm}} E_3$ .

- (10) Let  $A$  be the matrix whose eigenvalues are  $\lambda_1 = -1$ ,  $\lambda_2 = 1/2$ , and  $\lambda_3 = 1/3$ , and whose corresponding eigenvectors are  $\mathbf{e}_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$ ,  $\mathbf{e}_2 = \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}$ , and  $\mathbf{e}_3 = \begin{bmatrix} 0 \\ -1 \\ 0 \end{bmatrix}$ .

(a) Solve the difference equation  $\mathbf{x}_{k+1} = A\mathbf{x}_k$  (where  $\mathbf{x}_k = \begin{bmatrix} u_k \\ v_k \\ w_k \end{bmatrix}$ ) subject to the initial condition  $\mathbf{x}_0 = \begin{bmatrix} 10 \\ 20 \\ 30 \end{bmatrix}$ .

Answer:  $u_k = a(-b)^k - cd^k$ ,  $v_k = cd^k$ , and  $w_k = a(-b)^k$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ ,  
 $c = \underline{\hspace{1cm}}$ , and  $d = \underline{\hspace{1cm}}$ .

(b) Each  $\mathbf{x}_k$  can be written as a linear combination of the vectors (  $\underline{\hspace{1cm}}$ ,  $\underline{\hspace{1cm}}$ ,  $\underline{\hspace{1cm}}$  )  
 and (  $\underline{\hspace{1cm}}$ ,  $\underline{\hspace{1cm}}$ ,  $\underline{\hspace{1cm}}$  ).

(c) The value of  $\mathbf{x}_{1000}$  is approximately (  $\underline{\hspace{1cm}}$ ,  $\underline{\hspace{1cm}}$ ,  $\underline{\hspace{1cm}}$  ).

- (11) Let  $A$  be as in the preceding exercise. Solve the differential equation  $\frac{d\mathbf{x}}{dt} = A\mathbf{x}$  subject to

the initial conditions  $\mathbf{x}_0 = \begin{bmatrix} 10 \\ 20 \\ 30 \end{bmatrix}$ . Answer:  $\mathbf{x}(t) = (ae^{-t} - be^{ct}, be^{ct}, ae^{-t})$  where  $a = \underline{\hspace{1cm}}$ ,  
 $b = \underline{\hspace{1cm}}$ , and  $c = \underline{\hspace{1cm}}$ .

- (12) Suppose three cities A, B, and C are centers for trucks. Every month half of the trucks in A and half of those in B go to C. The other half of the trucks in A and the other half of the trucks in B stay where they are. Every month half of the trucks in C go to A and the other half go to B.

- (a) What is the (Markov) transition matrix which acts on the vector  $\begin{bmatrix} a_0 \\ b_0 \\ c_0 \end{bmatrix}$  (where  $a_0$  is the number of trucks initially in A, etc.)?

Answer:  $\begin{bmatrix} & & \\ & & \\ & & \end{bmatrix}$ .

- (b) If there are always 450 trucks, what is the long run distribution of trucks? Answer:  $a_\infty = \underline{\hspace{1cm}}$ ,  $b_\infty = \underline{\hspace{1cm}}$ ,  $c_\infty = \underline{\hspace{1cm}}$ .

## 15.3. Problems

- (1) Initially a 2100 gallon tank  $M$  is full of water and an 1800 gallon tank  $N$  is full of water in which 100 pounds of salt has been dissolved. Fresh water is pumped into tank  $M$  at a rate of 420 gallons per minute and salt water is released from tank  $N$  at the same rate. Additionally, the contents of  $M$  are pumped into  $N$  at a rate of 490 gallons per minute and the contents of  $N$  are pumped into  $M$  at a rate sufficient to keep both tanks full.

How long does it take (to the nearest second) for the concentration of salt in tank  $M$  to reach a maximum? And how much salt is there (to three significant figures) in  $M$  at that time?

- (2) Show that if  $A$  is a diagonalizable  $n \times n$  matrix, then

$$\det(\exp A) = \exp(\operatorname{tr} A).$$

*Hint.* What would be a reasonable definition of  $\exp A$  if  $A$  were a diagonal matrix?

- (3) Explain carefully how to use matrix methods to solve the initial value problem

$$y'' - y' - 6y = 0$$

under the initial conditions  $y_0 = -2$  and  $y'_0 = 14$ . Carry out the computations you describe.

- (4) Explain carefully how to use matrix methods to solve the initial value problem consisting of the system of differential equations

$$\begin{cases} \frac{dv}{dt} = -v + w \\ \frac{dw}{dt} = v - w \end{cases} \quad \text{and the initial conditions} \quad \begin{cases} v(0) = 5 \\ w(0) = -3. \end{cases}$$

Carry out the computation you describe.

- (5) Show how to use the *spectral theorem* to solve the initial value problem consisting of the system of differential equations

$$\begin{aligned} \frac{du}{dt} &= -7u - 5v + 5w \\ \frac{dv}{dt} &= 2u + 3v - 2w \\ \frac{dw}{dt} &= -8u - 5v + 6w \end{aligned}$$

and the initial conditions

$$u(0) = 2 \quad v(0) = 1 \quad w(0) = -1.$$

- (6) Explain carefully how to use the *spectral theorem* to find a square root of the matrix  $A = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}$ . Illustrate your discussion by carrying out the computation.

(7) Let  $A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$ .

- (a) Does  $A$  have a cube root? Explain.  
 (b) Does  $A$  have a square root? Explain.

- (8) Let  $A$  be a symmetric  $2 \times 2$  matrix whose trace is 20 and whose determinant is 64. Suppose that the eigenspace associated with the smaller eigenvalue of  $A$  is the line  $x - y = 0$ . Find a matrix  $B$  such that  $B^2 = A$ .

**15.4. Answers to Odd-Numbered Exercises**

- (1) (a)  $0, -2$   
(b)  $1$   
(c)  $\frac{1}{2}, 2$   
(d)  $2, 1, 2$   
(e)  $\begin{bmatrix} 2 \\ 2 \end{bmatrix}$
- (3)  $3, 2, 5, 4$
- (5)  $\frac{7}{2}, -3, \frac{3}{2}, 2$
- (7)  $2, -4, 5$
- (9) (a)  $-1, 1, 3, 0, \frac{1}{2}$   
(b)  $\begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$   
(c)  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$   
(d)  $-1, 1, 27$
- (11)  $30, 20, \frac{1}{2}$





## CHAPTER 16

# EVERY OPERATOR IS DIAGONALIZABLE PLUS NILPOTENT

### 16.1. Background

**Topics:** generalized eigenspaces, nilpotent operators

**16.1.1. Definition.** An operator  $T$  on a vector space is **NILPOTENT** if  $T^n = \mathbf{0}$  for some  $n \in \mathbb{N}$ . Similarly, a square matrix is **NILPOTENT** if some power of it is the zero matrix.

**16.1.2. Theorem.** Let  $T$  be an operator on a finite dimensional vector space  $V$ . Suppose that the minimal polynomial for  $T$  factors completely into linear factors

$$m_T(x) = (x - \lambda_1)^{r_1} \cdots (x - \lambda_k)^{r_k}$$

where  $\lambda_1, \dots, \lambda_k$  are the (distinct) eigenvalues of  $T$ . For each  $j$  let  $W_j = \ker(T - \lambda_j I)^{r_j}$  and  $E_j$  be the projection of  $V$  onto  $W_j$  along  $W_1 + \cdots + W_{j-1} + W_{j+1} + \cdots + W_k$ . Then

$$V = W_1 \oplus W_2 \oplus \cdots \oplus W_k,$$

each  $W_j$  is invariant under  $T$ , and  $I = E_1 + \cdots + E_k$ . Furthermore, the operator

$$D = \lambda_1 E_1 + \cdots + \lambda_k E_k$$

is diagonalizable, the operator

$$N = T - D$$

is nilpotent, and  $N$  commutes with  $D$ .

**16.1.3. Corollary.** Every operator on a finite dimensional complex vector space is the sum of a diagonal operator and a nilpotent one.

**16.1.4. Definition.** Since, in the preceding theorem,  $T = D + N$  where  $D$  is diagonalizable and  $N$  is nilpotent, we say that  $D$  is the **DIAGONALIZABLE PART** of  $T$  and  $N$  is the **NILPOTENT PART** of  $T$ . The subspace  $W_j = \ker(T - \lambda_j I)^{r_j}$  is **GENERALIZED EIGENSPACE** associated with the eigenvalue  $\lambda_j$ .

**16.2. Exercises**

- (1) Let  $T$  be the operator on  $\mathbb{R}^2$  whose matrix representation is  $\begin{bmatrix} 2 & 1 \\ -1 & 4 \end{bmatrix}$ .

(a) Explain briefly why  $T$  is not diagonalizable.

Answer: \_\_\_\_\_.

- (b) Find the diagonalizable and nilpotent parts of  $T$ . Answer:  $D = \begin{bmatrix} a & b \\ b & a \end{bmatrix}$  and

$$N = \begin{bmatrix} -c & c \\ -c & c \end{bmatrix} \text{ where } a = \_, b = \_, \text{ and } c = \_.$$

- (2) Let  $T$  be the operator on  $\mathbb{R}^2$  whose matrix representation is  $\begin{bmatrix} 4 & -2 \\ 2 & 0 \end{bmatrix}$ .

(a) Explain briefly why  $T$  is not diagonalizable.

Answer: \_\_\_\_\_.

- (b) Find the diagonalizable and nilpotent parts of  $T$ . Answer:  $D = \begin{bmatrix} & \\ & \end{bmatrix}$  and

$$N = \begin{bmatrix} & \\ & \end{bmatrix}$$

- (3) Let  $T$  be the operator on  $\mathbb{R}^3$  whose matrix representation is  $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$ .

(a) The characteristic polynomial of  $T$  is  $(\lambda)^p(\lambda - 1)^q$  where  $p = \_$  and  $q = \_$ .

(b) The minimal polynomial of  $T$  is  $(\lambda)^r(\lambda - 1)^s$  where  $r = \_$  and  $s = \_$ .

(c) Explain briefly how we know from (b) that  $T$  is not diagonalizable.

Answer: \_\_\_\_\_.

(d) The eigenspace  $M_1$  (corresponding to the smaller eigenvalue of  $T$ ) is  $\text{span}\{(\_, \_, 1)\}$  and the eigenspace  $M_2$  (corresponding to the larger eigenvalue) is  $\text{span}\{(1, \_, \_)\}$ .

(e) Explain briefly how we know from (d) that  $T$  is not diagonalizable.

Answer: \_\_\_\_\_.

(f) The generalized eigenspace  $W_1$  (corresponding to the smaller eigenvalue) is  $W_1 = \text{span}\{(\_, \_, 1)\}$  and the generalized eigenspace  $W_2$  (corresponding to the larger eigenvalue) is  $\text{span}\{(1, a, a), (a, 1, a)\}$ , where  $a = \_$ .

- (g) The (matrix representing the) projection  $E_1$  of  $\mathbb{R}^3$  along  $W_2$  onto  $W_1$  is  $\begin{bmatrix} a & a & a \\ a & a & a \\ a & a & b \end{bmatrix}$

where  $a = \_$  and  $b = \_$ .

- (h) The (matrix representing the) projection  $E_2$  of  $\mathbb{R}^3$  along  $W_1$  onto  $W_2$  is  $\begin{bmatrix} a & b & b \\ b & a & b \\ b & b & b \end{bmatrix}$

where  $a = \_$  and  $b = \_$ .

- (i) The diagonalizable part of  $T$  is  $D = \begin{bmatrix} a & b & b \\ b & a & b \\ b & b & b \end{bmatrix}$  and the nilpotent part of  $T$  is  $N = \begin{bmatrix} b & a & b \\ b & b & b \\ b & b & b \end{bmatrix}$  where  $a = \_$  and  $b = \_$ .

(j) A matrix  $S$  which diagonalizes  $D$  is  $\begin{bmatrix} a & b & a \\ a & a & b \\ b & a & a \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .

(k) The diagonal form  $\Lambda$  of the diagonalizable part  $D$  of  $T$  is  $\begin{bmatrix} a & a & a \\ a & b & a \\ a & a & b \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .

(l) Show that  $D$  commutes with  $N$  by computing  $DN$  and  $ND$ .

Answer:  $DN = ND = \begin{bmatrix} a & b & a \\ a & a & a \\ a & a & a \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .

(4) Let  $T$  be the operator on  $\mathbb{R}^3$  whose matrix representation is  $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{bmatrix}$ .

(a) The characteristic polynomial of  $T$  is  $(\lambda)^p(\lambda - 1)^q$  where  $p = \underline{\hspace{1cm}}$  and  $q = \underline{\hspace{1cm}}$ .

(b) The minimal polynomial of  $T$  is  $(\lambda)^r(\lambda - 1)^s$  where  $r = \underline{\hspace{1cm}}$  and  $s = \underline{\hspace{1cm}}$ .

(c) Explain briefly how we know from (b) that  $T$  is not diagonalizable.

Answer:  $\underline{\hspace{5cm}}$ .

(d) The eigenspace  $M_1$  (corresponding to the smaller eigenvalue of  $T$ ) is  $\text{span}\{(1, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})\}$  and the eigenspace  $M_2$  (corresponding to the larger eigenvalue) is  $\text{span}\{(1, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})\}$ .

(e) Explain briefly how we know from (d) that  $T$  is not diagonalizable.

Answer:  $\underline{\hspace{5cm}}$ .

(f) The generalized eigenspace  $W_1$  (corresponding to the smaller eigenvalue) is  $W_1 = \text{span}\{(1, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})\}$  and the generalized eigenspace  $W_2$  (corresponding to the larger eigenvalue) is  $\text{span}\{(1, a, a), (a, 1, a)\}$ , where  $a = \underline{\hspace{1cm}}$ .

(g) The (matrix representing the) projection  $E_1$  of  $\mathbb{R}^3$  along  $W_2$  onto  $W_1$  is  $\begin{bmatrix} a & a & b \\ a & a & -b \\ a & a & b \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .

(h) The (matrix representing the) projection  $E_2$  of  $\mathbb{R}^3$  along  $W_1$  onto  $W_2$  is  $\begin{bmatrix} a & b & -a \\ b & a & a \\ b & b & b \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .

(i) The diagonalizable part of  $T$  is  $D = \begin{bmatrix} a & b & -a \\ b & a & a \\ b & b & b \end{bmatrix}$  and the nilpotent part of  $T$  is  $N = \begin{bmatrix} b & a & a \\ b & b & b \\ b & b & b \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .

(j) A matrix  $S$  which diagonalizes  $D$  is  $\begin{bmatrix} a & a & b \\ -a & b & a \\ a & b & b \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .

(k) The diagonal form  $\Lambda$  of the diagonalizable part  $D$  of  $T$  is  $\begin{bmatrix} a & a & a \\ a & b & a \\ a & a & b \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .

(l) When comparing this exercise with the preceding one it may seem that the correct answer to part (i) should be that the diagonalizable part of  $T$  is  $D = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$  and

the nilpotent part of  $[T]$  is  $N = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$ , because  $D$  is diagonal,  $N$  is nilpotent, and  $[T] = D + N$ . Explain briefly why this is not correct.

Answer: \_\_\_\_\_.

(5) Let  $T$  be the operator on  $\mathbb{R}^3$  whose matrix representation is  $\begin{bmatrix} 3 & 1 & -1 \\ 2 & 2 & -1 \\ 2 & 2 & 0 \end{bmatrix}$ .

- (a) The characteristic polynomial of  $T$  is  $(\lambda - 1)^p(\lambda - 2)^q$  where  $p = \underline{\hspace{1cm}}$  and  $q = \underline{\hspace{1cm}}$ .
- (b) The minimal polynomial of  $T$  is  $(\lambda - 1)^r(\lambda - 2)^s$  where  $r = \underline{\hspace{1cm}}$  and  $s = \underline{\hspace{1cm}}$ .
- (c) The eigenspace  $M_1$  (corresponding to the smaller eigenvalue of  $T$ ) is  $\text{span}\{(1, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})\}$  and the eigenspace  $M_2$  (corresponding to the larger eigenvalue) is  $\text{span}\{(1, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})\}$ .
- (d) The generalized eigenspace  $W_1$  (corresponding to the smaller eigenvalue) is  $\text{span}\{(1, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})\}$  and the generalized eigenspace  $W_2$  (corresponding to the larger eigenvalue) is  $\text{span}\{(1, a, b), (0, b, a)\}$ , where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .

(e) The diagonalizable part of  $T$  is  $D = \begin{bmatrix} a & a & b \\ b & c & b \\ -c & c & c \end{bmatrix}$  and the nilpotent part of  $T$  is

$$N = \begin{bmatrix} c & b & -a \\ c & b & -a \\ 2c & b & -c \end{bmatrix} \text{ where } a = \underline{\hspace{1cm}}, b = \underline{\hspace{1cm}}, \text{ and } c = \underline{\hspace{1cm}}.$$

(f) A matrix  $S$  which diagonalizes  $D$  is  $\begin{bmatrix} a & a & b \\ b & a & b \\ c & b & a \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ , and  $c = \underline{\hspace{1cm}}$ .

(g) The diagonal form  $\Lambda$  of the diagonalizable part  $D$  of  $T$  is  $\begin{bmatrix} \lambda & 0 & 0 \\ 0 & \mu & 0 \\ 0 & 0 & \mu \end{bmatrix}$  where  $\lambda = \underline{\hspace{1cm}}$  and  $\mu = \underline{\hspace{1cm}}$ .

(h) Show that  $D$  commutes with  $N$  by computing  $DN$  and  $ND$ .

$$\text{Answer: } DN = ND = \begin{bmatrix} a & b & -c \\ a & b & -c \\ 2a & b & -a \end{bmatrix} \text{ where } a = \underline{\hspace{1cm}}, b = \underline{\hspace{1cm}}, \text{ and } c = \underline{\hspace{1cm}}.$$

(6) Let  $T$  be the operator on  $\mathbb{R}^4$  whose matrix representation is  $\begin{bmatrix} 0 & 1 & 0 & -1 \\ -2 & 3 & 0 & -1 \\ -2 & 1 & 2 & -1 \\ 2 & -1 & 0 & 3 \end{bmatrix}$ .

- (a) The characteristic polynomial of  $T$  is  $(\lambda - 2)^p$  where  $p = \underline{\hspace{1cm}}$ .
- (b) The minimal polynomial of  $T$  is  $(\lambda - 2)^r$  where  $r = \underline{\hspace{1cm}}$ .

(c) The diagonalizable part of  $T$  is  $D = \begin{bmatrix} a & b & b & b \\ b & a & b & b \\ b & b & a & b \\ b & b & b & a \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .

(c) The nilpotent part of  $T$  is  $N = \begin{bmatrix} -a & b & c & -b \\ -a & b & c & -b \\ -a & b & c & -b \\ a & -b & c & b \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ , and  $c = \underline{\hspace{1cm}}$ .

(7) Let  $T$  be the operator on  $\mathbb{R}^5$  whose matrix representation is  $\begin{bmatrix} 1 & 0 & 0 & 1 & -1 \\ 0 & 1 & -2 & 3 & -3 \\ 0 & 0 & -1 & 2 & -2 \\ 1 & -1 & 1 & 0 & 1 \\ 1 & -1 & 1 & -1 & 2 \end{bmatrix}$ .

(a) Find the characteristic polynomial of  $T$ .

Answer:  $c_T(\lambda) = (\lambda + 1)^p(\lambda - 1)^q$  where  $p = \underline{\hspace{1cm}}$  and  $q = \underline{\hspace{1cm}}$ .

(b) Find the minimal polynomial of  $T$ .

Answer:  $m_T(\lambda) = (\lambda + 1)^r(\lambda - 1)^s$  where  $r = \underline{\hspace{1cm}}$  and  $s = \underline{\hspace{1cm}}$ .

(c) Find the eigenspaces  $M_1$  and  $M_2$  of  $T$ .

Answer:  $M_1 = \text{span}\{(a, 1, b, a, a)\}$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ ; and

$M_2 = \text{span}\{(1, a, b, b, b), (b, b, b, 1, a)\}$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .

(d) Find the diagonalizable part of  $T$ .

Answer:  $D = \begin{bmatrix} a & b & b & b & b \\ b & a & -c & c & -c \\ b & b & -a & c & -c \\ b & b & b & a & b \\ b & b & b & b & a \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ , and  $c = \underline{\hspace{1cm}}$ .

(e) Find the nilpotent part of  $T$ .

Answer:  $N = \begin{bmatrix} a & a & a & b & -b \\ a & a & a & b & -b \\ a & a & a & a & a \\ b & -b & b & -b & b \\ b & -b & b & -b & b \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .

(f) Find a matrix  $S$  which diagonalizes the diagonalizable part  $D$  of  $T$ . What is the diagonal form  $\Lambda$  of  $D$  associated with this matrix?

Answer:  $S = \begin{bmatrix} a & b & a & a & a \\ b & a & b & a & a \\ b & a & a & b & a \\ a & a & a & b & b \\ a & a & a & a & b \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .

and  $\Lambda = \begin{bmatrix} -a & 0 & 0 & 0 & 0 \\ 0 & a & 0 & 0 & 0 \\ 0 & 0 & a & 0 & 0 \\ 0 & 0 & 0 & a & 0 \\ 0 & 0 & 0 & 0 & a \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$ .

## 16.3. Problems

- (1) Explain in detail how to find the diagonalizable and nilpotent parts of the matrix  $A = \begin{bmatrix} -3 & -4 & 5 \\ 6 & 8 & -6 \\ -2 & -1 & 4 \end{bmatrix}$ . Carry out the computations you describe.
- (2) Consider the matrix  $A = \begin{bmatrix} 2 & 0 & -2 \\ 0 & 0 & 2 \\ 0 & 2 & 0 \end{bmatrix}$ . In each part below explain carefully what you are doing.
- Find the characteristic polynomial for  $A$ .
  - Find the minimal polynomial for  $A$ . What can you conclude from the form of the minimal polynomial?
  - Find the eigenspace associated with each eigenvalue of  $A$ . Do the eigenvectors of  $A$  span  $\mathbb{R}^3$ ? What can you conclude from this?
  - Find the generalized eigenspaces  $W_1$  and  $W_2$  associated with the eigenvalues of  $A$ .
  - Find the projection operators  $E_1$  of  $\mathbb{R}^3$  onto  $W_1$  along  $W_2$  and  $E_2$  of  $\mathbb{R}^3$  onto  $W_2$  along  $W_1$ .
  - Find the diagonalizable part  $D$  of  $A$ . Express  $D$  both as a single matrix and as a linear combination of projections.
  - Find a matrix  $S$  which diagonalizes  $D$ . What is the resulting diagonal form  $\Lambda$  of  $D$ ?
  - Find the nilpotent part  $N$  of  $A$ . What is the smallest power of  $N$  which vanishes?
- (3) Let  $T$  be the operator on  $\mathbb{R}^3$  whose matrix representation is  $\begin{bmatrix} 1 & -1 & 0 \\ 1 & 3 & -1 \\ 0 & 0 & 1 \end{bmatrix}$ .
- Explain how to find the characteristic polynomial for  $T$ . Then carry out the computation.
  - What is the *minimal polynomial* for a matrix? Find the minimal polynomial for  $T$  and explain how you know your answer is correct. What can you conclude from the form of this polynomial?
  - Find the eigenspaces associated with each eigenvalue of  $T$ . Do the eigenvectors of  $T$  span  $\mathbb{R}^3$ ? What can you conclude from this?
  - Find the generalized eigenspaces  $W_1$  and  $W_2$  associated with the eigenvalues of  $T$ .
  - Find the projection operators  $E_1$  of  $\mathbb{R}^3$  onto  $W_1$  along  $W_2$  and  $E_2$  of  $\mathbb{R}^3$  onto  $W_2$  along  $W_1$ .
  - Find the diagonalizable part  $D$  of  $T$ . Express  $D$  both as a single matrix and as a linear combination of projections.
  - Find a matrix  $S$  which diagonalizes  $D$ . What is the resulting diagonal form  $\Lambda$  of  $D$ ?
  - Find the nilpotent part  $N$  of  $T$ . What is the smallest power of  $N$  which vanishes?

**16.4. Answers to Odd-Numbered Exercises**

- (1) (a) The single one-dimensional eigenspace does not span  $\mathbb{R}^2$ . (OR: the minimal polynomial  $(\lambda - 3)^2$  has a second degree factor—see theorem 13.1.11.)  
(b) 3, 0, 1
- (3) (a) 1, 2  
(b) 1, 2  
(c) The minimal polynomial has a second degree factor (see theorem 13.1.11).  
(d) 0, 0, 0, 0  
(e) The eigenspaces do not span  $\mathbb{R}^3$ .  
(f) 0, 0, 0  
(g) 0, 1  
(h) 1, 0  
(i) 1, 0  
(j) 0, 1  
(k) 0, 1  
(l) 0, 1
- (5) (a) 1, 2  
(b) 1, 2  
(c) 0, 2, 1, 2  
(d) 0, 2, 1, 0  
(e) 1, 0, 2  
(f) 1, 0, 2  
(g) 1, 2  
(h) 4, 0, 2
- (7) (a) 1, 4  
(b) 1, 2  
(c) 0, 1, 1, 0  
(d) 1, 0, 2  
(e) 0, 1  
(f) 0, 1, 1





**Part 5**

**THE GEOMETRY OF INNER PRODUCT  
SPACES**



## CHAPTER 17

# COMPLEX ARITHMETIC

### 17.1. Background

**Topics:** complex arithmetic, absolute value and argument of a complex number, *De Moivre's theorem*.

## 17.2. Exercises

- (1)  $\operatorname{Re}\left(\frac{2+3i}{3-4i}\right) = \underline{\hspace{2cm}}$ .
- (2)  $\operatorname{Im}\left(\frac{2-3i}{2+3i}\right) = \underline{\hspace{2cm}}$ .
- (3)  $\left|\frac{21+7i}{1-2i}\right| = a\sqrt{b}$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .
- (4)  $\operatorname{Arg}(-2\sqrt{3}+2i) = \underline{\hspace{2cm}}$ .
- (5) Let  $z = \frac{2-\sqrt{3}-(1+2\sqrt{3})i}{1+2i}$ . Then  $z^{10} = a(1-bi)$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ ,  
 $|z^{10}| = \underline{\hspace{1cm}}$ , and  $\operatorname{Arg} z^{10} = \underline{\hspace{1cm}}$ .
- (6) If  $z = \frac{1}{\sqrt{2}}(1-i)$ , then  $z^{365} = a+bi$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .
- (7) The cube roots of  $-27$  are  $a+bi$ ,  $a-bi$ , and  $c+di$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ ,  
 $c = \underline{\hspace{1cm}}$ , and  $d = \underline{\hspace{1cm}}$ .
- (8) The complex number  $w = 1+2i$  has 13 thirteenth roots. The sum of these roots is  $a+bi$   
 where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .
- (9) Let  $z_1 = (1, i, 1+i)$ ,  $z_2 = (i, 0, 1-i)$ , and  $z_3 = (-1, 1+2i, 7+3i)$ . Show that the set  
 $\{z_1, z_2, z_3\}$  is linearly dependent in  $\mathbb{C}^3$  by finding scalars  $\alpha$  and  $\beta$  such that  $\alpha z_1 + \beta z_2 - z_3 = 0$ .  
 Answer:  $\alpha = \underline{\hspace{2cm}}$  and  $\beta = \underline{\hspace{2cm}}$ .
- (10) Let  $A = \begin{bmatrix} 1 & 1 & 1 & i \\ 1 & i & 1 & i \\ 0 & 1+i & 0 & 0 \\ 2 & 0 & 2 & 2i \end{bmatrix}$ . Then the rank of  $A$  is  $\underline{\hspace{1cm}}$  and the nullity of  $A$  is  $\underline{\hspace{1cm}}$ .
- (11) Let  $A = \begin{bmatrix} 0 & 1 & 1 \\ 0 & i & 1 \\ i & i & i \end{bmatrix}$ . Then  $A^{-1} = \frac{1}{2} \begin{bmatrix} -2a & c & -2b \\ a+b & -a-b & c \\ a-b & a+b & c \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ ,  
 and  $c = \underline{\hspace{1cm}}$ .
- (12) If  $A = \begin{bmatrix} i & 0 & -i \\ -2i & 1 & -1-4i \\ 2-i & i & 3 \end{bmatrix}$ , then  $A^{-1} = \begin{bmatrix} -a-ai & b+ai & -a+bi \\ a-di & -c+ai & -a-di \\ -a+bi & b+ai & -a+bi \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$ ,  
 $b = \underline{\hspace{1cm}}$ ,  $c = \underline{\hspace{1cm}}$ , and  $d = \underline{\hspace{1cm}}$ .
- (13) Consider the initial value problem:  $y'' + 4y = 0$  with the initial conditions  $y_0 = 3$  and  $y'_0 = 2$ .
- (a) Express the differential equation in the form  $\frac{d\mathbf{u}}{dt} = A\mathbf{u}$  where  $\mathbf{u} = (y, z)$  and  $z = y'$ .  
 Then  $A$  is the matrix  $\begin{bmatrix} a & b \\ -c & a \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ , and  $c = \underline{\hspace{1cm}}$ .
- (b) The eigenvalues of  $A$  are  $\underline{\hspace{1cm}}$  and  $\underline{\hspace{1cm}}$ ; and the corresponding eigenvectors are  
 $(1, a)$  and  $(1, -a)$  where  $a = \underline{\hspace{1cm}}$ .
- (c) The diagonal form of  $A$  is  $\Lambda = \begin{bmatrix} a & 0 \\ 0 & -a \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$ .

(d) Find the diagonalizing matrix  $S$  for  $A$ . That is, find  $S$  so that  $\Lambda = S^{-1}AS$ . Answer:

$$S = \begin{bmatrix} 1 & 1 \\ a & -a \end{bmatrix} \text{ where } a = \underline{\hspace{2cm}} .$$

(e) Find the matrix  $e^{At}$ . Express your answer in terms of real trigonometric functions—

$$\text{not complex exponentials. Answer: } e^{At} = \begin{bmatrix} f(t) & a g(t) \\ -b g(t) & f(t) \end{bmatrix} \text{ where } a = \underline{\hspace{2cm}} , \\ b = \underline{\hspace{2cm}} , f(t) = \underline{\hspace{2cm}} , \text{ and } g(t) = \underline{\hspace{2cm}} .$$

(f) The solution to the initial value problem is  $y(t) = \underline{\hspace{4cm}} .$

(14) Let  $A = \begin{bmatrix} 5 & -12 \\ 2 & -5 \end{bmatrix}$ .

(a) Find the eigenvalues of  $A$ . Answer:  $\lambda_1 = \underline{\hspace{2cm}} ; \lambda_2 = \underline{\hspace{2cm}} .$

(b) Find a factorization of  $A$  in the form  $S\Lambda S^{-1}$  where  $\Lambda$  is a diagonal matrix.

$$\text{Answer: } S = \begin{bmatrix} \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} \end{bmatrix} ; \Lambda = \begin{bmatrix} \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} \end{bmatrix} ; S^{-1} = \begin{bmatrix} \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} \end{bmatrix} .$$

$$(c) \text{ Find a square root of } A. \text{ Answer: } \sqrt{A} = \begin{bmatrix} \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} \end{bmatrix} .$$

**17.3. Problems**

- (1) Explain in detail how to find all the cube roots of  $i$ .
- (2) Show that three points  $z_1$ ,  $z_2$ , and  $z_3$  in the complex plane form an equilateral triangle if and only if

$$z_1^2 + z_2^2 + z_3^2 = z_1z_2 + z_1z_3 + z_2z_3.$$

**17.4. Answers to Odd-Numbered Exercises**

- (1)  $-\frac{6}{25}$
- (3) 7, 2
- (5) 512,  $\sqrt{3}$ , 1024,  $-\frac{\pi}{3}$
- (7)  $\frac{3}{2}$ ,  $\frac{3\sqrt{3}}{2}$ ,  $-3$ , 0
- (9)  $2 - i$ ,  $1 + 3i$
- (11) 1,  $i$ , 0
- (13)
  - (a) 0, 1, 4
  - (b)  $2i$ ,  $-2i$ ,  $2i$
  - (c)  $2i$
  - (d)  $2i$
  - (e)  $\frac{1}{2}$ , 2,  $\cos 2t$ ,  $\sin 2t$ ,
  - (f)  $3 \cos 2t + \sin 2t$





## REAL AND COMPLEX INNER PRODUCT SPACES

### 18.1. Background

**Topics:** inner products in real and complex vector spaces, the *Schwarz inequality*, the *parallelogram law*, the *Pythagorean theorem*, the norm induced by an inner product, the metric induced by a norm, orthogonal (or perpendicular) vectors, the angle between two vectors, row space and column space of a matrix.

**18.1.1. Definition.** Let  $V$  be a (real or complex) vector space. A function which associates to each pair of vectors  $\mathbf{x}$  and  $\mathbf{y}$  in  $V$  a (real or complex) number  $\langle \mathbf{x}, \mathbf{y} \rangle$  (often written  $\mathbf{x} \cdot \mathbf{y}$ ) is an **INNER PRODUCT** (or a **DOT PRODUCT**) on  $V$  provided that the following four conditions are satisfied:

(a) If  $\mathbf{x}, \mathbf{y}, \mathbf{z} \in V$ , then

$$\langle \mathbf{x} + \mathbf{y}, \mathbf{z} \rangle = \langle \mathbf{x}, \mathbf{z} \rangle + \langle \mathbf{y}, \mathbf{z} \rangle.$$

(b) If  $\mathbf{x}, \mathbf{y} \in V$  and  $\alpha \in \mathbb{C}$  (or  $\mathbb{R}$ ), then

$$\langle \alpha \mathbf{x}, \mathbf{y} \rangle = \alpha \langle \mathbf{x}, \mathbf{y} \rangle.$$

(c) If  $\mathbf{x}, \mathbf{y} \in V$ , then

$$\langle \mathbf{x}, \mathbf{y} \rangle = \overline{\langle \mathbf{y}, \mathbf{x} \rangle}.$$

(d) For every nonzero  $\mathbf{x}$  in  $V$  we have  $\langle \mathbf{x}, \mathbf{x} \rangle > 0$ .

Conditions (a) and (b) show that an inner product is linear in its first variable. It is easy to see that an inner product is *conjugate linear* in its second variable. (A complex valued function  $f$  on  $V$  is **CONJUGATE LINEAR** if  $f(\mathbf{x} + \mathbf{y}) = f(\mathbf{x}) + f(\mathbf{y})$  and  $f(\alpha \mathbf{x}) = \overline{\alpha} f(\mathbf{x})$  for all  $\mathbf{x}, \mathbf{y} \in V$  and  $\alpha \in \mathbb{C}$ .) When a mapping is linear in one variable and conjugate linear in the other, it is often called **SESQUILINEAR** (the prefix “sesqui-” means “one and a half”). Taken together conditions (a)–(d) say that the inner product is a *positive definite conjugate symmetric sesquilinear form*. When a vector space has been equipped with an inner product we define the **NORM** (or **LENGTH**) of a vector  $\mathbf{x}$  by

$$\|\mathbf{x}\| := \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle};$$

(Notice that this is same definition we used in 4.1.2 for vectors in  $\mathbb{R}^n$ .)

**18.1.2. Notation.** There are many more or less standard notations for the inner product of two vectors  $\mathbf{x}$  and  $\mathbf{y}$ . The two that we will use interchangeably in these exercises are  $\mathbf{x} \cdot \mathbf{y}$  and  $\langle \mathbf{x}, \mathbf{y} \rangle$ .

**18.1.3. Example.** For vectors  $x = (x_1, x_2, \dots, x_n)$  and  $y = (y_1, y_2, \dots, y_n)$  belonging to  $\mathbb{R}^n$  define

$$\langle x, y \rangle = \sum_{k=1}^n x_k y_k.$$

Then  $\mathbb{R}^n$  is an inner product space.

**18.1.4. Example.** For vectors  $x = (x_1, x_2, \dots, x_n)$  and  $y = (y_1, y_2, \dots, y_n)$  belonging to  $\mathbb{C}^n$  define

$$\langle x, y \rangle = \sum_{k=1}^n x_k \overline{y_k}.$$

Then  $\mathbb{C}^n$  is an inner product space.

**18.1.5. Example.** Let  $l_2$  be the set of all square summable sequences of complex numbers. (A sequence  $x = (x_k)_{k=1}^{\infty}$  is SQUARE SUMMABLE if  $\sum_{k=1}^{\infty} |x_k|^2 < \infty$ .) (The vector space operations are defined pointwise.) For vectors  $x = (x_1, x_2, \dots)$  and  $y = (y_1, y_2, \dots)$  belonging to  $l_2$  define

$$\langle x, y \rangle = \sum_{k=1}^{\infty} x_k \overline{y_k}.$$

Then  $l_2$  is an inner product space. (It is important to recognize that in order for this definition to make sense, it must be verified that the infinite series actually converges.)

**18.1.6. Example.** For  $a < b$  let  $\mathcal{C}([a, b])$  be the family of all continuous complex valued functions on the interval  $[a, b]$ . For every  $f, g \in \mathcal{C}([a, b])$  define

$$\langle f, g \rangle = \int_a^b f(x) \overline{g(x)} dx.$$

Then  $\mathcal{C}([a, b])$  is an inner product space.

**18.1.7. Definition.** Let  $\mathbf{x}$  and  $\mathbf{y}$  be nonzero vectors in a real vector space  $V$ . Then  $\angle(\mathbf{x}, \mathbf{y})$ , the ANGLE between  $\mathbf{x}$  and  $\mathbf{y}$ , is defined by

$$\angle(\mathbf{x}, \mathbf{y}) = \arccos \frac{\langle \mathbf{x}, \mathbf{y} \rangle}{\|\mathbf{x}\| \|\mathbf{y}\|}$$

(Notice that this is the same definition as the one given in 4.1.3 for vectors in  $\mathbb{R}^n$ .)

**18.1.8. Definition.** Two vectors  $\mathbf{x}$  and  $\mathbf{y}$  in an inner product space  $V$  are ORTHOGONAL (or PERPENDICULAR) if  $\langle \mathbf{x}, \mathbf{y} \rangle = 0$ . In this case we write  $\mathbf{x} \perp \mathbf{y}$ . Similarly if  $M$  and  $N$  are nonempty subsets of  $V$  we write  $M \perp N$  if every vector in  $M$  is orthogonal to every vector in  $N$ . When one of the sets consists of a single vector we write  $x \perp N$  instead of  $\{x\} \perp N$ . When  $M$  is a nonempty subset of  $V$  we denote by  $M^\perp$  the set of all vectors  $x$  such that  $x \perp N$ . This is the ORTHOGONAL COMPLEMENT of  $M$ .

**18.1.9. Definition.** A real valued function  $f$  on an inner product space  $V$  is UNIFORMLY CONTINUOUS if for every number  $\epsilon > 0$  there exists a number  $\delta > 0$  such that  $|f(x) - f(y)| < \epsilon$  whenever  $\|x - y\| < \delta$  in  $V$ .

The following result is one we have already seen for  $\mathbb{R}^n$  (see 4.1.4).

**18.1.10. Theorem** (Cauchy-Schwarz inequality). *If  $\mathbf{x}$  and  $\mathbf{y}$  are vectors in a (real or complex) inner product space, then*

$$|\langle \mathbf{x}, \mathbf{y} \rangle| \leq \|\mathbf{x}\| \|\mathbf{y}\|.$$

## 18.2. Exercises

- (1) In  $\mathbb{C}^3$  let  $\mathbf{x} = (3 + 2i, 1 - i, i)$  and  $\mathbf{y} = (1 - 2i, 1 - i, 4 + i)$ . Then  $\|\mathbf{x}\| = \underline{\hspace{1cm}}$  ;  $\|\mathbf{y}\| = \underline{\hspace{1cm}}$  ; and  $\langle \mathbf{x}, \mathbf{y} \rangle = \underline{\hspace{1cm}}$  .
- (2) In  $\mathbb{C}^2$  let  $\mathbf{x} = (2 - 4i, 4i)$  and  $\mathbf{y} = (2 + 4i, 4)$ . Then  $\|\mathbf{x}\| = \underline{\hspace{1cm}}$  ;  $\|\mathbf{y}\| = \underline{\hspace{1cm}}$  ; and  $\langle \mathbf{x}, \mathbf{y} \rangle = \underline{\hspace{1cm}}$  .
- (3) In  $\mathbb{C}^2$  let  $\mathbf{x} = (3 - 2i, \sqrt{3}i)$  and  $\mathbf{y} = (1 + i, 1 - i)$ . Then  $\|\mathbf{x}\| = \underline{\hspace{1cm}}$  ;  $\|\mathbf{y}\| = \underline{\hspace{1cm}}$  ; and  $\langle \mathbf{x}, \mathbf{y} \rangle = 1 - \sqrt{a} + (\sqrt{a} - b)$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$  .
- (4) Make the family of  $2 \times 2$  matrices of real numbers into an inner product space by defining  $\langle A, B \rangle := \text{tr}(A^t B)$  (see problem 7) for  $A, B \in \mathbf{M}_{2,2}(\mathbb{R})$ . Let  $U = \begin{bmatrix} 1 & 4 \\ -3 & 5 \end{bmatrix}$  and  $V = \begin{bmatrix} \alpha^2 & \alpha - 1 \\ \alpha + 1 & -1 \end{bmatrix}$ . Find all values of  $\alpha$  such that  $U \perp V$  in the inner product space  $\mathbf{M}_{2,2}(\mathbb{R})$ .  
Answer:  $\alpha = \underline{\hspace{1cm}}$  and  $\underline{\hspace{1cm}}$  .
- (5) Let  $\mathbf{w} = \left( \frac{i}{\sqrt{3}}, \frac{i}{\sqrt{3}}, \frac{i}{\sqrt{3}} \right)$ ,  $\mathbf{x} = \left( -\frac{2i}{\sqrt{6}}, \frac{i}{\sqrt{6}}, \frac{i}{\sqrt{6}} \right)$ ,  $\mathbf{y} = \left( \frac{i}{\sqrt{6}}, \frac{i}{\sqrt{6}}, -\frac{2i}{\sqrt{6}} \right)$ , and  $\mathbf{z} = \left( 0, -\frac{i}{\sqrt{2}}, \frac{i}{\sqrt{2}} \right)$ .
- (a) Which three of these vectors form an orthonormal basis for  $\mathbb{C}^3$ ?  
Answer:  $\underline{\hspace{1cm}}$  ,  $\underline{\hspace{1cm}}$  , and  $\underline{\hspace{1cm}}$  .
- (b) Write  $(1, 0, 0)$  as a linear combination of the three basis vectors you chose in part (a). (Use 0 as the coefficient of the vector which does *not* belong to the basis.)  
Answer:  $(1, 0, 0) = -\frac{i}{\sqrt{a}} \mathbf{w} + \frac{2i}{\sqrt{2a}} \mathbf{x} + b\mathbf{y} + c\mathbf{z}$  where  $a = \underline{\hspace{1cm}}$  ,  $b = \underline{\hspace{1cm}}$  , and  $c = \underline{\hspace{1cm}}$  .
- (6) Find all real numbers  $\alpha$  such that the angle between the vectors  $2\mathbf{i} + 2\mathbf{j} + (\alpha - 2)\mathbf{k}$  and  $2\mathbf{i} + (\alpha - 2)\mathbf{j} + 2\mathbf{k}$  is  $\frac{\pi}{3}$ . Answer:  $\alpha = \underline{\hspace{1cm}}$  and  $\underline{\hspace{1cm}}$  .
- (7) Let  $\mathbf{f}(x) = x$  and  $\mathbf{g}(x) = x^2$  for  $0 \leq x \leq 1$ . Then the cosine of the angle between  $\mathbf{f}$  and  $\mathbf{g}$  in the inner product space  $\mathcal{C}([0, 1])$  of all continuous real valued functions on  $[0, 1]$  is  $\frac{a}{4}$  where  $a = \underline{\hspace{1cm}}$  .
- (8) Let  $\mathbf{f}(x) = x$  and  $\mathbf{g}(x) = \cos \pi x$  for  $0 \leq x \leq 1$ . In the inner product space  $\mathcal{C}([0, 1])$  of all continuous real valued functions on  $[0, 1]$  the cosine of the angle between  $\mathbf{f}$  and  $\mathbf{g}$  is  $-\frac{a\sqrt{b}}{c^2}$  where  $a = \underline{\hspace{1cm}}$  ,  $b = \underline{\hspace{1cm}}$  , and  $c = \underline{\hspace{1cm}}$  .
- (9) Let  $\mathbf{f}(x) = x^2$  and  $\mathbf{g}(x) = 1 - cx$  where  $0 \leq x \leq 1$  and  $c$  is a constant. If  $c = \underline{\hspace{1cm}}$  , then  $\mathbf{f} \perp \mathbf{g}$  in the inner product space  $\mathcal{C}([0, 1])$  of continuous real valued function on  $[0, 1]$ .
- (10) In  $\mathbb{R}^4$  the subspace perpendicular to both  $(1, 4, 4, 1)$  and  $(2, 9, 8, 2)$  is the span of the vectors  $(-4, a, b, a)$  and  $(-b, a, a, b)$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$  .
- (11) A complex valued function  $\mathbf{f}$  on the interval  $[-\pi, \pi]$  is said to be SQUARE INTEGRABLE if  $\int_{-\pi}^{\pi} |\mathbf{f}(x)|^2 dx < \infty$ . Let  $\mathcal{F}([-\pi, \pi])$  be the family of all square integrable complex valued functions on  $[-\pi, \pi]$ . This family is an inner product space under the usual pointwise operations of addition and scalar multiplication and the inner product defined by

$$\langle \mathbf{f}, \mathbf{g} \rangle = \frac{1}{2\pi} \int_{-\pi}^{\pi} \mathbf{f}(x) \overline{\mathbf{g}(x)} dx$$

for all  $\mathbf{f}, \mathbf{g} \in \mathcal{F}([-\pi, \pi])$ . Actually the preceding sentence is a lie: to be correct we should identify any two square integrable functions which differ only on a set of Lebesgue

measure zero and work with the resulting equivalence classes. *Then* we have an inner product space. What we actually have here is a so-called *semi-inner product*. For the purposes of the current exercise, however, this correction turns out to be unimportant; ignore it.

For each integer  $n$  (positive, negative, or zero) define a function  $\mathbf{e}_n$  by

$$\mathbf{e}_n(x) = e^{inx} \quad \text{for } -\pi \leq x \leq \pi.$$

- (a) Then each  $\mathbf{e}_n$  belongs to  $\mathcal{F}([-\pi, \pi])$  and  $\|\mathbf{e}_n\| = \underline{\hspace{1cm}}$  for every integer  $n$ .  
 (b) If  $m \neq n$ , then  $\langle \mathbf{e}_m, \mathbf{e}_n \rangle = \underline{\hspace{1cm}}$ .

$$\text{Now let } \mathbf{f}(x) = \begin{cases} 0 & \text{if } -\pi \leq x < 0, \\ 1 & \text{if } 0 \leq x \leq \pi. \end{cases}$$

- (c) Then  $\langle \mathbf{f}, \mathbf{e}_0 \rangle = \underline{\hspace{1cm}}$ .  
 (d) If  $n$  is odd, then  $\langle \mathbf{f}, \mathbf{e}_n \rangle = \underline{\hspace{1cm}}$ .  
 (e) If  $n$  is even but not zero, then  $\langle \mathbf{f}, \mathbf{e}_n \rangle = \underline{\hspace{1cm}}$ .  
 (f) Write the sum of the middle eleven terms of the Fourier series for  $\mathbf{f}$  in simplified form.

*Hint:* Use problem 8 in this chapter.

$$\text{Answer: } \sum_{n=-5}^5 \langle \mathbf{f}, \mathbf{e}_n \rangle \mathbf{e}_n = a + b \sin x + c \sin 3x + d \sin 5x \text{ where } a = \underline{\hspace{1cm}}, b = \underline{\hspace{1cm}}, \\ c = \underline{\hspace{1cm}}, \text{ and } d = \underline{\hspace{1cm}}.$$

## 18.3. Problems

- (1) Prove that if  $\mathbf{x} \in V$  and  $\langle \mathbf{x}, \mathbf{y} \rangle = 0$  for all  $\mathbf{y} \in V$ , then  $\mathbf{x} = \mathbf{0}$ .
- (2) Let  $S, T: V \rightarrow W$  be linear transformations between real inner product spaces. Prove that if  $\langle S\mathbf{v}, \mathbf{w} \rangle = \langle T\mathbf{v}, \mathbf{w} \rangle$  for all  $\mathbf{v} \in V$  and  $\mathbf{w} \in W$ , then  $S = T$ .
- (3) Prove that if  $V$  is a complex inner product space and  $T \in \mathfrak{L}(V)$  satisfies  $\langle T\mathbf{z}, \mathbf{z} \rangle = 0$  for all  $\mathbf{z} \in V$ , then  $T = 0$ . *Hint.* In the hypothesis replace  $\mathbf{z}$  first by  $\mathbf{x} + \mathbf{y}$  and then by  $\mathbf{x} + i\mathbf{y}$ .
- (4) Show that the preceding result does not hold for real inner product spaces.
- (5) Let  $V$  be a complex inner product space and  $S, T \in \mathfrak{L}(V)$ . Prove that if  $\langle S\mathbf{x}, \mathbf{x} \rangle = \langle T\mathbf{x}, \mathbf{x} \rangle$  for all  $\mathbf{x} \in V$ , then  $S = T$ .
- (6) Prove the *Schwarz inequality* 18.1.10. *Hint.* Let  $\alpha = \langle \mathbf{y}, \mathbf{y} \rangle$  and  $\beta = -\langle \mathbf{x}, \mathbf{y} \rangle$  and expand  $\|\alpha\mathbf{x} + \beta\mathbf{y}\|^2$ . This hint leads to a slick, easy, and totally unenlightening proof. Perhaps you can find a more perspicuous one.

the polarization identity If  $x$  and  $y$  are vectors in a complex inner product space, then

$$\langle x, y \rangle = \frac{1}{4}(\|x + y\|^2 - \|x - y\|^2 + i\|x + iy\|^2 - i\|x - iy\|^2).$$

What is the correct identity for a *real* inner product space?

- (7) Let  $\mathbf{M}_{2,2}(\mathbb{R})$  be the vector space of all  $2 \times 2$  matrices of real numbers. Show that this space can be made into an inner product space by defining  $\langle A, B \rangle := \text{tr}(A^t B)$  for all  $A, B \in \mathbf{M}_{2,2}(\mathbb{R})$ .
- (8) Prove that for every real number  $\theta$

$$e^{i\theta} = \cos \theta + i \sin \theta.$$

Derive from this that  $\cos \theta = \frac{1}{2}(e^{i\theta} + e^{-i\theta})$  and  $\sin \theta = \frac{1}{2i}(e^{i\theta} - e^{-i\theta})$ .

- (9) Let  $\mathbf{x}$  and  $\mathbf{y}$  be vectors in an inner product space. Prove that

$$\|\mathbf{x} + \mathbf{y}\|^2 + \|\mathbf{x} - \mathbf{y}\|^2 = 2\|\mathbf{x}\|^2 + 2\|\mathbf{y}\|^2.$$

Give a geometric interpretation of this result.

- (10) Show that the norm function  $\|\cdot\|: V \rightarrow \mathbb{R}$  on an inner product space is uniformly continuous.

**18.4. Answers to Odd-Numbered Exercises**

- (1)  $4, 2\sqrt{6}, 2 + 12i$
- (3)  $4, 2, 3, 5$
- (5) (a)  $w, x, y$   
(b)  $3, 0, 0$
- (7)  $\sqrt{15}$
- (9)  $\frac{4}{3}$
- (11) (a)  $1$   
(b)  $0$   
(c)  $\frac{1}{2}$   
(d)  $-\frac{i}{n\pi}$   
(e)  $0$   
(f)  $\frac{1}{2}, \frac{2}{\pi}, \frac{2}{3\pi}, \frac{2}{5\pi}$

## ORTHONORMAL SETS OF VECTORS

### 19.1. Background

**Topics:** orthonormal sets of vectors, orthonormal bases, orthogonal complements, orthogonal direct sums, *Gram-Schmidt orthonormalization*, the *QR-factorization* of a matrix.

**19.1.1. Definition.** A set  $B$  of vectors in an inner product space is ORTHONORMAL if  $\mathbf{x} \perp \mathbf{y}$  whenever  $\mathbf{x}$  and  $\mathbf{y}$  are distinct vectors in  $B$  and  $\|\mathbf{x}\| = 1$  for every  $\mathbf{x} \in B$ . The set  $B$  is a MAXIMAL!ORTHONORMAL SET provided that it is orthonormal and the only orthonormal set which contains  $B$  is  $B$  itself.

**19.1.2. Theorem.** Let  $B = \{\mathbf{e}^1, \dots, \mathbf{e}^n\}$  be an orthonormal set in an inner product space  $V$ . Then the following are equivalent.

- (a)  $B$  is a maximal orthonormal set in  $V$ .
- (b) If  $\langle \mathbf{x}, \mathbf{e}^k \rangle = 0$  for  $k = 1, \dots, n$ , then  $\mathbf{x} = \mathbf{0}$ .
- (c) The span of  $B$  is all of  $V$ .
- (d) If  $\mathbf{x} \in V$ , then  $\mathbf{x} = \sum_{k=1}^n \langle \mathbf{x}, \mathbf{e}^k \rangle \mathbf{e}^k$ . (the Fourier series for  $\mathbf{x}$ .)
- (e)  $\langle \mathbf{x}, \mathbf{y} \rangle = \sum_{k=1}^n \langle \mathbf{x}, \mathbf{e}^k \rangle \langle \mathbf{e}^k, \mathbf{y} \rangle$  for all  $\mathbf{x}, \mathbf{y} \in V$ .
- (f)  $\|\mathbf{x}\|^2 = \sum_{k=1}^n |\langle \mathbf{x}, \mathbf{e}^k \rangle|^2$  for every  $\mathbf{x} \in V$ .
- (g)  $\dim V = n$ .

**19.1.3. Definition.** An orthonormal set in an inner product space  $V$  which satisfies any (and hence all) of the conditions listed in the preceding theorem is called an ORTHONORMAL BASIS for  $V$ .

**19.1.4. Definition.** An  $n \times n$  matrix of real numbers is an ORTHOGONAL MATRIX if its column vectors are an orthonormal basis for  $\mathbb{R}^n$ . An  $n \times n$  matrix of complex numbers is a UNITARY MATRIX if its column vectors are an orthonormal basis for  $\mathbb{C}^n$ .

**19.1.5. Theorem (QR-factorization).** If  $A$  is an  $n \times n$  matrix of real numbers, then there exist an orthogonal matrix  $Q$  and an upper triangular matrix  $R$  such that  $A = QR$ . If  $A$  is an  $n \times n$  matrix of complex numbers, then there exist a unitary matrix  $Q$  and an upper triangular matrix  $R$  such that  $A = QR$ .

(For a proof of the preceding theorem see [9], pages 425–427.)

**19.1.6. Definition.** Let  $M$  and  $N$  be subspaces of an inner product space  $V$ . We say that the space  $V$  is the ORTHOGONAL DIRECT SUM of  $M$  and  $N$  if  $M + N = V$  and  $M \perp N$ . In this case we write

$$V = M \oplus N.$$

**Caution:** Since the same notation is used for direct sums of vector spaces and orthogonal direct sums of inner product spaces, close attention should be paid to the context in which these concepts arise. For example, if  $M$  is the  $x$ -axis and  $N$  is the line  $y = x$  in  $\mathbb{R}^2$ , is it true that  $\mathbb{R}^2 = M \oplus N$ ? Yes, if  $\mathbb{R}^2$  is regarded as a vector space. No, if it is regarded as an inner product space.

## 19.2. Exercises

- (1) Use the *Gram-Schmidt procedure* to find an orthonormal basis for the subspace of  $\mathbb{R}^4$  spanned by  $\mathbf{w}^1 = (1, 0, 0, 0)$ ,  $\mathbf{w}^2 = (1, 1, 1, 0)$ , and  $\mathbf{w}^3 = (1, 2, 0, 1)$ . The basis consists of the vectors  $\mathbf{e}^1 = (1, 0, 0, 0)$ ;  $\mathbf{e}^2 = \frac{1}{a}(0, 1, 1, b)$ ; and  $\mathbf{e}^3 = \frac{1}{c}(b, 1, -1, 1)$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ , and  $c = \underline{\hspace{1cm}}$ .
- (2) Let  $\mathcal{P}_4 = \mathcal{P}_4([0, 1])$  be the vector space of polynomials of degree strictly less than 4 with an inner product defined by

$$\langle \mathbf{p}, \mathbf{q} \rangle = \int_0^1 \mathbf{p}(x)\mathbf{q}(x) dx$$

for all  $\mathbf{p}, \mathbf{q} \in \mathcal{P}_4$ . Let  $\mathbf{w}^0(x) = 1$ ,  $\mathbf{w}^1(x) = x$ ,  $\mathbf{w}^2(x) = x^2$ , and  $\mathbf{w}^3(x) = x^3$  for  $0 \leq x \leq 1$ . Use the *Gram-Schmidt process* to convert the ordered basis  $\{\mathbf{w}^0, \mathbf{w}^1, \mathbf{w}^2, \mathbf{w}^3\}$  to an orthonormal basis  $\{\mathbf{e}^0, \mathbf{e}^1, \mathbf{e}^2, \mathbf{e}^3\}$  for  $\mathcal{P}_4$ .

Answer:  $\mathbf{e}^0(x) = \underline{\hspace{1cm}}$

$\mathbf{e}^1(x) = \sqrt{a}(bx - 1)$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$

$\mathbf{e}^2(x) = \sqrt{a}(bx^2 - bx + 1)$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$

$\mathbf{e}^3(x) = \sqrt{a}(bx^3 - cx^2 + dx - 1)$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ ,  
 $c = \underline{\hspace{1cm}}$ , and  $d = \underline{\hspace{1cm}}$

- (3) Find the *QR* factorization of  $A = \begin{bmatrix} 3 & 0 \\ 4 & 5 \end{bmatrix}$ .

Answer:  $A = QR = \frac{1}{a} \begin{bmatrix} b & -c \\ c & b \end{bmatrix} \begin{bmatrix} a & c \\ 0 & b \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ , and  $c = \underline{\hspace{1cm}}$ .

- (4) Let  $A = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}$ . The *QR*-factorization of  $A$  is  $A = QR$

where  $Q = \begin{bmatrix} & & \\ & & \\ & & \end{bmatrix}$  and  $R = \begin{bmatrix} & & \\ & & \\ & & \end{bmatrix}$ .

- (5) Let  $A = \begin{bmatrix} 1 & 4 & -2 \\ 1 & 3 & -1 \\ 1 & 2 & -1 \end{bmatrix}$ . The *QR*-factorization of  $A$  is  $A = QR$

where  $Q = \begin{bmatrix} a & b & -ab \\ a & 0 & 2ab \\ a & -b & -ab \end{bmatrix}$  and  $R = \begin{bmatrix} 3a & 9a & -4a \\ 0 & 2b & -b \\ 0 & 0 & ab \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .



## 19.3. Problems

- (1) Let  $\{\mathbf{e}^1, \mathbf{e}^2, \dots, \mathbf{e}^n\}$  be a finite orthonormal subset of an inner product space  $V$  and  $\mathbf{x} \in V$ . Show that

$$\sum_{k=1}^n |\langle \mathbf{x}, \mathbf{e}^k \rangle|^2 \leq \|\mathbf{x}\|^2.$$

*Hint.* Multiply out  $\langle \mathbf{x} - \sum_{k=1}^n \langle \mathbf{x}, \mathbf{e}^k \rangle \mathbf{e}^k, \mathbf{x} - \sum_{k=1}^n \langle \mathbf{x}, \mathbf{e}^k \rangle \mathbf{e}^k \rangle$ .

- (2) Let  $M$  be a subspace of an inner product space  $V$ .  
 (a) Show that  $M \subseteq M^{\perp\perp}$ .  
 (b) Prove that equality need not hold in (a).  
 (c) Show that if  $V$  is finite dimensional, then  $M = M^{\perp\perp}$ .  
 (3) Let  $M$  and  $N$  be subspaces of an inner product space. Prove that

$$(M + N)^{\perp} = M^{\perp} \cap N^{\perp}.$$

- (4) Let  $M$  be a subspace of a finite dimensional inner product space  $V$ . Prove that  $V = M \oplus M^{\perp}$ .  
 (5) Give an example to show that the conclusion of the preceding problem need not hold in an infinite dimensional space.  
 (6) Prove that if an inner product space  $V$  is the orthogonal direct sum  $M \oplus N$  of two subspaces  $M$  and  $N$ , then  $N = M^{\perp}$ .  
 (7) Prove that if  $f$  is a linear functional on a finite dimensional inner product space  $V$ , then there exists a unique vector  $\mathbf{a} \in V$  such that

$$f(\mathbf{x}) = \langle \mathbf{x}, \mathbf{a} \rangle$$

for every  $\mathbf{x} \in V$ .

- (8) In beginning calculus you found (by making use of the  $p$ -test) that the series  $\sum_{k=1}^{\infty} \frac{1}{k^2}$  converges. But you were not given a means of discovering what the series converges to. Now you have enough machinery to accomplish this.

We denote by  $L_2[0, 2\pi]$  the vector space of all complex valued functions  $\mathbf{f}$  defined on the interval  $[0, 2\pi]$  such that

$$\int_0^{2\pi} |\mathbf{f}(t)|^2 dt < \infty.$$

(As in exercise 11 of chapter 18 this isn't quite correct: the members of  $L_2$  are technically equivalence classes of functions. For the purposes of this problem, use the preceding not-quite-right definition.)

On the space  $L_2[0, 2\pi]$  we define the following inner product

$$\langle \mathbf{f}, \mathbf{g} \rangle = \frac{1}{2\pi} \int_0^{2\pi} \mathbf{f}(t) \overline{\mathbf{g}(t)} dt.$$

For each integer  $n$  (positive, negative, or zero) define the function  $\mathbf{e}^n$  on  $[0, 2\pi]$  by

$$\mathbf{e}^n(x) = e^{inx}$$

for  $0 \leq x \leq 2\pi$ .

- (a) Show that  $\{\mathbf{e}^n : n \text{ is an integer}\}$  is an orthonormal set in  $L_2[0, 2\pi]$ .

In part (b) you may use without proof the following fact: for every function  $\mathbf{f}$  in the inner product space  $L_2[0, 2\pi]$

$$\|\mathbf{f}\|^2 = \sum_{k=-\infty}^{\infty} |\langle \mathbf{f}, \mathbf{e}^k \rangle|^2 \quad (*)$$

That is, in  $L_2[0, 2\pi]$  the square of the length of a vector is the sum of the squares of its Fourier coefficients with respect to the orthonormal family given in part (a). This is the infinite dimensional version of *Parseval's formula*.

- (b) Find the sum of the infinite series  $\sum_{k=1}^{\infty} \frac{1}{k^2}$ . *Hint.* Apply (\*) to the function  $\mathbf{f}(x) = x$ .

**19.4. Answers to Odd-Numbered Exercises**

(1)  $\sqrt{2}, 0, \sqrt{3}$

(3) 5, 3, 4

(5)  $\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{2}}$



## QUADRATIC FORMS

### 20.1. Background

**Topics:** quadratic forms, quadric surfaces, positive (and negative) definite, positive (and negative) semidefinite, indefinite.

**20.1.1. Definition.** A symmetric matrix  $A$  on a real inner product  $V$  is **POSITIVE DEFINITE** if  $\langle A\mathbf{x}, \mathbf{x} \rangle > 0$  for every  $\mathbf{x} \neq \mathbf{0}$  in  $V$ . It is **NEGATIVE DEFINITE** if  $\langle A\mathbf{x}, \mathbf{x} \rangle < 0$  for every  $\mathbf{x} \neq \mathbf{0}$  in  $V$ . It is **INDEFINITE** if there are vectors  $\mathbf{x}$  and  $\mathbf{y}$  in  $V$  such that  $\langle A\mathbf{x}, \mathbf{x} \rangle > 0$  and  $\langle A\mathbf{y}, \mathbf{y} \rangle < 0$ . Of course an operator on a finite dimensional vector space is positive definite, negative definite, or indefinite of its matrix representation is positive definite, *etc.*

The following useful result (and its proof) can be found on page 250 of [10].

**20.1.2. Theorem.** *Let  $A$  be an  $n \times n$  matrix. Then the following conditions are equivalent:*

- (a)  *$A$  is positive definite;*
- (b)  *$\mathbf{x}^t A \mathbf{x} > 0$  for every  $\mathbf{x} \neq \mathbf{0}$  in  $\mathbb{R}^n$ ;*
- (c) *every eigenvalue  $\lambda$  of  $A$  is strictly positive;*
- (d) *every leading principal submatrix  $A_k$  ( $k = 1, \dots, n$ ) has strictly positive determinant; and*
- (e) *when  $A$  has been put in echelon form (without row exchanges) the pivots are all strictly positive.*

In the preceding, the *leading principal submatrix*  $A_k$  is the  $k \times k$  matrix which appears in the upper left corner of  $A$ .

## 20.2. Exercises

- (1) Suppose that  $A$  is a  $3 \times 3$  matrix such that  $\langle A\mathbf{x}, \mathbf{x} \rangle = x_1^2 + 5x_2^2 - 3x_3^2 + 6x_1x_2 - 4x_1x_3 + 8x_2x_3$  for all  $\mathbf{x} \in \mathbb{R}^3$ . Then  $A = \begin{bmatrix} a & b & c \\ b & d & e \\ c & e & f \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ ,  $c = \underline{\hspace{1cm}}$ ,  $d = \underline{\hspace{1cm}}$ ,  $e = \underline{\hspace{1cm}}$ , and  $f = \underline{\hspace{1cm}}$ .
- (2) A curve  $C$  is given by the equation  $2x^2 - 72xy + 23y^2 = 50$ . What kind of curve is  $C$ ?  
Answer: It is a(n)  $\underline{\hspace{3cm}}$ .
- (3) The equation  $5x^2 + 8xy + 5y^2 = 1$  describes an ellipse. The principal axes of the ellipse lie along the lines  $y = \underline{\hspace{1cm}}$  and  $y = \underline{\hspace{1cm}}$ .
- (4) The graph of the equation  $13x^2 - 8xy + 7y^2 = 45$  is an ellipse. The length of its semimajor axis is  $\underline{\hspace{1cm}}$  and the length of its semiminor axis is  $\underline{\hspace{1cm}}$ .
- (5) Consider the equation  $2x^2 + 2y^2 - z^2 - 2xy + 4xz + 4yz = 3$ .
- (a) The graph of the equation is what type of quadric surface?  
Answer:  $\underline{\hspace{3cm}}$ .
- (b) In standard form the equation for this surface is  $u^2 + v^2 + \underline{\hspace{1cm}} w^2 = \underline{\hspace{1cm}}$ .
- (c) Find three orthonormal vectors with the property that in the coordinate system they generate, the equation of the surface is in standard form.  
Answer:  $\frac{1}{\sqrt{6}}(1, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})$ ,  $\frac{1}{\sqrt{2}}(\underline{\hspace{1cm}}, \underline{\hspace{1cm}}, 0)$ , and  $\frac{1}{\sqrt{3}}(1, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})$ .
- (6) Determine for each of the following matrices whether it is positive definite, positive semidefinite, negative definite, negative semidefinite, or indefinite.
- (a) The matrix  $\begin{bmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{bmatrix}$  is  $\underline{\hspace{3cm}}$ .
- (b) The matrix  $\begin{bmatrix} -2 & 1 & 1 \\ 1 & -2 & -1 \\ 1 & -1 & -2 \end{bmatrix}$  is  $\underline{\hspace{3cm}}$ .
- (7) Determine for each of the following matrices whether it is positive definite, positive semidefinite, negative definite, negative semidefinite, or indefinite.
- (a) The matrix  $\begin{bmatrix} 1 & 2 & 3 \\ 2 & 5 & 4 \\ 3 & 4 & 9 \end{bmatrix}$  is  $\underline{\hspace{3cm}}$ .
- (b) The matrix  $\begin{bmatrix} 1 & 2 & 0 & 0 \\ 2 & 6 & -2 & 0 \\ 0 & -2 & 5 & -2 \\ 0 & 0 & -2 & 3 \end{bmatrix}$  is  $\underline{\hspace{3cm}}$ .
- (c) The matrix  $\begin{bmatrix} 0 & 1 & 2 \\ 1 & 0 & 1 \\ 2 & 1 & 0 \end{bmatrix}^2$  is  $\underline{\hspace{3cm}}$ .
- (8) Let  $B = \begin{bmatrix} 2 & 2 & 4 \\ 2 & b & 8 \\ 4 & 8 & 7 \end{bmatrix}$ . For what range of values of  $b$  is  $B$  positive definite?  
Answer:  $\underline{\hspace{3cm}}$ .

- (9) Let  $A = \begin{bmatrix} a & 1 & 1 \\ 1 & a & 1 \\ 1 & 1 & a \end{bmatrix}$ . For what range of values of  $a$  is  $A$  positive definite?

Answer: \_\_\_\_\_ .

**20.3. Problems**

- (1) You are given a quadratic form  $q(x, y, z) = ax^2 + by^2 + cz^2 + 2dxy + 2exz + 2fyz$ . Explain in detail how to determine whether the associated level surface  $q(x, y, z) = c$  encloses a region of finite volume in  $\mathbb{R}^3$  and, if it does, how to find that volume. Justify carefully all claims you make. Among other things, explain how to use the *change of variables theorem* for multiple integrals to express the volume of an ellipsoid in terms of the lengths of the principal axes of the ellipsoid.

Apply the method you have developed to the equation

$$11x^2 + 4y^2 + 11z^2 + 4xy - 10xz + 4yz = 8.$$



**20.4. Answers to Odd-Numbered Exercises**

- (1)  $1, 3, -2, 5, 4, -3$
- (3)  $-x, x$
- (5) (a) hyperboloid of one sheet  
(b)  $-1, 1$   
(c)  $1, -2, 1, -1, 1, 1$
- (7) (a) indefinite  
(b) positive definite  
(c) positive definite
- (9)  $a > 1$



## CHAPTER 21

# OPTIMIZATION

### 21.1. Background

**Topics:** critical (stationary) points of a function of several variables; local (relative) maxima and minima; global (absolute) maxima and minima.

**21.1.1. Definition.** Let  $f: \mathbb{R}^n \rightarrow \mathbb{R}$  be a smooth scalar field (that is, a real valued function on  $\mathbb{R}^n$  with derivatives of all orders) and  $\mathbf{p} \in \mathbb{R}^n$ . The HESSIAN MATRIX (or SECOND DERIVATIVE MATRIX) of  $f$  at  $\mathbf{p}$ , denoted by  $H_f(\mathbf{p})$ , is the symmetric  $n \times n$  matrix

$$H_f(\mathbf{p}) = \left[ \frac{\partial^2 f}{\partial x_i \partial x_j}(\mathbf{p}) \right]_{i=1, j=1}^n = [f_{ij}(\mathbf{p})].$$

**21.1.2. Theorem** (Second Derivative Test). *Let  $\mathbf{p}$  be a critical point of a smooth scalar field  $f$  (that is, a point where the gradient of  $f$  is zero). If the Hessian matrix  $H_f$  is positive definite at  $\mathbf{p}$ , then  $f$  has a local minimum there. If  $H_f$  is negative definite at  $\mathbf{p}$ , then  $f$  has a local maximum there. If  $H_f$  is indefinite at  $\mathbf{p}$ , then  $f$  has a saddle point there.*

**21.2. Exercises**

- (1) Notice that the function  $f$  defined by  $f(x, y) = (x^2 - 2x) \cos y$  has a critical point (stationary point) at the point  $(1, \pi)$ . The eigenvalues of the Hessian matrix of  $f$  are \_\_\_\_\_ and \_\_\_\_\_; so we conclude that the point  $(1, \pi)$  is a \_\_\_\_\_ (local minimum, local maximum, saddle point).

- (2) Use matrix methods to classify the critical point of the function

$$f(x, y) = 2x^2 + 2xy + 2x + y^4 - 4y^3 + 7y^2 - 4y + 5$$

as a local maximum, local minimum, or saddle point.

- (a) The only critical point is located at ( \_\_\_\_\_ , \_\_\_\_\_ ).

- (b) It is a \_\_\_\_\_.

- (3) Use matrix methods to classify the critical point of the function

$$f(x, y, z) = \frac{1}{2}x^4 - xy + y^2 - xz + z^2 - x + 3$$

as a local maximum, local minimum, or saddle point.

- (a) The only critical point is located at ( \_\_\_\_\_ , \_\_\_\_\_ , \_\_\_\_\_ ).

- (b) It is a \_\_\_\_\_.

- (4) Notice that the function  $f$  defined by  $f(x, y) = -1 + 4(e^x - x) - 5x \sin y + 6y^2$  has a critical point (stationary point) at the origin. Since the eigenvalues of the Hessian matrix of  $f$  are \_\_\_\_\_ (both positive, both negative, of different signs) we conclude that the origin is a \_\_\_\_\_ (local minimum, local maximum, saddle point).

- (5) Use matrix methods to classify each critical point of the function

$$f(x, y) = y^3 - \frac{4}{3}x^3 - 2y^2 + 2x^2 + y - 7$$

as a local maximum, local minimum, or saddle point.

Answer:  $(0, \frac{1}{3})$  is a \_\_\_\_\_.

$(0, \text{_____})$  is a \_\_\_\_\_.

$(\text{_____}, \frac{1}{3})$  is a \_\_\_\_\_.

$(\text{_____}, \text{_____})$  is a \_\_\_\_\_.

- (6) Use matrix methods to classify each critical point of the function

$$f(x, y, z) = x^2y - 4x - y \sin z \quad \text{for } 0 < z < \pi$$

as a local maximum, local minimum, or saddle point.

Answer: The critical points are  $(-1, \text{_____}, \text{_____})$ , which is a \_\_\_\_\_; and  $(1, \text{_____}, \text{_____})$ , which is a \_\_\_\_\_ (local minimum, local maximum, saddle point).

- (7) The function  $f$  defined by  $f(x, y) = x^2y^2 - 2x - 2y$  has a stationary point at  $(\text{_____}, \text{_____})$ . At this stationary point  $f$  has a \_\_\_\_\_ (local minimum, local maximum, saddle point).

**21.3. Problems**

- (1) Use matrix methods to classify each critical point of the function

$$f(x, y, z) = x^3y + z^2 - 3x - y + 4z + 5$$

as a local maximum, local minimum, or saddle point. Justify your conclusions carefully.

- (2) Let  $f(x, y, z) = x^2y - ye^z + 2x + z$ . The only critical point of  $f$  is located at  $(-1, 1, 0)$ . Use the second derivative test to classify this point as a local maximum, local minimum, or saddle point. State the reasons for your conclusion clearly.

- (3) Notice that the function  $f$  defined by  $f(x, y, z) = x^2y + 2xy + y - ye^{z-1} + 2x + z + 7$  has a critical point (stationary point) at  $(-2, 1, 1)$ . Use the second derivative test to classify this point as a local maximum, local minimum, or saddle point. State the reasons for your conclusion clearly.

- (4) Explain in detail how to use matrix methods to classify each critical point of the function

$$f(x, y) = -\frac{1}{2}xy + \frac{2}{x} - \frac{1}{y}$$

as a local maximum, local minimum, or saddle point. Carry out the computations you describe.

**21.4. Answers to Odd-Numbered Exercises**

- (1)  $-2, -1$ , local maximum
- (3) (a)  $1, \frac{1}{2}, \frac{1}{2}$   
(b) local minimum
- (5) saddle point, 1, local minimum, 1, local maximum, 1, 1, saddle point
- (7) 1, 1, saddle point

## Part 6

# ADJOINT OPERATORS





## CHAPTER 22

# ADJOINTS AND TRANSPOSES

### 22.1. Background

**Topics:** adjoint of an operator, transpose of an operator, conjugate transpose.

**22.1.1. Definition.** Let  $T: V \rightarrow W$  be a linear transformation between real inner product spaces. If there exists a linear map  $T^t: W \rightarrow V$  which satisfies

$$\langle T\mathbf{v}, \mathbf{w} \rangle = \langle \mathbf{v}, T^t\mathbf{w} \rangle$$

for all  $\mathbf{v} \in V$  and  $\mathbf{w} \in W$ , then  $T^t$  is the TRANSPOSE of  $T$ .

In connection with the definition above see problem 1.

**22.1.2. Theorem.** Let  $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation. Then the transpose linear transformation  $T^t$  exists. Furthermore, the matrix representation  $[T^t]$  of this transformation is the transpose of the matrix representation of  $T$ .

**22.1.3. Definition.** Let  $V$  be a real inner product space and  $T$  be an operator on  $V$  whose transpose exists. If  $T = T^t$ , then  $T$  is SYMMETRIC. If  $T$  commutes with its transpose (that is, if  $TT^t = T^tT$ ) it is NORMAL.

**22.1.4. Definition.** If  $[a_{ij}]$  is an  $m \times n$  matrix, its CONJUGATE TRANSPOSE is the  $n \times m$  matrix  $[\overline{a_{ji}}]$ .

**22.1.5. Definition.** Let  $T: V \rightarrow W$  be a linear transformation between complex inner product spaces. If there exists a linear map  $T^*: W \rightarrow V$  which satisfies

$$\langle T\mathbf{v}, \mathbf{w} \rangle = \langle \mathbf{v}, T^*\mathbf{w} \rangle$$

for all  $\mathbf{v} \in V$  and  $\mathbf{w} \in W$ , then  $T^*$  is the ADJOINT (or CONJUGATE TRANSPOSE, or HERMITIAN CONJUGATE) of  $T$ . (In many places,  $T^*$  is denoted by  $T^H$  or by  $T^\dagger$ .)

**22.1.6. Theorem.** Let  $T: \mathbb{C}^n \rightarrow \mathbb{C}^m$  be a linear transformation. Then the adjoint linear transformation  $T^*$  exists. Furthermore, the matrix representation  $[T^*]$  of this transformation is the conjugate transpose of the matrix representation of  $T$ .

**22.1.7. Definition.** Let  $V$  be a complex inner product space and  $T$  be an operator on  $V$  whose adjoint exists. If  $T = T^*$ , then  $T$  is SELF-ADJOINT (or HERMITIAN). If  $T$  commutes with its adjoint (that is, if  $TT^* = T^*T$ ) it is NORMAL. A matrix is NORMAL if it is the representation of a normal operator.

**22.1.8. Definition.** Let  $V$  and  $W$  be inner product spaces. We make the vector space  $V \oplus W$  into an inner product space as follows. For  $\mathbf{v}_1, \mathbf{v}_2 \in V$  and  $\mathbf{w}_1, \mathbf{w}_2 \in W$  let

$$\langle (\mathbf{v}_1, \mathbf{w}_1), (\mathbf{v}_2, \mathbf{w}_2) \rangle := \langle \mathbf{v}_1, \mathbf{v}_2 \rangle + \langle \mathbf{w}_1, \mathbf{w}_2 \rangle.$$

(It is an easy exercise to verify that this is indeed an inner product on  $V \oplus W$ .)

**22.2. Exercises**

- (1) Let  $\mathcal{C}([0, 1], \mathbb{C})$  be the family of all continuous complex valued functions on the interval  $[0, 1]$ . The usual inner product on this space is given by

$$\langle \mathbf{f}, \mathbf{g} \rangle = \int_0^1 \mathbf{f}(x) \overline{\mathbf{g}(x)} dx.$$

Let  $\phi$  be a fixed continuous complex valued function on  $[0, 1]$ . Define the operator  $M_\phi$  on the complex inner product space  $\mathcal{C}([0, 1], \mathbb{C})$  by  $M_\phi(\mathbf{f}) = \phi \mathbf{f}$ . Then

$$M_\phi^* = \underline{\hspace{2cm}}.$$

- (2) Let  $A = \begin{bmatrix} 3-i & 2+2i \\ \frac{1}{1-i} & 3i \end{bmatrix}$ . Find Hermitian (that is, self-adjoint) matrices  $B$  and  $C$  such that  $A = B + iC$ . *Hint.* Consider  $A \pm A^*$ .

$$\text{Answer: } B = \frac{1}{a} \begin{bmatrix} 4c & b+ci \\ b-ci & d \end{bmatrix} \text{ and } C = \frac{1}{a} \begin{bmatrix} -a & b-ci \\ b+ci & 4c \end{bmatrix}, \text{ where } a = \underline{\hspace{1cm}},$$

$$b = \underline{\hspace{1cm}}, c = \underline{\hspace{1cm}}, \text{ and } d = \underline{\hspace{1cm}}.$$

- (3) Let  $\mathcal{P}_3$  be the space of polynomial functions of degree strictly less than 3 defined on the interval  $[0, 1]$ . Define the inner product of two polynomials  $p, q \in \mathcal{P}_3$  by  $\langle p, q \rangle = \int_0^1 p(t)q(t) dt$ . Then the matrix representation of the transpose of the differentiation operator  $D$  on the

space  $\mathcal{P}_3$  (with respect to its usual basis  $\{1, t, t^2\}$ ) is  $\left[ \begin{array}{ccc} & & \\ & & \\ & & \end{array} \right]$ . *Hint.* The answer is **not** the transpose of the matrix representation of  $D$ .

- (4) Let  $V$  be a complex inner product space. Define an operator  $T: V \oplus V \rightarrow V \oplus V$  by

$$T(\mathbf{x}, \mathbf{y}) = (\mathbf{y}, -\mathbf{x}).$$

Then  $T^*(\mathbf{u}, \mathbf{v}) = (\underline{\hspace{1cm}}, \underline{\hspace{1cm}})$ .

## 22.3. Problems

- (1) Let  $T: V \rightarrow W$  be a linear map between real inner product spaces. If  $S: W \rightarrow V$  is a function which satisfies

$$\langle T\mathbf{v}, \mathbf{w} \rangle = \langle \mathbf{v}, S\mathbf{w} \rangle$$

for all  $\mathbf{v} \in V$  and all  $\mathbf{w} \in W$ , then  $S$  is linear (and is therefore the transpose of  $T$ ).

- (2) Prove theorem 22.1.2. Show that, in fact, every linear map between finite dimensional real inner product spaces has a transpose.
- (3) Let  $T$  be a self-adjoint operator on a complex inner product space  $V$ . Prove that  $\langle T\mathbf{x}, \mathbf{x} \rangle$  is real for every  $\mathbf{x} \in V$ .
- (4) Let  $T$  be an operator on a complex inner product space whose adjoint  $T^*$  exists. Prove that  $T^*T = 0$  if and only if  $T = \mathbf{0}$ .
- (5) Let  $V$  be a complex inner product space and let  $\phi$  be defined on the set  $\mathfrak{A}(V)$  of operators on  $V$  whose adjoint exists by

$$\phi(T) = T^*.$$

Show that if  $S, T \in \mathfrak{A}(V)$  and  $\alpha \in \mathbb{C}$ , then  $(S + T)^* = S^* + T^*$  and  $(\alpha T)^* = \bar{\alpha} T^*$ . *Hint.* Use problem 5 in chapter 18.

**Note:** Similarly, if  $V$  is a real inner product space,  $\mathfrak{A}(V)$  is the set of operators whose transpose exists, and  $\phi(T) := T^t$ , then  $\phi$  is linear.

- (6) Let  $T$  be a linear operator on a complex inner product space  $V$ . Show if  $T$  has an adjoint, then so does  $T^*$  and  $T^{**} = T$ . *Hint:* Use problem 5 in chapter 18. (Here  $T^{**}$  means  $(T^*)^*$ .)

**Note:** The real inner product space version of this result says that if  $T$  is an operator on a real inner product space whose transpose exists, then the transpose of  $T^t$  exists and  $T^{tt} = T$ .

- (7) Let  $S$  and  $T$  be operators on a complex inner product space  $V$ . Show that if  $S$  and  $T$  have adjoints, then so does  $ST$  and  $(ST)^* = T^*S^*$ . *Hint.* Use problem 5 in chapter 18.

**Note:** The real inner product space version of this says that if  $S$  and  $T$  are operators on a real inner product space and if  $S$  and  $T$  both have transposes, then so does  $ST$  and  $(ST)^t = T^t S^t$ .

- (8) Let  $A: V \rightarrow V$  be an operator on a real inner product space. Suppose that  $A^t$  exists and that it commutes with  $A$  (that is, suppose  $AA^t = A^t A$ ). Show that  $\ker A = \ker A^t$ .

- (9) Let  $A$  and  $B$  be Hermitian operators on a complex inner product space. Prove that  $AB$  is Hermitian if and only if  $AB = BA$ .

- (10) Show that if  $T: V \rightarrow W$  is an invertible linear map between complex inner product spaces and both  $T$  and  $T^{-1}$  have adjoints, then  $T^*$  is invertible and  $(T^*)^{-1} = (T^{-1})^*$ .

**Note:** The real inner product space version of this says that if  $T: V \rightarrow W$  is an invertible linear map between real inner product spaces and both  $T$  and  $T^{-1}$  have transposes, then  $T^t$  is invertible and  $(T^t)^{-1} = (T^{-1})^t$ .

- (11) Every eigenvalue of a self-adjoint operator on a complex inner product space is real. *Hint.* Let  $\mathbf{x}$  be an eigenvector associated with an eigenvalue  $\lambda$  of an operator  $A$ . Consider  $\lambda \|\mathbf{x}\|^2$ .

- (12) Let  $A$  be a self-adjoint operator on a complex inner product space. Prove that eigenvectors associated with distinct eigenvalues of  $A$  are orthogonal. *Hint.* Use problem 11. Let  $x$  and  $y$  be eigenvectors associated with distinct eigenvalues  $\lambda$  and  $\mu$  of  $A$ . Start your proof by showing that  $\lambda \langle x, y \rangle = \mu \langle x, y \rangle$ .

**22.4. Answers to Odd-Numbered Exercises**

(1)  $M_{\bar{\phi}}$

(3)  $\begin{bmatrix} -6 & 2 & 3 \\ 12 & -24 & -26 \\ 0 & 30 & 30 \end{bmatrix}$

## THE FOUR FUNDAMENTAL SUBSPACES

### 23.1. Background

**Topics:** column space; row space; nullspace; left nullspace, lead variables and free variables in a matrix, rank, row rank and column rank of a matrix.

**23.1.1. Definition.** A linear transformation  $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$  (and its associated standard matrix  $[T]$ ) have *four* fundamental subspaces: the kernels and ranges of  $T$  and  $T^t$ . Over the years a rather elaborate terminology has grown up around these basic notions.

The NULLSPACE of the matrix  $[T]$  is the kernel of the linear map  $T$ .

The LEFT NULLSPACE of the matrix  $[T]$  is the kernel of the linear map  $T^t$ .

The COLUMN SPACE of  $[T]$  is the subspace of  $\mathbb{R}^m$  spanned by the column vectors of the matrix  $[T]$ . This is just the range of the linear map  $T$ .

And finally, the ROW SPACE of  $[T]$  is the subspace of  $\mathbb{R}^n$  spanned by the row vectors of the matrix  $[T]$ . This is just the range of the linear map  $T^t$ .

For a linear transformation  $T: \mathbb{C}^n \rightarrow \mathbb{C}^m$  the terminology is the same. **EXCEPT:** in the preceding five paragraphs each appearance of “ $T^t$ ” must be replaced by a “ $T^*$ ” (and, of course,  $\mathbb{R}^n$  by  $\mathbb{C}^n$  and  $\mathbb{R}^m$  by  $\mathbb{C}^m$ ).

**23.1.2. Definition.** The ROW RANK of a matrix is the dimension of its row space and the COLUMN RANK of a matrix is the dimension of its column space.

**23.1.3. Proposition.** *The rank of a matrix  $A$  is the dimension of the largest square submatrix of  $A$  with nonzero determinant.*

Two useful facts that you may wish to keep in mind are:

- (i) row equivalent matrices have the same row space (for a proof see [8], page 56); and
- (ii) the row rank of a matrix is the same as its column rank (for a proof see [8], page 72).

Note that according to the second assertion the rank of a linear map  $T$ , the row rank of its matrix representation  $[T]$ , and the column rank of  $[T]$  are all equal.

**23.1.4. Theorem** (Fundamental Theorem of Linear Algebra). *If  $T$  is an operator on a finite dimensional complex inner product space, then*

$$\ker T = (\text{ran } T^*)^\perp.$$

**23.1.5. Corollary.** *If  $T$  is an operator on a finite dimensional complex inner product space, then*

$$\ker T^* = (\text{ran } T)^\perp.$$

**23.1.6. Corollary.** *If  $T$  is an operator on a finite dimensional complex inner product space, then*

$$\text{ran } T = (\ker T^*)^\perp.$$

**23.1.7. Corollary.** *If  $T$  is an operator on a finite dimensional complex inner product space, then*

$$\text{ran } T^* = (\ker T)^\perp.$$

**Note:** With the obvious substitutions of  $T^t$  for  $T^*$ , the preceding theorem and its three corollaries remain true for finite dimensional real inner product spaces.

**23.2. Exercises**

- (1) Let  $T: \mathbb{C}^3 \rightarrow \mathbb{C}^3$  be the operator whose matrix representation is

$$[T] = \begin{bmatrix} 1 & i & -1 \\ 1+i & 3-i & -2 \\ 1-2i & -6+5i & 1 \end{bmatrix}.$$

- (a) The kernel of  $T$  (the nullspace of  $[T]$ ) is the span of

$$\{ ( \rule{1cm}{0.4pt} , \rule{1cm}{0.4pt} , 10 ) \}.$$

- (b) The range of  $T$  (the column space of  $[T]$ ) is the span of

$$\{ ( 1 , 0 , \rule{1cm}{0.4pt} ) , ( 0 , 1 , \rule{1cm}{0.4pt} ) \}.$$

- (c) The kernel of  $T^*$  (the left nullspace of  $T^*$ ) is the span of

$$\{ ( 3 , \rule{1cm}{0.4pt} , \rule{1cm}{0.4pt} ) \}.$$

- (d) The range of  $T^*$  (the row space of  $[T]$ ) is the span of

$$\{ ( 10 , 0 , \rule{1cm}{0.4pt} ) , ( 0 , 10 , \rule{1cm}{0.4pt} ) \}.$$

- (2) Find a basis for each of the four fundamental subspaces associated with the matrix

$$A = \begin{bmatrix} 1 & 2 & 0 & 1 \\ 0 & 1 & 1 & 0 \\ 1 & 2 & 0 & 1 \end{bmatrix}.$$

- (a) The column space of  $A$  is the plane in  $\mathbb{R}^3$  whose equation is  $\rule{2cm}{0.4pt}$ .  
It is the span of  $\{ ( 1 , 0 , \rule{1cm}{0.4pt} ) , ( 0 , 1 , \rule{1cm}{0.4pt} ) \}$ .

- (b) The nullspace of  $A$  is the span of  $\{ ( \rule{1cm}{0.4pt} , -1 , \rule{1cm}{0.4pt} , 0 ) , ( \rule{1cm}{0.4pt} , 0 , 0 , 1 ) \}$ .

- (c) The row space of  $A$  is the span of  $\{ ( 1 , 0 , \rule{1cm}{0.4pt} , 1 ) , ( 0 , \rule{1cm}{0.4pt} , 1 , 0 ) \}$ .

- (d) The left nullspace of  $A$  is the line in  $\mathbb{R}^3$  whose equations are  $\rule{1cm}{0.4pt} = \rule{1cm}{0.4pt} = 0$ .  
It is the span of  $\{ ( \rule{1cm}{0.4pt} , \rule{1cm}{0.4pt} , 1 ) \}$ .

(3) Let  $A = \begin{bmatrix} 1 & 2 & 0 & 2 & -1 & 1 \\ 3 & 6 & 1 & 1 & -2 & 1 \\ 5 & 10 & 1 & 5 & -4 & 3 \end{bmatrix}$ .

- (a) Find a basis for the column space of  $A$ .

Answer:

$$\{ ( 1 , 0 , \rule{1cm}{0.4pt} ) , ( 0 , \rule{1cm}{0.4pt} , 1 ) \}.$$

- (b) The column space of  $A$  is a plane in  $\mathbb{R}^3$ . What is its equation?

Answer:  $\rule{2cm}{0.4pt}$ .

- (c) The dimension of the row space of  $A$  is  $\rule{1cm}{0.4pt}$ .

- (d) Fill in the missing coordinates of the following vector so that it lies in the row space of  $A$ .

$$(4, \rule{1cm}{0.4pt} , 6 , \rule{1cm}{0.4pt} , \rule{1cm}{0.4pt} , \rule{1cm}{0.4pt} ).$$

- (e) The dimension of the nullspace of  $A$  is  $\rule{1cm}{0.4pt}$ .

- (f) Fill in the missing coordinates of the following vector so that it lies in the nullspace of  $A$ .

$$( \rule{1cm}{0.4pt} , 1 , \rule{1cm}{0.4pt} , 1 , 1 , 1 ) .$$

- (4) Let  $T: \mathbb{R}^5 \rightarrow \mathbb{R}^3$  be defined by

$$T(v, w, x, y, z) = (v - x + z, v + w - y, w + x - y - z).$$

- (a) Find the matrix representation of  $T$ .

- (b) The kernel of  $T$  (the nullspace of  $[T]$ ) is the span of

$$\{ (\_, \_, 1, 0, 0), (0, 1, 0, 1, 0), (-1, 1, 0, \_, 1) \}.$$

- (c) The range of  $T$  (the column space of  $[T]$ ) is the span of

$$\{ (1, \_, -1), (0, 1, \_) \}.$$

Geometrically this is a \_\_\_\_\_.

- (d) The range of  $T^t$  (the row space of  $[T]$ ) is the span of

$$\{ (1, 0, \_, 0, 1), (\_, \_, \_, -1, -1) \}.$$

- (e) The kernel of  $T^t$  (the left nullspace of  $[T]$ ) is the span of

$$\{ (1, \_, \_) \}.$$

Geometrically this is a \_\_\_\_\_.

- (5) Let  $A$  be the matrix  $\begin{bmatrix} 1 & 0 & 2 & 0 & -1 \\ 1 & 2 & 4 & -2 & -1 \\ 0 & 1 & 1 & -1 & 0 \\ 2 & 3 & 7 & -3 & -2 \end{bmatrix}$ . Find the following subspaces associated with  $A$ .

- (a) The column space of  $A$  is the span of

$$\{ (1, 0, -1/2, \_, \_), (0, 1, \_, \_, \_) \}.$$

- (b) The row space of  $A$  is the span of

$$\{ (1, 0, \_, 0, \_), (0, 1, 1, \_, 0) \}.$$

- (c) The nullspace of  $A$  is the span of

$$\{ (\_, -1, 1, 0, 0), (0, 1, 0, \_, 0), (1, 0, 0, \_, \_) \}.$$

- (d) The nullspace of  $A^t$  is the span of

$$\{ (\_, \_, 1, 0), (-1/2, \_, 0, 1) \}.$$

- (6) Let  $A = \begin{bmatrix} 1 & -2 & -1 & 3 & 2 \\ -2 & 4 & 2 & -6 & -4 \\ 5 & -10 & -1 & 15 & 0 \\ 3 & -6 & 1 & 9 & -4 \\ 3 & -6 & -1 & 9 & 1 \\ 0 & 0 & 2 & 0 & -5 \end{bmatrix}$ .

- (a) The nullspace of  $A$  is the span of

$$\{ (2, 1, 0, 0, 0), (\_, 0, 0, 1, 0), (\_, 0, \_, 0, 1) \}.$$



(b) The row space of  $A$  is the span of

$$\{(1, -2, 0, \_, -1/2), (0, 0, 1, \_, \_) \}.$$

(c) The column space of  $A$  is the span of

$$\{(1, -2, 0, \_, 1/2, \_), (0, 0, 1, \_, \_, 1/2) \}.$$

(d) The left nullspace of  $A$  is the span of

$$\{(2, \_, 0, -1, 1, 0, 0), (\_, \_, 1, 0, 0, 0, 0), (-1/2, 0, \_, 0, 1, 0), (\_, \_, 0, -1/2, 0, 0, 1) \}$$

(7) Let  $A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 4 & 3 \\ 3 & 6 & 4 \end{bmatrix}$ .

(a) Fill in coordinates of the following vector  $\mathbf{x}$  so that it is perpendicular to the row space of  $A$ . Answer:  $\mathbf{x} = (10, \_, \_)$ .

(b) Fill in coordinates of the following vector  $\mathbf{y}$  so that it is perpendicular to the column space of  $A$ . Answer:  $\mathbf{y} = (3, \_, \_)$ .

(8) In this exercise we prove a slightly different version of the *fundamental theorem of linear algebra* than the one given in theorem 23.1.4. Here we work with real inner product spaces and the scope of the result is not restricted to finite dimensional spaces, but we must assume that the linear map with which we are dealing has a transpose.

**23.2.1. Theorem** (Fundamental theorem of linear algebra). *Suppose that  $V$  and  $W$  are arbitrary real inner product spaces and that the linear transformation  $T: V \rightarrow W$  has a transpose. Then*

$$\ker T = (\operatorname{ran} T^t)^\perp.$$

We prove the preceding theorem. For each step in the proof give the appropriate reason. Choose from the following list.

DK	Definition of “Kernel”
DO	Definition of “Orthogonal”
DOC	Definition of “Orthogonal Complement”
DR	Definition of “Range”
DT	Definition of “Transpose”
H	Hypothesis
PIP	Elementary Property of Inner Products

PROOF. We must show two things: (i)  $\ker T \subseteq (\operatorname{ran} T^t)^\perp$  and (ii)  $(\operatorname{ran} T^t)^\perp \subseteq \ker T$ .

To prove (i) we suppose that  $x \in \ker T$  and prove that  $x \in (\operatorname{ran} T^t)^\perp$ . Let  $v$  be a vector in  $\operatorname{ran} T^t$ . Then there exists a vector  $w$  in  $W$  such that  $v = T^t w$  (reason:  $\_$ ). We compute the inner product of  $x$  and  $v$ .

$$\begin{aligned} \langle x, v \rangle &= \langle x, T^t w \rangle \\ &= \langle Tx, w \rangle \quad (\text{reason: } \_) \\ &= \langle 0, w \rangle \quad (\text{reason: } \_ \text{ and } \_) \\ &= 0 \quad (\text{reason: } \_) \end{aligned}$$

From this we infer that  $x \perp v$  (reason:  $\_$ ) and consequently that  $x \in (\operatorname{ran} T^t)^\perp$  (reason:  $\_$ ).

To prove the converse we suppose that  $x \in (\text{ran } T^t)^\perp$  and show that  $x \in \ker T$ . We know that  $x \perp \text{ran } T^t$  (reason: \_\_\_\_\_ and \_\_\_\_\_). If  $w \in W$  then the vector  $T^t w$  belongs to  $\text{ran } T^t$  (reason: \_\_\_\_\_); so  $x \perp T^t w$  for all  $w \in W$ . Thus for all  $w \in W$

$$\begin{aligned} 0 &= \langle x, T^t w \rangle && (\text{reason: } \_\_\_\_\_\_) \\ &= \langle Tx, w \rangle && (\text{reason: } \_\_\_\_\_\_) \end{aligned}$$

It follows from this that  $Tx = 0$  (reason: \_\_\_\_\_). That is,  $x \in \ker T$  (reason: \_\_\_\_\_).  $\square$

- (9) The matrix  $\begin{bmatrix} x & y & z \\ y & 1 & x \end{bmatrix}$  has rank one if and only if the point  $(x, y, z)$  lies on the parametrized curve  $\mathbf{r}(t) = ( \_\_\_\_\_\_, t, \_\_\_\_\_\_ )$  in  $\mathbb{R}^3$ . *Hint.* Use proposition 23.1.3.
- (10) Let  $A$  be the  $3 \times 4$  matrix whose nullspace is the subspace of  $\mathbb{R}^4$  spanned by the vectors  $(1, 0, 1, 0)$  and  $(0, 1, 1, 0)$ . Then the vectors  $( \_\_\_\_\_\_, \_\_\_\_\_\_, \_\_\_\_\_\_, 0 )$  and  $( 0, \_\_\_\_\_\_, \_\_\_\_\_\_, \_\_\_\_\_\_ )$  form an orthonormal basis for the row space of  $A$ .
- (11) Let  $T: \mathbb{R}^3 \rightarrow \mathbb{R}^2$  be the linear transformation whose matrix representation is  $\begin{bmatrix} 1 & 0 & 2 \\ 1 & 1 & 4 \end{bmatrix}$  and let  $\mathbf{x} = (5, 4, -9)$ .
- (a) Find  $\mathbf{u} \in \ker T$  and  $\mathbf{v} \in \text{ran } T^t$  such that  $\mathbf{x} = \mathbf{u} + \mathbf{v}$ .  
 Answer:  $\mathbf{u} = ( \_\_\_\_\_\_, \_\_\_\_\_\_, \_\_\_\_\_\_ )$  and  $\mathbf{v} = ( \_\_\_\_\_\_, \_\_\_\_\_\_, \_\_\_\_\_\_ )$ .
- (b) Find  $\mathbf{y} \in \text{ran } T$  and  $\mathbf{z} \in \ker T^t$  such that  $T\mathbf{x} = \mathbf{y} + \mathbf{z}$ .  
 Answer:  $\mathbf{y} = ( \_\_\_\_\_\_, \_\_\_\_\_\_ )$  and  $\mathbf{z} = ( \_\_\_\_\_\_, \_\_\_\_\_\_ )$ .

### 23.3. Problems

- (1) Let  $T: \mathbb{R}^5 \rightarrow \mathbb{R}^4$  be a linear transformation whose matrix representation is

$$[T] = \begin{bmatrix} 1 & 2 & 0 & -5 & 3 \\ -2 & -4 & 3 & 1 & 0 \\ -1 & -2 & 3 & -4 & 3 \\ 1 & 2 & 3 & -14 & 9 \end{bmatrix}.$$

Gauss-Jordan reduction applied to  $[T]$  yields the matrix  $B = \begin{bmatrix} 1 & 2 & 0 & -5 & 3 \\ 0 & 0 & 1 & -3 & 2 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$  and applied to the transpose of  $[T]$  yields  $C = \begin{bmatrix} 1 & 0 & 1 & 3 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$ .

- From the matrices above we can read off the dimension of the range of  $T$  and write down a basis for it. Explain carefully.
  - From the matrices above we can read off the dimension of the range of the transpose of  $T$  and write down a basis for it. Explain carefully.
  - From the matrices above we can write down two equations which a vector  $(v, w, x, y, z)$  must satisfy to be in the kernel of  $T$ . Explain carefully. What are the equations? Also explain carefully how we obtain from these equations the dimension of the kernel of  $T$  and find a basis for it. Carry out the calculation you describe.
  - From the matrices above we can write down two equations which a vector  $(w, x, y, z)$  must satisfy to be in the kernel of the transpose of  $T$ . Explain carefully. What are the equations? Also explain carefully how we obtain from these equations the dimension of the kernel of  $T^t$  and find a basis for it. Carry out the calculation you describe.
- (2) Let  $T: \mathbb{R}^6 \rightarrow \mathbb{R}^3$  be a linear transformation whose matrix representation is

$$[T] = \begin{bmatrix} 1 & 2 & 0 & 2 & -1 & 1 \\ 3 & 6 & 1 & 1 & -2 & 1 \\ 5 & 10 & 1 & 5 & -4 & 3 \end{bmatrix},$$

Gauss-Jordan reduction applied to  $[T]$  yields the matrix  $B = \begin{bmatrix} 1 & 2 & 0 & 2 & -1 & 1 \\ 0 & 0 & 1 & -5 & 1 & -2 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$ , and applied to the transpose of  $[T]$  yields  $C = \begin{bmatrix} 1 & 0 & 2 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$ .

- From the matrices above we can read off the dimension of the range of  $T$  and write down a basis for it. Explain carefully.
- From the matrices above we can read off the dimension of the range of the transpose of  $T$  and write down a basis for it. Explain carefully.
- From the matrices above we can write down two equations which a vector  $(u, v, w, x, y, z)$  in  $\mathbb{R}^6$  must satisfy to be in the kernel of  $T$ . Explain carefully. What are the equations? Also explain carefully how we obtain from these equations the dimension of the kernel of  $T$  and find a basis for it. Carry out the calculation you describe.
- From the matrices above we can write down two equations which a vector  $(x, y, z)$  in  $\mathbb{R}^3$  must satisfy to be in the kernel of the transpose of  $T$ . Explain carefully. What are the equations? Also explain carefully how we obtain from these equations the

dimension of the kernel of  $T^t$  and find a basis for it. Carry out the calculation you describe.

- (3) Let  $T: \mathbb{R}^6 \rightarrow \mathbb{R}^5$  be the linear transformation whose matrix representation is

$$A = [T] = \begin{bmatrix} 1 & 2 & -1 & -2 & 3 & 0 \\ 0 & 0 & 0 & 1 & -1 & 2 \\ 2 & 4 & -2 & -4 & 7 & -4 \\ 0 & 0 & 0 & -1 & 1 & -2 \\ 3 & 6 & -3 & -6 & 7 & 8 \end{bmatrix}.$$

You may use the following fact: the reduced row echelon forms of the augmented matrix  $[A | \mathbf{b}]$  and of  $A^t$  are

$$B = \begin{bmatrix} 1 & 2 & -1 & 0 & 0 & 8 & 3b_1 + 2b_2 - b_3 \\ 0 & 0 & 0 & 1 & 0 & -2 & -2b_1 + b_2 + b_3 \\ 0 & 0 & 0 & 0 & 1 & -4 & -2b_1 + b_3 \\ 0 & 0 & 0 & 0 & 0 & 0 & b_2 + b_4 \\ 0 & 0 & 0 & 0 & 0 & 0 & -7b_1 + 2b_3 + b_5 \end{bmatrix} \quad \text{and} \quad C = \begin{bmatrix} 1 & 0 & 0 & 0 & 7 \\ 0 & 1 & 0 & -1 & 0 \\ 0 & 0 & 1 & 0 & -2 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}, \text{ respectively.}$$

Suppose that  $\mathbf{x} = \begin{bmatrix} t \\ u \\ v \\ w \\ y \\ z \end{bmatrix}.$

- What are the free variables of the system  $A\mathbf{x} = \mathbf{0}$  and which are the lead variables? How do you know?
- What is the rank of  $A$ ? Why?
- Write a general solution to the homogeneous equation  $A\mathbf{x} = \mathbf{0}$  as a linear combination of vectors in  $\mathbb{R}^6$  using the free variables as coefficients. Explain.
- Explain how to find the dimension of and a basis for the kernel of  $T$ . Do so.
- Explain how to find the dimension of and a basis for the range of  $T$ . Do so.

- What conditions must the vector  $\mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \\ b_5 \end{bmatrix}$  satisfy in order that the nonhomogeneous equation  $A\mathbf{x} = \mathbf{b}$  have solutions?

- Find, if possible, the general solution to the nonhomogeneous equation  $A\mathbf{x} = \begin{bmatrix} 1 \\ -3 \\ 2 \\ 3 \\ 3 \end{bmatrix}.$

(Write your answer as a general solution to the homogeneous equation plus a particular solution.)

- Explain how to find the dimension of and a basis for the range of  $T^t$ . Do so.
  - Explain how to find the dimension of and a basis for the kernel of  $T^t$ . Do so.
- (4) Prove the three corollaries to the *fundamental theorem of linear algebra* 23.1.4 for complex inner product spaces.

**23.4. Answers to Odd-Numbered Exercises**

- (1) (a)  $9 - 3i$ ,  $3 - i$   
(b)  $3$ ,  $-2$   
(c)  $-2$ ,  $-1$   
(d)  $-9 - 3i$ ,  $-3 - i$
- (3) (a)  $2$ ,  $1$   
(b)  $2x + y - z = 0$   
(c)  $2$   
(d)  $8$ ,  $-22$ ,  $2$ ,  $-8$   
(e)  $4$   
(f)  $-4$ ,  $6$
- (5) (a)  $\frac{1}{2}$ ,  $\frac{1}{2}$ ,  $\frac{3}{2}$   
(b)  $2$ ,  $-1$ ,  $-1$   
(c)  $-2$ ,  $1$ ,  $0$ ,  $1$   
(d)  $\frac{1}{2}$ ,  $-\frac{1}{2}$ ,  $-\frac{3}{2}$
- (7) (a)  $-5$ ,  $0$   
(b)  $3$ ,  $-3$
- (9)  $t^2$ ,  $t^3$
- (11) (a)  $6$ ,  $6$ ,  $-3$ ,  $-1$ ,  $-2$ ,  $-6$   
(b)  $-13$ ,  $-27$ ,  $0$ ,  $0$



## CHAPTER 24

# ORTHOGONAL PROJECTIONS

### 24.1. Background

**Topics:** orthogonal and unitary operators, orthogonal projections

**24.1.1. Definition.** A linear operator  $T: V \rightarrow V$  on a real inner product space is **ORTHOGONAL** if it is invertible and  $T^t = T^{-1}$ . A matrix is **ORTHOGONAL** if it is the representation of an orthogonal operator. An operator  $T$  on a complex inner product space  $V$  is **UNITARY** if it is invertible and  $T^* = T^{-1}$ . A matrix is **UNITARY** if it is the representation of a unitary operator.

The definitions for *orthogonal* and *unitary* matrices given above differs from the ones offered in 19.1.4. In problem 8 you are asked to show that in both cases the definitions are equivalent.

**24.1.2. Definition.** An operator  $T$  on an inner product space  $V$  is an **ISOMETRY** if it preserves the distance between vectors. Equivalently,  $T$  is an isometry if  $\|T\mathbf{x}\| = \|\mathbf{x}\|$  for every  $\mathbf{x} \in V$ .

**24.1.3. Definition.** Let  $V$  be an inner product space and suppose that it is the vector space direct sum of  $M$  and  $N$ . Then the projection  $E_{MN}: V \rightarrow V$  is an **ORTHOGONAL PROJECTION** if  $M \perp N$  (that is if  $V$  is the orthogonal direct sum of  $M$  and  $N$ ).

**24.1.4. Proposition.** *A projection  $E$  on a complex inner product space  $V$  is an orthogonal projection if and only if  $E$  is self-adjoint. On a real inner product space a projection is orthogonal if and only if it is symmetric.*

For a proof of (the real case of) this result see exercise 4.

**24.1.5. Definition.** We say that a linear operator  $T$  on an inner product space  $V$  is **POSITIVE** (and write  $T \geq 0$ ) if  $\langle T\mathbf{x}, \mathbf{x} \rangle \geq 0$  for all  $\mathbf{x} \in V$ . If  $S$  and  $T$  are two linear operators on  $V$ , we say that  $Q$  **DOMINATES** (or **MAJORIZES**)  $P$  if  $Q - P \geq 0$ . In this case we write  $P \leq Q$ .

## 24.2. Exercises

- (1) The matrix representation of the orthogonal projection operator taking  $\mathbb{R}^3$  onto the plane

$$x + y + z = 0 \text{ is } \begin{bmatrix} \phantom{0} \\ \phantom{0} \\ \phantom{0} \end{bmatrix}.$$

- (2) Find a vector  $\mathbf{u} = (u_1, u_2, u_3)$  in  $\mathbb{C}^3$  such that the matrix  $\begin{bmatrix} \frac{1}{m} & \frac{-1}{m} & u_1 \\ \frac{i}{m} & \frac{-i}{m} & u_2 \\ \frac{1-i}{m} & \frac{1-i}{m} & u_3 \end{bmatrix}$  is unitary.

Answer:  $\mathbf{u} = \frac{1}{\sqrt{n}}(2 + ai, 3 - bi, c + di)$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ ,  $c = \underline{\hspace{1cm}}$ ,  $d = \underline{\hspace{1cm}}$ ,  $m = \underline{\hspace{1cm}}$ , and  $n = \underline{\hspace{1cm}}$ .

- (3) The orthogonal projection of the vector  $(2, 0, -1, 3)$  on the plane spanned by  $(-1, 1, 0, 1)$  and  $(0, 1, 1, 1)$  in  $\mathbb{R}^4$  is  $\frac{1}{5}(1, a, b, a)$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ . The matrix which

implements this orthogonal projection is  $\frac{1}{5} \begin{bmatrix} c & -d & e & -d \\ -d & e & d & e \\ e & d & c & d \\ -d & e & d & e \end{bmatrix}$  where  $c = \underline{\hspace{1cm}}$ ,  $d = \underline{\hspace{1cm}}$ ,

and  $e = \underline{\hspace{1cm}}$ .

- (4) Let  $E$  be a projection operator on a real inner product space. Below we prove (the real case of) proposition 24.1.4: that  $E$  is an orthogonal projection if and only if  $E = E^t$ . Fill in the missing reasons and steps. Choose reasons from the following list.

(DK) Definition of “kernel”.

(DL) Definition of “linear”.

(DO) Definition of “orthogonal”.

(DOP) Definition of “orthogonal projection”.

(DT) Definition of “transpose”.

(GPa) Problem 5 in chapter 6.

(GPb) Problem 1 in chapter 11.

(GPc) Problem 5 in chapter 18.

(GPD) Problem 6 in chapter 22.

(H1) Hypothesis that  $M \perp N$ .

(H2) Hypothesis that  $E = E^t$ .

(PIP) Elementary property of Inner Products.

(Ti) Theorem 11.1.2, part (i).

(Tiii) Theorem 11.1.2, part (iii).

(Tiv) Theorem 11.1.2, part (iv).

(VA) Vector space arithmetic (consequences of vector space axioms, etc.)

Let  $E = E_{MN}$  be a projection operator on a real inner product space  $V = M \oplus N$ . Suppose first that  $E$  is an orthogonal projection. Then  $M \perp N$  (reason:  $\underline{\hspace{2cm}}$ ). If  $\mathbf{x}$  and  $\mathbf{y}$  are elements in  $V$ , then there exist unique vectors  $\mathbf{m}, \mathbf{p} \in M$  and  $\mathbf{n}, \mathbf{q} \in N$  such



that  $\mathbf{x} = \mathbf{m} + \mathbf{n}$  and  $\mathbf{y} = \mathbf{p} + \mathbf{q}$  (reason: \_\_\_\_\_) . Then

$$\begin{aligned}
 \langle E\mathbf{x}, \mathbf{y} \rangle &= \langle E(\mathbf{m} + \mathbf{n}), \mathbf{p} + \mathbf{q} \rangle \\
 &= \langle E\mathbf{m} + E\mathbf{n}, \mathbf{p} + \mathbf{q} \rangle \quad \text{reason: _____ and _____} \\
 &= \langle \mathbf{0} + E\mathbf{n}, \mathbf{p} + \mathbf{q} \rangle \quad \text{reason: _____ and _____} \\
 &= \langle E\mathbf{n}, \mathbf{p} + \mathbf{q} \rangle \quad \text{reason: _____} \\
 &= \langle \mathbf{n}, \mathbf{p} + \mathbf{q} \rangle \quad \text{reason: _____ and _____} \\
 &= \langle \mathbf{n}, \mathbf{p} \rangle + \langle \mathbf{n}, \mathbf{q} \rangle \quad \text{reason: _____} \\
 &= \mathbf{0} + \langle \mathbf{n}, \mathbf{q} \rangle \quad \text{reason: _____ and _____} \\
 &= \langle \mathbf{m}, \mathbf{q} \rangle + \langle \mathbf{n}, \mathbf{q} \rangle \quad \text{reason: _____ and _____} \\
 &= \langle \mathbf{x}, \mathbf{q} \rangle \quad \text{reason: _____} \\
 &= \langle \mathbf{x}, E\mathbf{q} \rangle \quad \text{reason: _____ and _____} \\
 &= \langle \mathbf{x}, \mathbf{0} + E\mathbf{q} \rangle \quad \text{reason: _____} \\
 &= \langle \mathbf{x}, E\mathbf{p} + E\mathbf{q} \rangle \quad \text{reason: _____ and _____} \\
 &= \langle \mathbf{x}, E(\mathbf{p} + \mathbf{q}) \rangle \quad \text{reason: _____ and _____} \\
 &= \langle \mathbf{x}, E^{tt}\mathbf{y} \rangle \quad \text{reason: _____} \\
 &= \langle E^t\mathbf{x}, \mathbf{y} \rangle \quad \text{reason: _____} .
 \end{aligned}$$

From this we conclude that  $E = E^t$  (reason: \_\_\_\_\_) .

Conversely, suppose that  $E = E^t$ . To show that  $M \perp N$  it is enough to show that  $\mathbf{m} \perp \mathbf{n}$  for arbitrary elements  $\mathbf{m} \in M$  and  $\mathbf{n} \in N$ .

$$\begin{aligned}
 \langle \mathbf{n}, \mathbf{m} \rangle &= \langle E\mathbf{n}, \mathbf{m} \rangle \quad \text{reason: _____ and _____} \\
 &= \langle \mathbf{n}, E^t\mathbf{m} \rangle \quad \text{reason: _____} \\
 &= \langle \mathbf{n}, E\mathbf{m} \rangle \quad \text{reason: _____} \\
 &= \langle \mathbf{n}, \mathbf{0} \rangle \quad \text{reason: _____ and _____} \\
 &= \mathbf{0} \quad \text{reason: _____}
 \end{aligned}$$

Thus  $\mathbf{m} \perp \mathbf{n}$  (reason: \_\_\_\_\_) .

**Note:** Of course, the complex inner product space version of the preceding result says that if  $E$  is a projection operator on a complex inner product space, then  $E$  is an orthogonal projection if and only if it is self-adjoint.

- (5) Let  $P$  be the orthogonal projection of  $\mathbb{R}^3$  onto the subspace spanned by the vectors  $(1, 0, 1)$

and  $(1, 1, -1)$ . Then  $[P] = \frac{1}{6} \begin{bmatrix} a & b & c \\ b & b & -b \\ c & -b & a \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ , and  $c = \underline{\hspace{1cm}}$ .

- (6) Find the image of the vector  $\mathbf{b} = (1, 2, 7)$  under the orthogonal projection of  $\mathbb{R}^3$  onto the

column space of the matrix  $A = \begin{bmatrix} 1 & 1 \\ 2 & -1 \\ -2 & 4 \end{bmatrix}$ .

Answer: ( \_\_\_\_\_ , \_\_\_\_\_ , \_\_\_\_\_ ).

- (7) Let  $\mathbf{u} = (3, -1, 1, 4, 2)$  and  $\mathbf{v} = (1, 2, -1, 0, 1)$ . Then the orthogonal projection of  $\mathbf{u}$  onto  $\mathbf{v}$  is

( \_\_\_\_\_ , \_\_\_\_\_ , \_\_\_\_\_ , \_\_\_\_\_ , \_\_\_\_\_ ).

- (8) Let  $\mathbf{u} = (8, \sqrt{3}, \sqrt{7}, -1, 1)$  and  $\mathbf{v} = (1, -1, 0, 2, \sqrt{3})$ . Then the orthogonal projection of  $\mathbf{u}$  onto  $\mathbf{v}$  is  $\frac{a}{b}\mathbf{v}$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .
- (9) Let  $\mathbf{u} = (5, 4, 3, \frac{1}{2})$  and  $\mathbf{v} = (1, 2, 0, -2)$ . Then the orthogonal projection of  $\mathbf{u}$  onto  $\mathbf{v}$  is  $\frac{a}{b}\mathbf{v}$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .
- (10) Find the point  $\mathbf{q}$  in  $\mathbb{R}^3$  on the ray connecting the origin to the point  $(2, 4, 8)$  which is closest to the point  $(1, 1, 1)$ .  
 Answer:  $\mathbf{q} = \frac{1}{3} (\underline{\hspace{1cm}}, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})$ .
- (11) Let  $\mathbf{e}^1 = (\frac{2}{3}, \frac{2}{3}, -\frac{1}{3})$  and  $\mathbf{e}^2 = (-\frac{1}{3}, \frac{2}{3}, \frac{2}{3})$  be vectors in  $\mathbb{R}^3$ . Notice that  $\{\mathbf{e}^1, \mathbf{e}^2\}$  is an orthonormal set.
- (a) Find a vector  $\mathbf{e}^3$  whose first coordinate is positive such that  $B = \{\mathbf{e}^1, \mathbf{e}^2, \mathbf{e}^3\}$  is an orthonormal basis for  $\mathbb{R}^3$ . Answer:  $\frac{1}{3} (\underline{\hspace{1cm}}, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})$ .
- (b) Suppose that  $\mathbf{x}$  is a vector in  $\mathbb{R}^3$  whose Fourier coefficients with respect to the basis  $B$  are:  $\langle \mathbf{x}, \mathbf{e}^1 \rangle = -2$ ;  $\langle \mathbf{x}, \mathbf{e}^2 \rangle = -1$ ; and  $\langle \mathbf{x}, \mathbf{e}^3 \rangle = 3$ . Then  $\mathbf{x} = (\underline{\hspace{1cm}}, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})$ .
- (c) Let  $\mathbf{y}$  be a vector in  $\mathbb{R}^3$  whose Fourier coefficients with respect to  $B$  are

$$\langle \mathbf{y}, \mathbf{e}_1 \rangle = \sqrt{8 - \sqrt{37}};$$

$$\langle \mathbf{y}, \mathbf{e}_2 \rangle = \sqrt{5 - \sqrt{13}}; \quad \text{and}$$

$$\langle \mathbf{y}, \mathbf{e}_3 \rangle = \sqrt{3 + \sqrt{13} + \sqrt{37}}.$$

Then the length of the vector  $\mathbf{y}$  is  $\underline{\hspace{1cm}}$ .

- (d) The orthogonal projection of the vector  $\mathbf{b} = (0, 3, 0)$  onto the plane spanned by  $\mathbf{e}_1$  and  $\mathbf{e}_2$  is  $\frac{2}{3} (\underline{\hspace{1cm}}, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})$ .
- (e) The orthogonal projection of the vector  $\mathbf{b} = (0, 3, 0)$  onto the line spanned by  $\mathbf{e}_3$  is  $\frac{1}{3} (\underline{\hspace{1cm}}, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})$ .
- (f) What vector do you get when you add the results of the projections you found in parts (d) and (e)? Answer:  $(\underline{\hspace{1cm}}, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})$ .

## 24.3. Problems

- (1) Prove that an operator  $T: V \rightarrow V$  on a finite dimensional real inner product space  $V$  is orthogonal if and only if it is an isometry. Similarly, on a finite dimensional complex inner product space an operator is unitary if and only if it is an isometry.
- (2) Prove that an operator  $T: V \rightarrow V$  on a finite dimensional real inner product space  $V$  is orthogonal if and only if  $T^t T = I$ . What is the corresponding necessary and sufficient condition on a finite dimensional complex inner product space for an operator to be unitary?
- (3) Show that if an operator  $U$  on a complex inner product space is both Hermitian and unitary, then  $\sigma(U) \subseteq \{-1, 1\}$ .
- (4) Let  $P$  and  $Q$  be orthogonal projections on a real inner product space. Show that their sum  $P + Q$  is an orthogonal projection if and only if  $PQ = QP = \mathbf{0}$ . *Hint.* Use proposition 24.1.4.
- (5) Explain in detail how to find the matrix which represents the orthogonal projection of  $\mathbb{R}^3$  onto the plane  $x + y - 2z = 0$ . Carry out the computation you describe.
- (6) Let  $P$  and  $Q$  be orthogonal projection operators on a real inner product space  $V$ .
  - (a) Show that the operator  $PQ$  is an orthogonal projection if and only if  $P$  commutes with  $Q$ .
  - (b) Show that if  $P$  commutes with  $Q$ , then

$$\text{ran}(PQ) = \text{ran } P \cap \text{ran } Q.$$

*Hint.* To show that  $\text{ran } P \cap \text{ran } Q \subseteq \text{ran}(PQ)$  start with a vector  $\mathbf{y}$  in  $\text{ran } P \cap \text{ran } Q$  and examine  $PQ\mathbf{y}$ .

- (7) Let  $P$  and  $Q$  be orthogonal projections on an inner product space  $V$ . Prove that the following are equivalent:
  - (a)  $P \leq Q$ ;
  - (b)  $\|P\mathbf{x}\| \leq \|Q\mathbf{x}\|$  for all  $\mathbf{x} \in V$ ;
  - (c)  $\text{ran } P \subseteq \text{ran } Q$ ;
  - (d)  $QP = P$ ; and
  - (e)  $PQ = P$ .

*Hint.* First show that (d) and (e) are equivalent. Then show that (a)  $\Rightarrow$  (b)  $\Rightarrow$  (c)  $\Rightarrow$  (d)  $\Rightarrow$  (a). To prove that (b)  $\Rightarrow$  (c) take an arbitrary element  $x$  in the range of  $P$ ; show that  $\|Q\mathbf{x}\| = \|\mathbf{x}\|$  and that consequently  $\|(I - Q)\mathbf{x}\| = 0$ . To prove that (d)  $\Rightarrow$  (a) show that  $(I - P)Q$  is an orthogonal projection; then consider  $\|(I - P)Q\|^2$ .

- (8) In 19.1.4 and 24.1.1 the definitions for *unitary* matrices differ. Show that they are, in fact, equivalent. Argue that the same is true for the definitions given for *orthogonal* matrices.

**24.4. Answers to Odd-Numbered Exercises**

$$(1) \frac{1}{3} \begin{bmatrix} 2 & -1 & -1 \\ -1 & 2 & -1 \\ -1 & -1 & 2 \end{bmatrix}$$

$$(3) 3, 4, 3, 1, 2$$

$$(5) 5, 2, 1$$

$$(7) \frac{2}{7}, \frac{4}{7}, -\frac{2}{7}, 0, \frac{2}{7}$$

$$(9) 4, 3$$

$$(11) \text{ (a) } 2, -1, 2$$

$$\text{ (b) } 1, -3, 2$$

$$\text{ (c) } 4$$

$$\text{ (d) } 1, 4, 1$$

$$\text{ (e) } -2, 1, -2$$

$$\text{ (f) } 0, 3, 0$$

## CHAPTER 25

# LEAST SQUARES APPROXIMATION

### 25.1. Background

**Topics:** least squares approximation.

## 25.2. Exercises

(1) Let  $A = \begin{bmatrix} 1 & 1 \\ 2 & -1 \\ -2 & 4 \end{bmatrix}$ .

- (a) Find an orthonormal basis  $\{\mathbf{e}^1, \mathbf{e}^2, \mathbf{e}^3\}$  for  $\mathbb{R}^3$  such that  $\{\mathbf{e}^1, \mathbf{e}^2\}$  spans the column space of  $A$ .

$$\mathbf{e}^1 = \frac{1}{n}(a, b, -b)$$

$$\mathbf{e}^2 = \frac{1}{n}(b, a, b)$$

$$\mathbf{e}^3 = \frac{1}{n}(b, -b, -a)$$

where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ , and  $n = \underline{\hspace{1cm}}$ .

- (b) To which of the four fundamental subspaces of  $A$  does  $\mathbf{e}^3$  belong?

Answer:  $\mathbf{e}^3$  belongs to the    of  $A$ .

- (c) What is the least squares solution to  $A\mathbf{x} = \mathbf{b}$  when  $\mathbf{b} = (1, 2, 7)$ ?

Answer:  $\hat{\mathbf{x}} = (\underline{\hspace{1cm}}, \underline{\hspace{1cm}})$ .

- (2) Find the best least squares fit by a straight line to the following data:  $x = 1$  when  $t = -1$ ;  $x = 3$  when  $t = 0$ ;  $x = 2$  when  $t = 1$ ; and  $x = 3$  when  $t = 2$ .

Answer:  $x = \underline{\hspace{1cm}} + \underline{\hspace{1cm}} t$ .

- (3) At times  $t = -2, -1, 0, 1$ , and  $2$  the data  $y = 4, 2, -1, 0$ , and  $0$ , respectively, are observed. Find the best line to fit this data. Answer:  $y = Ct + D$  where  $C = \underline{\hspace{1cm}}$  and  $D = \underline{\hspace{1cm}}$ .

- (4) The best (least squares) line fit to the data:  $y = 2$  at  $t = -1$ ;  $y = 0$  at  $t = 0$ ;  $y = -3$  at  $t = 1$ ;  $y = -5$  at  $t = 2$  is  $y = -\frac{a}{10} - \frac{b}{5}t$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .

- (5) Consider the following data:  $y = 20$  when  $t = -2$ ;  $y = 6$  when  $t = -1$ ;  $y = 2$  when  $t = 0$ ;  $y = 8$  when  $t = 1$ ;  $y = 24$  when  $t = 2$ . Find the parabola which best fits the data in the least squares sense.

Answer:  $y = C + Dt + Et^2$  where  $C = \underline{\hspace{1cm}}$ ,  $D = \underline{\hspace{1cm}}$ , and  $E = \underline{\hspace{1cm}}$ .

- (6) Consider the following data:  $y = 2$  when  $t = -1$ ;  $y = 0$  when  $t = 0$ ;  $y = -3$  when  $t = 1$ ;  $y = -5$  when  $t = 2$ . Find the parabola which best fits the data in the least squares sense.

Answer:  $y = C + Dt + Et^2$  where  $C = \underline{\hspace{1cm}}$ ,  $D = \underline{\hspace{1cm}}$ , and  $E = \underline{\hspace{1cm}}$ .

- (7) Find the plane  $50z = a + bu + cv$  which is the best least squares fit to the following data:  $z = 3$  when  $u = 1$  and  $v = 1$ ;  $z = 6$  when  $u = 0$  and  $v = 3$ ;  $z = 5$  when  $u = 2$  and  $v = 1$ ;  $z = 0$  when  $u = 0$  and  $v = 0$ .

Answer:  $a = \underline{\hspace{1cm}}$ ;  $b = \underline{\hspace{1cm}}$ ;  $c = \underline{\hspace{1cm}}$ .

- (8) Consider the following data:  $y = 4$  at  $t = -1$ ;  $y = 5$  at  $t = 0$ ;  $y = 9$  at  $t = 1$ .

- (a) Then the best (least squares) line which fits the data is  $y = c + dt$  where  $c = \underline{\hspace{1cm}}$  and  $d = \underline{\hspace{1cm}}$ .

- (b) The orthogonal projection of  $\mathbf{b} = (4, 5, 9)$  onto the column space of  $A = \begin{bmatrix} 1 & -1 \\ 1 & 0 \\ 1 & 1 \end{bmatrix}$  is  $(\underline{\hspace{1cm}}, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})$ .

- (9) The best least squares solution to the following (inconsistent) system of equations  $\begin{cases} u & = 1 \\ v & = 1 \\ u + v & = 0 \end{cases}$  is  $u = \underline{\hspace{1cm}}$  and  $v = \underline{\hspace{1cm}}$ .

**25.3. Problems**

- (1) Explain in detail how to use matrix methods to find the best (least squares) solution to the following (inconsistent) system of equations  $\begin{cases} u &= 1 \\ v &= 1 \\ u + v &= 0 \end{cases}$ . Carry out the computation you describe.
- (2) The following data  $y$  are observed at times  $t$ :  $y = 4$  when  $t = -2$ ;  $y = 3$  when  $t = -1$ ;  $y = 1$  when  $t = 0$ ; and  $y = 0$  when  $t = 2$ .
- (a) Explain how to use matrix methods to find the best (least squares) straight line approximation to the data. Carry out the computation you describe.
- (b) Find the orthogonal projection of  $\mathbf{y} = (4, 3, 1, 0)$  on the column space of the matrix
- $$A = \begin{bmatrix} 1 & -2 \\ 1 & -1 \\ 1 & 0 \\ 1 & 2 \end{bmatrix}.$$
- (c) Explain carefully what your answer in (b) has to do with part (a).
- (d) At what time does the largest error occur? That is, when does the observed data differ most from the values your line predicts?

**25.4. Answers to Odd-Numbered Exercises**

- (1) (a) 1, 2, 3
  - (b) left nullspace
  - (c) 1, 2
- (3) -1, 1
- (5) 2, 1, 5
- (7) -6, 73, 101
- (9)  $\frac{1}{3}, \frac{1}{3}$



**Part 7**

**SPECTRAL THEORY OF INNER  
PRODUCT SPACES**



## SPECTRAL THEOREM FOR REAL INNER PRODUCT SPACES

### 26.1. Background

**Topics:** the spectral theorem for finite dimensional real inner product spaces.

**26.1.1. Definition.** An operator  $T$  on a finite dimensional real inner product space with an orthonormal basis is ORTHOGONALLY DIAGONALIZABLE if there exists an orthogonal matrix which diagonalizes  $T$ .

The following theorem (together with its analog for complex spaces) is the fundamental structure theorem for inner product spaces. It says that any symmetric operator on a finite dimensional real inner product space can be written as a linear combination of orthogonal projections. The coefficients are the eigenvalues of the operator and the ranges of the orthogonal projections are the eigenspaces of the operator.

**26.1.2. Theorem** (Spectral Theorem for Finite Dimensional Real Inner Product Spaces). *Let  $T$  be a symmetric operator on a finite dimensional real inner product space  $V$ , and  $\lambda_1, \dots, \lambda_k$  be the (distinct) eigenvalues of  $T$ . For each  $j$  let  $M_j$  be the eigenspace associated with  $\lambda_j$  and  $E_j$  be the projection of  $V$  onto  $M_j$  along  $M_1 + \dots + M_{j-1} + M_{j+1} + \dots + M_k$ . Then  $T$  is orthogonally diagonalizable, the eigenspaces of  $T$  are mutually orthogonal, each  $E_j$  is an orthogonal projection, and the following hold:*

- (i)  $T = \lambda_1 E_1 + \dots + \lambda_k E_k$ ,
- (ii)  $I = E_1 + \dots + E_k$ , and
- (iii)  $E_i E_j = 0$  when  $i \neq j$ .

## 26.2. Exercises

- (1) Let
- $T$
- be the operator on
- $\mathbb{R}^3$
- whose matrix representation is

$$\begin{bmatrix} \frac{1}{3} & -\frac{2}{3} & -\frac{2}{3} \\ -\frac{2}{3} & \frac{5}{6} & -\frac{7}{6} \\ -\frac{2}{3} & -\frac{7}{6} & \frac{5}{6} \end{bmatrix}.$$

- (a) Find the characteristic polynomial and minimal polynomial for
- $T$
- .

Answer:  $c_T(\lambda) = \underline{\hspace{2cm}}$ . $m_T(\lambda) = \underline{\hspace{2cm}}$ .

- (b) The eigenspace  $M_1$  associated with the smallest eigenvalue  $\lambda_1$  is the span of  $(1, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})$ .
- (c) The eigenspace  $M_2$  associated with the middle eigenvalue  $\lambda_2$  is the span of  $(\underline{\hspace{1cm}}, \underline{\hspace{1cm}}, -1)$ .
- (d) The eigenspace  $M_3$  associated with the largest eigenvalue  $\lambda_3$  is the span of  $(\underline{\hspace{1cm}}, 1, \underline{\hspace{1cm}})$ .
- (e) Find the (matrix representations of the) orthogonal projections  $E_1$ ,  $E_2$ , and  $E_3$  onto the eigenspaces  $M_1$ ,  $M_2$ , and  $M_3$ , respectively.

Answer:  $E_1 = \frac{1}{m} \begin{bmatrix} a & a & a \\ a & a & a \\ a & a & a \end{bmatrix}$ ;  $E_2 = \frac{1}{n} \begin{bmatrix} b & -c & -c \\ -c & a & a \\ -c & a & a \end{bmatrix}$ ;  $E_3 = \frac{1}{2} \begin{bmatrix} d & d & d \\ d & a & -a \\ d & -a & a \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ ,  $c = \underline{\hspace{1cm}}$ ,  $d = \underline{\hspace{1cm}}$ ,  $m = \underline{\hspace{1cm}}$ , and  $n = \underline{\hspace{1cm}}$ .

- (f) Write
- $T$
- as a linear combination of the projections found in (e).

Answer:  $[T] = \underline{\hspace{1cm}} E_1 + \underline{\hspace{1cm}} E_2 + \underline{\hspace{1cm}} E_3$ .

- (g) Find an orthogonal matrix
- $Q$
- (that is, a matrix such that
- $Q^t = Q^{-1}$
- ) which diagonalizes
- $T$
- . What is the associated diagonal form
- $\Lambda$
- of
- $T$
- ?

Answer:  $Q = \begin{bmatrix} \frac{a}{\sqrt{b}} & \frac{c}{\sqrt{bc}} & 0 \\ \frac{a}{\sqrt{b}} & -\frac{a}{\sqrt{bc}} & \frac{a}{\sqrt{c}} \\ \frac{a}{\sqrt{b}} & -\frac{a}{\sqrt{bc}} & -\frac{a}{\sqrt{c}} \end{bmatrix}$  and  $\Lambda = \begin{bmatrix} \lambda & 0 & 0 \\ 0 & \mu & 0 \\ 0 & 0 & \nu \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ ,  $c = \underline{\hspace{1cm}}$ ,  $\lambda = \underline{\hspace{1cm}}$ ,  $\mu = \underline{\hspace{1cm}}$ , and  $\nu = \underline{\hspace{1cm}}$ .

- (2) Let
- $T$
- be the operator on
- $\mathbb{R}^3$
- whose matrix representation is

$$\begin{bmatrix} 2 & 2 & 1 \\ 2 & 2 & -1 \\ 1 & -1 & -1 \end{bmatrix}.$$

- (a) The eigenspace  $M_1$  associated with the smallest eigenvalue  $\lambda_1$  is the span of  $(1, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})$ .
- (b) The eigenspace  $M_2$  associated with the middle eigenvalue  $\lambda_2$  is the span of  $(1, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})$ .
- (c) The eigenspace  $M_3$  associated with the largest eigenvalue  $\lambda_3$  is the span of  $(1, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})$ .
- (d) Find the (matrix representations of the) orthogonal projections  $E_1$ ,  $E_2$ , and  $E_3$  onto the eigenspaces  $M_1$ ,  $M_2$ , and  $M_3$ , respectively.

Answer:  $E_1 = \frac{1}{mn} \begin{bmatrix} a & -a & -b \\ -a & a & b \\ -b & b & c \end{bmatrix}$ ;  $E_2 = \frac{1}{m} \begin{bmatrix} a & -a & a \\ -a & a & -a \\ a & -a & a \end{bmatrix}$ ;  $E_3 = \frac{1}{n} \begin{bmatrix} a & a & d \\ a & a & d \\ d & d & d \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ ,  $c = \underline{\hspace{1cm}}$ ,  $d = \underline{\hspace{1cm}}$ ,  $m = \underline{\hspace{1cm}}$ , and  $n = \underline{\hspace{1cm}}$ .

- (e) Write
- $T$
- as a linear combination of the projections found in (d).

Answer:  $[T] = \underline{\hspace{1cm}} E_1 + \underline{\hspace{1cm}} E_2 + \underline{\hspace{1cm}} E_3$ .

- (f) Find an orthogonal matrix
- $Q$
- (that is, a matrix such that
- $Q^t = Q^{-1}$
- ) which diagonalizes
- $T$
- . What is the associated diagonal form
- $\Lambda$
- of
- $T$
- ?

Answer:  $Q = \begin{bmatrix} \frac{a}{\sqrt{bc}} & \frac{a}{\sqrt{b}} & \frac{a}{\sqrt{c}} \\ -\frac{a}{\sqrt{bc}} & -\frac{a}{\sqrt{b}} & \frac{a}{\sqrt{c}} \\ -\frac{c}{\sqrt{bc}} & \frac{a}{\sqrt{b}} & 0 \end{bmatrix}$  and  $\Lambda = \begin{bmatrix} \lambda & 0 & 0 \\ 0 & \mu & 0 \\ 0 & 0 & \nu \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ ,  
 $c = \underline{\hspace{1cm}}$ ,  $\lambda = \underline{\hspace{1cm}}$ ,  $\mu = \underline{\hspace{1cm}}$ , and  $\nu = \underline{\hspace{1cm}}$ .

**26.3. Problem**

(1) Let  $A = \begin{bmatrix} 1 & -4 & 2 \\ -4 & 1 & -2 \\ 2 & -2 & -2 \end{bmatrix}$ .

- (a) Does  $A$  satisfy the hypotheses of the *spectral theorem* 26.1.2 for symmetric operators on a finite dimensional real inner product space? Explain.
- (b) Explain how to find an orthogonal matrix which diagonalizes the matrix  $A$ . Carry out the computation you describe.
- (c) Explain in careful detail how to write the matrix  $A$  in part (b) as a linear combination of orthogonal projections. Carry out the computations you describe.

**26.4. Answers to the Odd-Numbered Exercise**

- (1) (a)  $\lambda^3 - 2\lambda^2 - \lambda + 2$  (or  $(\lambda + 1)(\lambda - 1)(\lambda - 2)$ );  $\lambda^3 - 2\lambda^2 - \lambda + 2$  (or  $(\lambda + 1)(\lambda - 1)(\lambda - 2)$ )  
(b) 1, 1  
(c) 2, -1  
(d) 0, -1  
(e) 1, 4, 2, 0, 3, 6  
(f) -1, 1, 2  
(g) 1, 3, 2, -1, 1, 2





## SPECTRAL THEOREM FOR COMPLEX INNER PRODUCT SPACES

### 27.1. Background

**Topics:** the spectral theorem for finite dimensional complex inner product spaces.

**27.1.1. Definition.** An operator  $T$  on a finite dimensional real inner product space with an orthonormal basis is UNITARILY DIAGONALIZABLE if there exists an orthogonal matrix which diagonalizes  $T$ .

**27.1.2. Theorem** (Spectral Theorem for Finite Dimensional Complex Inner Product Spaces). *Let  $T$  be a normal operator on a finite dimensional complex inner product space  $V$ , and  $\lambda_1, \dots, \lambda_k$  be the (distinct) eigenvalues of  $T$ . For each  $j$  let  $M_j$  be the eigenspace associated with  $\lambda_j$  and  $E_j$  be the projection of  $V$  onto  $M_j$  along  $M_1 + \dots + M_{j-1} + M_{j+1} + \dots + M_k$ . Then  $T$  is unitarily diagonalizable, the eigenspaces of  $T$  are mutually orthogonal, each  $E_j$  is an orthogonal projection, and the following hold:*

- (i)  $T = \lambda_1 E_1 + \dots + \lambda_k E_k$ ,
- (ii)  $I = E_1 + \dots + E_k$ , and
- (iii)  $E_i E_j = 0$  when  $i \neq j$ .

**27.1.3. Theorem.** *Let  $T$  be an operator on a finite dimensional complex inner product space  $V$ . Then the following are equivalent:*

- (1)  $T$  is normal;
- (2)  $T$  is unitarily diagonalizable; and
- (3)  $V$  has an orthonormal basis consisting of eigenvectors of  $T$ .

## 27.2. Exercises

(1) Let  $A = \begin{bmatrix} 2 & 1+i \\ 1-i & 3 \end{bmatrix}$ .

- (a) Use the *spectral theorem* 27.1.2 to write  $A$  as a linear combination of orthogonal projections.

Answer:  $A = \alpha E_1 + \beta E_2$  where  $\alpha = \underline{\hspace{1cm}}$ ,  $\beta = \underline{\hspace{1cm}}$ ,  $E_1 = \frac{1}{3} \begin{bmatrix} 2 & -1-i \\ 1-i & 3 \end{bmatrix}$ ,  
and  $E_2 = \frac{1}{3} \begin{bmatrix} 1 & 1+i \\ 1-i & 3 \end{bmatrix}$ .

- (b) Find a square root of  $A$ .

Answer:  $\sqrt{A} = \frac{1}{3} \begin{bmatrix} 4 & 1+i \\ 1-i & 3 \end{bmatrix}$ .

(2) Let  $T$  be the operator on  $\mathbb{C}^2$  whose matrix representation is  $\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$ .

- (a) The eigenspace  $V_1$  associated with the eigenvalue  $-i$  is the span of  $(1, \underline{\hspace{1cm}})$ .  
(b) The eigenspace  $V_2$  associated with the eigenvalue  $i$  is the span of  $(1, \underline{\hspace{1cm}})$ .  
(c) The (matrix representations of the) orthogonal projections  $E_1$  and  $E_2$  onto the eigenspaces

$V_1$  and  $V_2$ , respectively, are  $E_1 = \begin{bmatrix} a & b \\ -b & a \end{bmatrix}$ ; and  $E_2 = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .

- (d) Write  $T$  as a linear combination of the projections found in (c).

Answer:  $[T] = \underline{\hspace{1cm}} E_1 + \underline{\hspace{1cm}} E_2$ .

- (e) A unitary matrix  $U$  which diagonalizes  $[T]$  is  $\begin{bmatrix} a & a \\ -b & b \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .  
The associated diagonal form  $\Lambda = U^*[T]U$  of  $[T]$  is  $\begin{bmatrix} \underline{\hspace{1cm}} & 0 \\ 0 & \underline{\hspace{1cm}} \end{bmatrix}$ .

(3) Let  $N = \frac{1}{3} \begin{bmatrix} 4+2i & 1-i & 1-i \\ 1-i & 4+2i & 1-i \\ 1-i & 1-i & 4+2i \end{bmatrix}$ .

- (a) The matrix  $N$  is normal because  $NN^* = N^*N = \begin{bmatrix} a & b & b \\ b & a & b \\ b & b & a \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .

- (b) Thus according to the *spectral theorem* 27.1.2  $N$  can be written as a linear combination of orthogonal projections. Written in this form  $N = \lambda_1 E_1 + \lambda_2 E_2$  where  $\lambda_1 = \underline{\hspace{1cm}}$ ,  
 $\lambda_2 = \underline{\hspace{1cm}}$ ,  $E_1 = \begin{bmatrix} a & a & a \\ a & a & a \\ a & a & a \end{bmatrix}$ , and  $E_2 = \begin{bmatrix} b & -a & -a \\ -a & b & -a \\ -a & -a & b \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$  and  $b = \underline{\hspace{1cm}}$ .

- (c) A unitary matrix  $U$  which diagonalizes  $N$  is  $\begin{bmatrix} a & -b & -c \\ a & b & -c \\ a & d & 2c \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$ ,  
 $b = \underline{\hspace{1cm}}$ ,  $c = \underline{\hspace{1cm}}$ , and  $d = \underline{\hspace{1cm}}$ . The associated diagonal form  $\Lambda = U^*NU$   
of  $N$  is  $\begin{bmatrix} \underline{\hspace{1cm}} & 0 & 0 \\ 0 & \underline{\hspace{1cm}} & 0 \\ 0 & 0 & \underline{\hspace{1cm}} \end{bmatrix}$ .

(4) Let  $T$  be an operator whose matrix representation is  $\begin{bmatrix} 1 & 2 \\ -1 & -1 \end{bmatrix}$ .

(a) Regarded as an operator on  $\mathbb{R}^2$  is  $T$  triangulable? \_\_\_\_\_. As an operator on  $\mathbb{R}^2$  is it diagonalizable? \_\_\_\_\_.

(b) Show that  $T$  regarded as an operator on  $\mathbb{C}^2$  is diagonalizable by finding numbers  $c$  and  $d$  such that the matrix  $S = \begin{bmatrix} -2 & -2 \\ c & d \end{bmatrix}$  is invertible and  $S^{-1}TS$  is diagonal.

Answer:  $c = \underline{\hspace{1cm}}$  and  $d = \underline{\hspace{1cm}}$ .

(c) Show that despite being diagonalizable (as an operator on  $\mathbb{C}^2$ )  $T$  is not normal.

Answer:  $TT^* = \begin{bmatrix} \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} \end{bmatrix} \neq \begin{bmatrix} \phantom{0} & \phantom{0} \\ \phantom{0} & \phantom{0} \end{bmatrix} = T^*T.$

(d) Explain briefly why the result of part (c) does not contradict Theorem 27.1.3.

(5) Let  $T$  be the operator on  $\mathbb{C}^3$  whose matrix representation is  $\frac{1}{6} \begin{bmatrix} 8-i & 5-2i & 2+4i \\ -5+2i & 8-i & -4+2i \\ -2-4i & -4+2i & 14+2i \end{bmatrix}$ .

(a) Find the characteristic polynomial and minimal polynomial for  $T$ .

Answer:  $c_T(\lambda) = \underline{\hspace{4cm}}$ .

$m_T(\lambda) = \underline{\hspace{4cm}}$ .

(b) The eigenspace  $M_1$  associated with the real eigenvalue  $\lambda_1$  is the span of  $(1, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})$ .

(c) The eigenspace  $M_2$  associated with the complex eigenvalue  $\lambda_2$  with negative imaginary part is the span of  $(1, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})$ .

(d) The eigenspace  $M_3$  associated with the remaining eigenvalue  $\lambda_3$  is the span of  $(1, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})$ .

(e) Find the (matrix representations of the) orthogonal projections  $E_1$ ,  $E_2$ , and  $E_3$  onto the eigenspaces  $M_1$ ,  $M_2$ , and  $M_3$ , respectively.

Answer:  $E_1 = \frac{1}{m} \begin{bmatrix} 1 & -b & bc \\ b & a & -c \\ -bc & -c & d \end{bmatrix}$ ;  $E_2 = \frac{1}{n} \begin{bmatrix} 1 & b & e \\ -b & a & e \\ e & e & e \end{bmatrix}$ ;  $E_3 = \frac{1}{p} \begin{bmatrix} 1 & -b & -b \\ b & a & a \\ b & a & a \end{bmatrix}$   
 where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ ,  $c = \underline{\hspace{1cm}}$ ,  $d = \underline{\hspace{1cm}}$ ,  $e = \underline{\hspace{1cm}}$ ,  $m = \underline{\hspace{1cm}}$ ,  $n = \underline{\hspace{1cm}}$ , and  $p = \underline{\hspace{1cm}}$ .

(f) Write  $T$  as a linear combination of the projections found in (e).

Answer:  $[T] = \underline{\hspace{1cm}} E_1 + \underline{\hspace{1cm}} E_2 + \underline{\hspace{1cm}} E_3.$

(g) Find a unitary matrix  $U$  which diagonalizes  $T$ . What is the associated diagonal form  $\Lambda$  of  $T$ ?

Answer:  $U = \begin{bmatrix} \frac{a}{\sqrt{bc}} & \frac{a}{\sqrt{b}} & \frac{a}{\sqrt{c}} \\ \frac{d}{\sqrt{bc}} & -\frac{d}{\sqrt{b}} & \frac{d}{\sqrt{c}} \\ -\frac{bd}{\sqrt{bc}} & \frac{e}{\sqrt{b}} & \frac{d}{\sqrt{c}} \end{bmatrix}$  and  $\Lambda = \begin{bmatrix} \lambda & 0 & 0 \\ 0 & \mu & 0 \\ 0 & 0 & \nu \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ ,  
 $c = \underline{\hspace{1cm}}$ ,  $d = \underline{\hspace{1cm}}$ ,  $e = \underline{\hspace{1cm}}$ ,  $\lambda = \underline{\hspace{1cm}}$ ,  $\mu = \underline{\hspace{1cm}}$ , and  $\nu = \underline{\hspace{1cm}}$ .

(h) The operator  $T$  is normal because  $TT^* = T^*T = \frac{1}{6} \begin{bmatrix} a & -bc & 2bc \\ bc & a & -2b \\ -2bc & -2b & d \end{bmatrix}$  where  
 $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ ,  $c = \underline{\hspace{1cm}}$ , and  $d = \underline{\hspace{1cm}}$ .

(6) Let  $T$  be the operator on  $\mathbb{C}^3$  whose matrix representation is  $\frac{1}{3} \begin{bmatrix} 5+2i & 2-i & 2-i \\ 2-i & 5-i & 2+2i \\ 2-i & 2+2i & 5-i \end{bmatrix}$ .

(a) Find the characteristic polynomial and minimal polynomial for  $T$ .

Answer:  $c_T(\lambda) = \underline{\hspace{4cm}}$ .

$m_T(\lambda) = \underline{\hspace{4cm}}$ .

(b) The eigenspace  $M_1$  associated with the real eigenvalue  $\lambda_1$  is the span of  $(1, \underline{\hspace{1cm}}, \underline{\hspace{1cm}})$ .

(c) The eigenspace  $M_2$  associated with the complex eigenvalue  $\lambda_2$  with negative imaginary part is the span of  $(\underline{\hspace{1cm}}, \underline{\hspace{1cm}}, -1)$ .

(d) The eigenspace  $M_3$  associated with the remaining eigenvalue  $\lambda_3$  is the span of  $(\underline{\hspace{1cm}}, -1, \underline{\hspace{1cm}})$ .

(e) Find the (matrix representations of the) orthogonal projections  $E_1$ ,  $E_2$ , and  $E_3$  onto the eigenspaces  $M_1$ ,  $M_2$ , and  $M_3$ , respectively.

Answer:  $E_1 = \frac{1}{m} \begin{bmatrix} a & a & a \\ a & a & a \\ a & a & a \end{bmatrix}$ ;  $E_2 = \frac{1}{n} \begin{bmatrix} b & b & b \\ b & c & -c \\ b & -c & c \end{bmatrix}$ ;  $E_3 = \frac{1}{6} \begin{bmatrix} d & -e & -e \\ -e & a & a \\ -e & a & a \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ ,  $c = \underline{\hspace{1cm}}$ ,  $d = \underline{\hspace{1cm}}$ ,  $e = \underline{\hspace{1cm}}$ ,  $m = \underline{\hspace{1cm}}$ , and  $n = \underline{\hspace{1cm}}$ .

(f) Write  $T$  as a linear combination of the projections found in (e).

Answer:  $[T] = \underline{\hspace{1cm}} E_1 + \underline{\hspace{1cm}} E_2 + \underline{\hspace{1cm}} E_3$ .

(g) Find an orthogonal matrix  $Q$  (that is, a matrix such that  $Q^t = Q^{-1}$ ) which diagonalizes  $T$ . What is the associated diagonal form  $\Lambda$  of  $T$ ?

Answer:  $Q = \begin{bmatrix} \frac{a}{\sqrt{b}} & 0 & \frac{c}{\sqrt{bc}} \\ \frac{a}{\sqrt{b}} & \frac{a}{\sqrt{c}} & -\frac{a}{\sqrt{bc}} \\ \frac{a}{\sqrt{b}} & -\frac{a}{\sqrt{c}} & -\frac{a}{\sqrt{bc}} \end{bmatrix}$  and  $\Lambda = \begin{bmatrix} \lambda & 0 & 0 \\ 0 & \mu & 0 \\ 0 & 0 & \nu \end{bmatrix}$  where  $a = \underline{\hspace{1cm}}$ ,  $b = \underline{\hspace{1cm}}$ ,  $c = \underline{\hspace{1cm}}$ ,  $\lambda = \underline{\hspace{1cm}}$ ,  $\mu = \underline{\hspace{1cm}}$ , and  $\nu = \underline{\hspace{1cm}}$ .

**27.3. Problems**

- (1) Let  $N$  be a normal operator on a finite dimensional complex inner product space  $V$ . Show that  $\|N\mathbf{x}\| = \|N^*\mathbf{x}\|$  for all  $\mathbf{x} \in V$ .
- (2) Let  $N$  be a normal operator on a complex finite dimensional inner product space  $V$ . Show that if  $\lambda_1, \dots, \lambda_k$  are the eigenvalues of  $N$ , then  $\overline{\lambda_1}, \dots, \overline{\lambda_k}$  are the eigenvalues of  $N^*$ .
- (3) Let  $T$  be as in exercise 4. Show by direct computation that there is no invertible  $2 \times 2$  matrix  $S = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$  of real numbers such that  $S^{-1}TS$  is upper triangular.

**27.4. Answers to Odd-Numbered Exercises**

- (1) (a)  $1, 4, -1 + i, 1, 1 - i, 2$   
 (b)  $1 - i, 5$
- (3) (a)  $\frac{8}{3}, \frac{2}{3}$   
 (b)  $2, 1 + i, \frac{1}{3}, \frac{2}{3}$   
 (c)  $\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{6}}, 0, \begin{bmatrix} 2 & 0 & 0 \\ 0 & 1 + i & 0 \\ 0 & 0 & 1 + i \end{bmatrix}$
- (5) (a)  $\lambda^3 - 5\lambda^2 + 8\lambda - 6$  (or  $(\lambda^2 - 2\lambda + 2)(\lambda - 3)$ );  $\lambda^3 - 5\lambda^2 + 8\lambda - 6$  (or  $(\lambda^2 - 2\lambda + 2)(\lambda - 3)$ )  
 (b)  $i, -2i$   
 (c)  $-i, 0$   
 (d)  $i, i$   
 (e)  $1, i, 2, 4, 0, 6, 2, 3$   
 (f)  $3, 1 - i, 1 + i$   
 (g)  $1, 2, 3, i, 0, 3, 1 - i, 1 + i$   
 (h)  $19, 7, i, 40$

## Bibliography

1. Howard Anton and Chris Rorres, *Elementary Linear Algebra: Applications Version*, eighth ed., John Wiley and Sons, New York, 2000. vii, 51, 85
2. Robert A. Beezer, *A First Course in Linear Algebra*, 2004, <http://linear.ups.edu/download/fcla-electric-2.00.pdf>. vii
3. Przemyslaw Bogacki, *Linear Algebra Toolkit*, 2005, <http://www.math.odu.edu/~bogacki/lat>. vii
4. William C. Brown, *A second Course in Linear Algebra*, John Wiley, New York, 1988. vii
5. Charles W. Curtis, *Linear Algebra: An Introductory Approach*, Springer, New York, 1984. vii
6. Paul R. Halmos, *Finite-Dimensional Vector Spaces*, D. Van Nostrand, Princeton, 1958. vii
7. Jim Hefferon, *Linear Algebra*, 2006, <http://joshua.smcvt.edu/linearalgebra>. vii, 51
8. Kenneth Hoffman and Ray Kunze, *Linear Algebra*, second ed., Prentice Hall, Englewood Cliffs, N.J., 1971. vii, 93, 149
9. Steven Roman, *Advanced Linear Algebra*, second ed., Springer-Verlag, New York, 2005. vii, 127
10. Gilbert Strang, *Linear Algebra and its Applications*, second ed., Academic Press, New York, 1980. vii, 133
11. Eric W. Weisstein, *MathWorld*, A Wolfram Web Resource, <http://mathworld.wolfram.com>. vii
12. Wikipedia, *Wikipedia, The Free Encyclopedia*, <http://en.wikipedia.org>. vii





# Index

- $\angle(\mathbf{x}, \mathbf{y})$  (angle between  $\mathbf{x}$  and  $\mathbf{y}$ ), 25, 122
- $\mathbf{x} \cdot \mathbf{y}$  (inner product of  $\mathbf{x}$  and  $\mathbf{y}$ ), 121
- $M \perp N$  (two sets are orthogonal), 122
- $\mathbf{x} \perp \mathbf{y}$  (two vectors are orthogonal), 122
- $M \leq V$  ( $M$  is a subspace of  $V$ ), 39
- $\langle \mathbf{x}, \mathbf{y} \rangle$  (inner product), 121
- $[T]$  (matrix representation of  $T$ ), 71
- $[A, B]$  (commutator of  $A$  and  $B$ ), 9
- $TS$  (notation for composition of linear maps), 61
- $\langle \mathbf{x}, \mathbf{y} \rangle$  (inner product of  $\mathbf{x}$  and  $\mathbf{y}$ ), 25, 121
- $\mathbf{x} \cdot \mathbf{y}$  (inner product of  $\mathbf{x}$  and  $\mathbf{y}$ ), 25, 121
- $M \oplus N$  (direct sum of  $M$  and  $N$ ), 39
- $\|x\|$  (norm of a vector  $x$ ), 121
- $A^t$  (transpose of  $A$ ), 9
- $M^\perp$  (the orthogonal complement of  $M$ ), 122
- action
  - of a matrix, 15
- additive
  - inverses, 33
- adjoint
  - of a linear map, 145
- angle, 25, 122
- annihilating polynomial, 87
- associative, 33
- basis, 55
  - orthonormal, 127
  - standard, 55
- bijective, 61
- bounded function, 39
- $\mathbb{C}^n$ 
  - as an inner product space, 121
- $\mathcal{C}([a, b])$ 
  - as an inner product space, 122
- Cauchy-Schwarz inequality, 25, 122
- Cayley-Hamilton theorem, 87
- characteristic polynomial, 83
- cofactor, 16
- column
  - rank, 149
- column index, 9
- column space, 149
- combination
  - linear, 47
- commutative, 33
  - diagram, 68
- commutator, 9
- commute, 9, 68
- complement
  - orthogonal, 122
- complementary subspaces, 39
- conjugate
  - Hermitian, 145
  - linear, 121
  - transpose, 145
    - of a linear map, 145
    - of a matrix, 145
- continuous
  - uniformly, 122
- continuously differentiable, 64
- cross product, 25
- definite
  - negative, 133
  - positive, 133
- dependence
  - linear, 47
- determinant, 16
- $\det A$  (determinant of  $A$ ), 16
- diagonal matrix, 87
- diagonalizable, 87
  - conditions to be, 88
  - orthogonally, 171
  - part of an operator, 105
  - unitarily, 177
- diagonalizing matrix, 87
- diagram, 68
  - commutative, 68
- differentiable
  - continuously, 64
- differential equations, 97
- dimension, 55
- $\dim V$  (dimension of  $V$ ), 55
- direct sum, 39
  - orthogonal, 127
- dominate, 159
- $E_{MN}$  (projection along  $M$  onto  $N$ ), 77
- eigenspace, 83
  - generalized, 105
- eigenvalue, 83
- eigenvector, 83
- electrical networks
  - problem on, 51

- equivalent
    - row, 15
  - even function, 41
  - exact sequence, 67
  - expansion
    - Laplace, 16
  - factorization
    - $QR$ -, 127
  - finite dimension, 55
  - Fourier series, 127
  - function
    - even, 41
    - odd, 41
    - square integrable, 123
    - uniformly continuous, 122
  - functional
    - linear, 61
  - fundamental theorem of linear algebra, 149, 153
  - generalized eigenspace, 105
  - Hermitian
    - conjugate, 145
    - operator, 145
  - Hessian matrix, 139
  - $I_V$  (identity operator on  $V$ ), 62
  - identity
    - additive, 33
  - identity map, 62
  - indefinite, 133
  - independence
    - linear, 47
  - index
    - column, 9
    - row, 9
  - inequality
    - (Cauchy-)Schwarz, 122
  - infinite dimensional, 55
  - injective, 61
  - inner product, 25, 121
    - space
      - $\mathbb{C}^n$  as a, 121
      - $\mathbb{R}^n$  as a, 121
      - $\mathcal{C}([a, b])$  as a, 122
      - $l_2$  as a, 122
  - integrable, 39
  - inverse
    - additive, 33
    - left, 62
    - right, 62
  - invertible, 62
  - isometry, 159
  - isomorphic, 62
  - isomorphism, 62
  - $\ker T$  (the kernel of  $T$ ), 61
  - kernel, 61
  - $\mathcal{L}(V)$ 
    - linear operators on  $V$ , 61
  - $\mathcal{L}(V, W)$ 
    - linear maps between vector spaces, 61
  - $l_2$ 
    - as an inner product space, 122
    - square summable sequences, 122
  - Laplace expansion, 16
  - leading principal submatrix, 133
  - left
    - inverse, 12, 62
    - nullspace, 149
  - length, 25
  - Lie bracket, 9
  - linear, 61
    - combination, 47
    - trivial, 47
  - conjugate, 121
  - dependence, 47
  - functional, 61
  - independence, 47
  - map, 61
    - adjoint of a, 145
    - conjugate transpose of a, 145
    - Hermitian conjugate of a, 145
    - transpose of a, 145
  - operator, 61
  - sesqui-, 121
  - transformation, 61
- majorize, 159
- map
  - linear, 61
- Markov matrix, 97
- matrix
  - conjugate transpose of a, 145
  - diagonal, 87
  - diagonalizing, 87
  - Hessian, 139
  - Markov, 97
  - nilpotent, 105
  - nonsingular, 33
  - normal, 145
  - notation for, 9
  - orthogonal, 127, 159
  - representation of an linear map, 71
  - second derivative, 139
  - standard, 71
  - symmetric, 9
  - transpose of a, 9
  - unitary, 127, 159
  - upper triangular, 9
- maximal
  - linearly independent set, 57
  - orthonormal set, 127
- minimal polynomial, 87
  - existence and uniqueness of, 91
- minimal spanning set, 57
- minor, 16
- monic polynomial, 87
- negative
  - definite, 133

- networks
  - problem on, 51
- nilpotent, 88, 105
  - part of an operator, 105
- nonsingular, 33
- norm, 25, 121
- normal
  - matrix, 145
  - operator, 145
- nullity, 61
- nullspace, 61, 149
  - left, 149
- odd function, 41
- one-to-one, 61
  - correspondence, 61
- onto, 61
- operator
  - adjoint of an, 145
  - diagonalizable part of an, 105
  - Hermitian, 145
  - linear, 61
  - nilpotent, 105
  - nilpotent part of an, 105
  - normal, 145
  - orthogonal, 159
  - positive, 159
  - self-adjoint, 145
  - symmetric, 145
  - transpose of an, 145
  - unitary, 159
- orthogonal, 122
  - complement, 122
  - diagonalization, 171
  - direct sum, 127
  - matrix, 127, 159
  - operator, 159
  - projection, 159
- orthonormal, 127
  - basis, 127
- perpendicular, 122
- pivoting, 3
- point spectrum, 83
- polarization identity, 125
- polynomial
  - annihilating, 87
  - characteristic, 83
  - minimal, 87
    - existence and uniqueness of, 91
  - monic, 87
- positive
  - definite, 133
  - operator, 159
- principal submatrix, 133
- product
  - cross, 25
  - inner, 25, 121
- projection
  - along one subspace onto another, 77
  - orthogonal, 159
- QR*-factorization, 127
- $\mathbb{R}^n$ 
  - as an inner product space, 121
- $\text{ran } T$  (the range of  $T$ ), 61
- range, 61
- rank
  - column, 149
  - of a linear map, 61
  - row, 149
- representation
  - matrix, 71
- Riemann integrable function, 39
- right
  - inverse, 12, 62
- row
  - equivalent matrices, 15
  - operations, 3
  - rank, 149
- row index, 9
- row space, 149
- scaling, 3
- Schwarz inequality, 25, 122
- second derivative
  - matrix, 139
  - test, 139
- self-adjoint
  - operator, 145
- sequence
  - exact, 67
- sesquilinear, 121
- short exact sequence, 67
- $\sigma(T)$  (spectrum of  $T$ ), 83
- similar, 87
- space
  - vector, 33
- span, 47
- span  $A$  (the span of the set  $A$ ), 47
- spectral mapping theorem, 83
- spectral theorem
  - for finite dimensional complex inner product spaces, 177
  - for finite dimensional real inner product spaces, 171
  - for finite dimensional vector spaces, 93
- spectrum, 83
- square integrable function, 123
- square summable, 122
- standard
  - basis for  $\mathbb{R}^n$ , 55
  - matrix for a linear map, 71
- submatrix
  - leading principal, 133
- subspace, 39
- subspaces
  - complementary, 39
- sum
  - direct, 39
- summable
  - square, 122
- surjective, 61

- swapping, 3
- symmetric
  - matrix, 9
  - operator, 145
- system
  - of differential equations, 97
- trace, 9
- $\text{tr } A$  (trace of  $A$ ), 9
- transformation
  - linear, 61
- transpose
  - conjugate, 145
  - of a matrix, 9
  - of an operator, 145
- triangulable, 88
  - conditions to be, 88
- triangular
  - upper, 9
- trivial
  - linear combination, 47
- uniformly continuous, 122
- unitary
  - diagonalization, 177
  - matrix, 127, 159
  - operator, 159
- upper
  - triangular, 9
- upper triangular, 88
- vector
  - space, 33
    - complex, 33
    - real, 33