hovid

COSAMINE

Glucosamine Sulphate
250 mg & 500 mg

- Lubricate Joints
- Adjunctive Therapy in Osteoarthritis

as POTASSIUM salt
GLUCOSAMINE
MARINE SOURCE

COSAMINE 250 mg

Product description/产品简述/Deskripsi keluaran:
Reddish orange opaque / white opaque capsule with 'CM250' printed on one end and 'hovid' printed on the other end of the capsule.
'CM250' 和 'hovid' 字眼各印在橙红色和白色不透明胶囊上。
Kapsul jingga kemerahan opaque / putih opaque dengan 'CM250' tercetak pada satu hujung dan 'hovid' pada hujung yang satu lagi.

Recommended dosage/推荐用量/Dos cadangan:
For mild to moderate osteoarthritis symptoms:
Oral, 2 capsules 2 times daily for 6 weeks, taken 15- 30 minutes before meal.
For severe osteoarthritis symptoms:
Oral, 2 capsules 3 times daily for 8 weeks, taken 15- 30 minutes before meal.
Maintenance therapy should be followed for 3-4 months by administration of 2 capsules twice daily.

轻微骨关节炎:
口服：每天两次，每次两粒，使用六星期，在用餐的15至30分钟前食用。
严重骨关节炎:
口服：每天三次，每次两粒，使用八星期，在用餐的15至30分钟前食用。
治疗须继续三至四个月，每天两次，每次两粒。

Bagi simptom osteoarthritis ringan sehingga sederhana:
Oral, 2 kapsul dua kali sehari selama 6 minggu, diambil 15-30 minit sebelum makan.
Bagi simptom osteoarthritis serius:
Oral, 2 kapsul tiga kali sehari selama 8 minggu, diambil 15-30 minit sebelum makan.
Terapi jagaan perlu diteruskan sehingga 3-4 bulan dengan pengambilan 2 kapsul dua kali sehari.

COSAMINE 500 mg

Product description/产品简述/Deskripsi keluaran:
Reddish orange opaque / white opaque capsule with 'CM500' printed on one end and 'hovid' printed on the other end of the capsule.
'CM500' 和 'hovid' 字眼各印在橙红色和白色不透明胶囊上。
Kapsul jingga kemerahan opaque / putih opaque dengan 'CM500' tercetak pada satu hujung dan 'hovid' pada hujung yang satu lagi.

Recommended dosage/推荐用量/Dos cadangan:
For mild to moderate osteoarthritis symptoms:
Oral, 1 capsule 2 times daily for 6 weeks, taken 15- 30 minutes before meal.
For severe osteoarthritis symptoms:
Oral, 1 capsule 3 times daily for 8 weeks, taken 15- 30 minutes before meal.
Maintenance therapy should be followed for 3-4 months by administration of 1 capsule twice daily.

轻微骨关节炎:
口服：每天两次，每次一粒，使用六星期，在用餐的15至30分钟前食用。
严重骨关节炎:
口服：每天三次，每次一粒，使用八星期，在用餐的15至30分钟前食用。
治疗须继续三至四个月，每天两次，每次一粒。

Bagi simptom osteoarthritis ringan sehingga sederhana:
Oral, 1 kapsul dua kali sehari selama 6 minggu, diambil 15-30 minit sebelum makan.
Bagi simptom osteoarthritis serius:
Oral, 1 kapsul tiga kali sehari selama 8 minggu, diambil 15-30 minit sebelum makan.
Terapi jagaan perlu diteruskan sehingga 3-4 bulan dengan pengambilan 1 kapsul dua kali sehari.

hovid

COSAMINE

Glucosamine Sulphate
250 mg & 500 mg

- Lubricate Joints
- Adjunctive Therapy in Osteoarthritis



as POTASSIUM salt
GLUCOSAMINE
MARINE SOURCE

COSAMINE

COSAMINE 250 mg
Each capsule contains :
Crystalline Glucosamine Potassium Sulphate 331.60 mg equivalent to Glucosamine Sulphate 250 mg.
每胶囊含量：
葡萄糖胺硫酸晶体331.60mg相等于葡萄糖胺硫酸250 mg.

Pack size/包装/Saiz pembungkusan:
Glass/Plastic bottle of 100's.
每瓶（玻璃瓶/塑料瓶）100粒。
100 biji setiap botol kaca/plastik.

Storage condition/储存方法/
Cara penyimpanan:
Store below 30°C. Protect from moisture.
存放在摄氏三十度以下。避免潮湿。
Simpan di bawah suhu 30°C.
Lindungi dari lembapan.

Shelf life/有效期/Tarikh luput:
3 years from the date of manufacture.
三年自生产期计算。
3 tahun dari tarikh pengilangan.

COSAMINE 500 mg
Each capsule contains :
Crystalline Glucosamine Potassium Sulphate 663.20 mg equivalent to Glucosamine Sulphate 500 mg.
每胶囊含量：
葡萄糖胺硫酸晶体663.20mg相等于葡萄糖胺硫酸500 mg.

Pack size/包装/Saiz pembungkusan:
Glass/Plastic bottle of 60's.
每瓶（玻璃瓶/塑料瓶）60粒。
60 biji setiap botol kaca/plastik.

Storage condition/储存方法/
Cara penyimpanan:
Store below 30°C. Protect from moisture.
存放在摄氏三十度以下。避免潮湿。
Simpan di bawah suhu 30°C.
Lindungi dari lembapan.

Shelf life/有效期/Tarikh luput:
3 years from the date of manufacture.
三年自生产期计算。
3 tahun dari tarikh pengilangan.

Indications/指示/Indikasi:
Adjunctive therapy in osteoarthritis.
关节炎的附加治疗。
Terapi tambahan osteoarthritis.

Contraindications/禁忌/Kontraindikasi:
Contraindicated in patients sensitive to glucosamine sulphate and shellfish.
对葡萄糖胺硫酸和水生贝壳类动物过敏的病人不能使用。
Tidak dibenarkan bagi pesakit yang sensitif terhadap glucosamine sulphate dan kerang-kerangan.

Precautions/提防/Perhatian:
Cosamine is not meant to replace NSAIDs in cases of intense pain, as this is a causal therapy. It is advisable to take an anti-inflammatory drug in the initial treatment of osteoarthritis together with glucosamine sulphate.
Cosamine is to be taken under medical supervision in pregnant and lactating mothers, and in geriatric patients.
Therapeutic effect can only be seen after approximately 1 week from the beginning.

在骨关节炎病人极其疼痛的情况下Cosamine不能用来取代NSAIDs当作止痛药，比较好的疗法是让初期骨关节炎病人一起使用抗发炎药和葡萄糖胺硫酸。
服用约一星期后方见效。

Pengambilan Cosamine bukan untuk menggantikan ubatan NSAIDs akibat kes sakit yang tersangat. Adalah dinasihatkan supaya mengambil ubatan anti-inflamasi pada rawatan awal osteoarthritis bersama dengan glucosamine sulphate.
Pengambilan Cosamine perlu di bawah nasihat doktor bagi pesakit mengandung dan menyusu dan pesakit yang berumur.
Kesan terapeutik hanya akan berkesan selepas penggunaan ubat selama seminggu.

DERIVED FROM SEAFOOD.
DIPEROLEHI DARI SUMBER LAUT.
取源于海产。

Incompatibility/不适应性/Ketakserasian:
None known.
未有记录。
Tidak diketahui.

Symptoms of overdose and treatment/
过量的症状及处理法/
Simptom akibat dos lebihan dan rawatan:
None known.
未有记录。
Tidak diketahui.

Drug interactions/药性交互作用/
Tindakbalas ubat:
None known.
没有记录。
Tidak diketahui.

7 8 2

COSAMINE

Pharmacodynamics/药物效应动力学/ Farmakodinamik:

Glucosamine possesses antiarthrosic-arthrotrophic activities. It stimulates the anabolic metabolism of osteo-cartilagineous tissues via stimulation of the biosynthesis of the mucopolysaccharides (which are the essential components of the cartilage ground substances) and of the bone mesenchymal tissues.
Glucosamine also acts to improve the viscosity of synovial fluid by increasing synovial fluid production, thereby providing lubricant activity.

葡萄糖胺有抗关节炎及保护关节的作用。通过刺激体内细胞合成粘多糖(是软骨组织基质的构成成分)及骨间质组织。
它能刺激骨及软骨组织的同化新陈代谢。
葡萄糖胺也能改善滑液的生产及粘性，从而供应润滑的需要。

Glucosamine mengandungi aktiviti antiarthrosik-arthrotrofik. Ia merangsangkan metabolisme anabolik tisu rawan otot melalui rangsangan biosintesis mukopolisakarida (komponen penting bagi rawan) dan tisu mesenkima tulang.

Pharmacokinetics/药物代谢动力学/ Farmakokinetik:

Absorption
Glucosamine sulphate is well absorbed (with a proportion close to 90%) following oral administration.

吸收
葡萄糖胺硫酸在口服后吸收良好(比例接近90%)。

Penyerapan
Pengambilan glucosamine sulphate secara oral mudah diserap (berhampiran dengan kadar 90%).

Protein binding
It is known that glucosamine and its metabolites bind to α - and β -globulins.

蛋白结合
葡萄糖胺及其代谢产物会与 α -和 β -球蛋白结合。

Ikatan protein
Adalah diketahui bahawa glucosamine dan metabolitnya mengikat kepada α - dan β -globulin.

Distribution
Mainly distributed to bone tissues and articular cartilages.

分布
主要分布于骨组织和关节软骨

Pengedaran
Diedarkan terutamanya kepada tisu tulang dan rawan artikular.

Side effects/副作用/Kesan sampingan:

Cardiovascular
Peripheral oedema, tachycardia were reported in a few patients following larger clinical trials investigating oral administration in osteoarthritis. Causal relationship has not been established.

心血管
研究口服葡萄糖胺用予治疗骨关节炎的大型临床试验中，少数患者出现心率加快、末梢水肿。因果关系尚未确立。

Kardiovaskular
Edema peripheral, takikardia telah dilaporkan oleh beberapa pesakit berikutan ujian klinikal secara pengambilan oral yang dijalankan ke atas osteoarthritis. Perhubungan penyebab belum ditubuhkan.

Central nervous system
Drowsiness, headache, insomnia have been observed rarely during therapy (less than 1%).

中枢神经系统
治疗期间极少出现嗜睡、头痛、失眠等不良反应（少于1%）。

Sistem Saraf Pusat
Mengantuk, sakit kepala, insomnia jarang diperhatikan semasa terapi (kurang daripada 1%).

Gastrointestinal
Nausea, vomiting, diarrhoea, dyspepsia or epigastric pain, constipation, heartburn and anorexia have been described rarely during oral therapy with glucosamine.

胃肠道
葡萄糖胺口服治疗期间极少出现恶心、呕吐、腹泻、消化不良或上腹部疼痛、便秘、胃灼热和厌食症等不良反应。

Gastrousus
Loya, muntah, cirit-birit, dispepsia atau sakit epigastrik, sembelit, pedih ulu hati dan anoreksia jarang diperhatikan semasa terapi oral dengan Glucosamine.

Skin
Skin reactions such as erythema and pruritus have been reported with therapeutic administration of glucosamine.

皮肤
使用葡萄糖胺治疗可导致皮肤过敏，包括红斑和瘙痒。

Kulit
Reaksi kulit seperti eritema dan pruritus telah dilaporkan dengan pemberian terapeutik Glucosamine.

VICOS19-var1 (MY)

COSAMINE

What is Osteoarthritis?

Osteoarthritis (OA) is a degenerative joint disease that most commonly affects middle-aged and older people. This occurs equally in both sexes, however, after the age of 55, the incidence is higher in women.

OA is marked by the degeneration of joint cartilage and enlargement of bone at the joints, often accompanied by pain and stiffness. As the disease progresses, the cartilage wears out completely, causing bone to rub against bone.

While there may be occasions where no symptoms may surface, people with OA are most commonly presented with gradual or subtle onset of deep aching joint pain (worsens after exercise, relieved by rest), joint swelling, morning stiffness and grating of joint motion.

Glucosamine Sulphate

Glucosamine is a simple molecule that the body makes from sugar and uses as a major building block of cartilage. It is an important component of the mucopolysaccharides, which provides structure to bone, cartilage and other body tissues.

Several studies have shown that glucosamine stimulates the production of cartilage, and improve joint functions in those with osteoarthritis.

Studies conducted showed favourable results of glucosamine relieving OA pain over a period of time. Unlike non-steroidal anti-inflammatory drugs, glucosamine does not cause gastrointestinal side effects, e.g. heartburn, stomach pain, gastritis and nausea.

References/参考资料：
1) Thienm, etal,: J. Rheumatol 2001 Jun; 28 (6): 1347 - 55
2) Lopes Vaz A. Curr Med Res Opin 1982; 8 (3): 145 - 9
3) Qiu Gx,: etal,: Arzneimittelforschung 1998 May; 48 (5): 469 - 74
4) J.Y Reginster et al. Lancet 2001, vol 357, 251 - 256

什么是骨关节炎?

骨关节炎是一种关节退化的疾病，多发生在中年及老年人身上。男女得病的机会是相同的，但是妇女年龄超过55岁得病率比男人高。

骨关节炎的特征是关节骨退化及关节骨头增大，通常带有关节疼痛及关节僵硬。随着关节发炎恶化，关节软骨完全耗损，使骨头与骨头直接摩檫。

虽然有时候骨关节炎病人的症状无马上出现，不过大多数的病人会有慢性的疼痛(运动后会恶化，休息可减轻关节疼痛)，关节肿胀， 早晨起床关节僵硬及身体行动缓慢的现象逐渐发生。

葡萄糖胺硫酸盐

葡萄糖胺是在我们体内糖合成的一种简单的分子，也是形成软骨最主要构造的成分。葡萄糖胺是粘多糖的主要成分，而粘多糖组成骨头，软骨及其他身体细胞的结构。

几项试验及研究已证明葡萄糖胺除了刺激软骨的形成也帮助减轻骨关节炎病人的关节疼痛及改善关节功能。

试验也证明葡萄糖胺能减轻骨关节炎患者的关节疼痛。除此之外葡萄糖胺也没有非类固醇消炎药副作用的现象（胃灼热，胃痛，胃炎和反胃）。

Product Registration Holder: **HOVID Bhd.** (58476-A)
121, Jalan Tunku Abdul Rahman,
30010 Ipoh, Malaysia.

Manufactured by: **HOVID Bhd.** (58476-A)
Lot 56442, 7$^{1}/_{2}$ Miles, Jalan Ipoh/Chemor,
31200 Chemor, Malaysia

Revision date: May 2023