

FOR VETERINARY USE ONLY

INECTIL 20%

INJECTABLE SOLUTION



COMPOSITION

Tylosin base	20 g
Benzyl alcohol	4 g
Diluent to	100 ml

PRODUCT DESCRIPTION

Yellowish liquid solution.

PHARMACODYNAMIC PROPERTIES

Tylosin is a mixture of macrolide antibiotics produced by the growth of certain strains of *Streptomyces fradiae*. It consists basically of tylosin A, which must not be lower than 80%, but tylosins B (desmycosin), C (macrocin) and D (relomycin) may also be present. These 4 factors represent at least 95% of the real content. Chemically it is characterized by a lactone ring. At usual doses it exerts a bacteriostatic effect, but at high concentrations it may be bactericidal. Tylosin penetrates the bacteria by passive diffusion and blocks the bacterial protein synthesis by binding to 50s ribosomal subunit.

Tylosin is effective against:

- *Mycoplasma* spp.
- *Spirochaetes*.
- *Chlamydias* spp.
- *Gram (+) germs*: *Staphylococcus*.

Streptococcus. *Erysipella*. *Corynebacterium*. *Clostridium* spp.

- *Gram (-) germs*: *Fusobacterium*.

Pasteurella spp. *Bordetella*.

The resistant micro-organisms may present cross-resistance with other macrolides such as erythromycin, lincomycin, spiramycin and spectinomycin.

PHARMACOKINETIC PROPERTIES

When administered parenterally, Tylosin reaches maximum blood concentrations within 1-2 hours. Tylosin combines with bovine plasma proteins at 40%. Plasma levels are lower than those in the tissues. It is metabolized in the liver and excreted unchanged in the urine and bile.

ENVIRONMENTAL PROPERTIES

Not applicable.

SPECIES FOR WHICH INTENDED AND INDICATIONS

Cattle: Treatment of bovine respiratory complex (shipping fever, pneumonia) usually associated with *Pasteurella multocida* and *Arcanobacterium pyogenes*; foot rot (necrotic pododermatitis) and calf diptheria caused by *Fusobacterium necrophorum* and metritis caused by *Arcanobacterium pyogenes*.

Pigs: Treatment of swine arthritis caused by *Mycoplasma hyosynoviae*; swine erysipelas caused by *Erysipelothrix rhusiopathiae*; swine dysentery associated with *Treponema hyodysenteriae* when followed by appropriate medication in the drinking water and/or feed.

Dogs: Treatment of upper respiratory infections such as bronchitis, tracheobronchitis, tracheitis laryngitis, tonsillitis, and pneumonia caused by *Staphylococci* species, hemolytic *Streptococci* species and *Pasteurella multocida*.

Cats: Treatment of upper respiratory infections when caused by *Staphylococci* species and hemolytic *Streptococci* species and for feline pneumonitis when caused by tylosin susceptible organisms.

TARGET SPECIES

Cattle, pigs, dogs and cats.

ADMINISTRATION AND DOSAGE

Cattle: 18 mg/kg bw a day, equivalent to 0.9 ml of product/10 kg bw a day, up to 5 consecutive days.

Swine: 9 mg/kg bw twice a day, equivalent to 0.45 ml of product/10 kg bw twice a day, up to 3 consecutive days.

Dogs and Cats: 6 - 10 mg/kg bw, equivalent to 0.03 to 0.05 ml of product/kg bw, at 12 to 24 hours interval. A one ml fine dosage syringe should be used.

ROUTE OF ADMINISTRATION

Intramuscular injection.

CONTRAINDICATIONS

It should not be given to chickens or turkeys. Do not administer to horses or other equines in which injection of tylosin may be fatal.

WARNING AND PRECAUTIONS

SPECIAL CAUTIONS AND WARNINGS FOR ITS USE

Cattle: When administered more than 10 ml, distribute the dose in different sites of injection.

Pigs: When administered more than 5 ml, distribute the dose in different sites of injection.

Cats and Dogs: For repeated use, change injections site. Irritation and pain at injection site may be observed occasionally.

Special cautions for people administering or handling the product: Use under veterinarian recommendation.

SPECIAL CAUTIONS FOR PEOPLE ADMINISTERING OR HANDLING THE PRODUCT

Not required.

Use under veterinarian recommendation.

INTERACTIONS WITH OTHER MEDICINAL PRODUCTS

None observed.

INCOMPATIBILITIES

Do not administer along with bactericide antibiotics.

USE DURING PREGNANCY AND LACTATION

Do not use in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older, including dry dairy cows. Use in these cattle may cause drug residues in milk and/or in calves born to these cows.

SIDE EFFECTS (FREQUENCY AND SEVERITY)

In very rare cases the following have been observed;

- Swelling/inflammation at the site of injection,
- Vulvular swelling in cattle,
- Oedema of the rectal mucosa, partial anal protrusion ('rosebudding'), erythema and pruritus in pigs.

SYMPTOMS AND TREATMENT OF OVERDOSAGE

Discontinue medication and seek for veterinarian advice if there is sign of overdose.

WITHDRAWAL PERIOD

Pigs: Meat: 14 days.

Cattle: Meat: 21 days.

PROPOSED SHELF LIFE (AS PACKAGE FOR SALE)

3 years.

PROPOSED SHELF LIFE (AFTER FIRST OPENING OF THE CONTAINER)

28 days.

PRESENTATIONS

100 ml and 250 ml

DATE OF REVISION OF PACKAGE INSERT

8th July 2014

Store in a cool, dry place (recommended below 25° C), protected from intensive light.

Use under veterinarian recommendation

keep medicine out of reach of children
Jauhkan ubat dari jangkauan kanak-kanak

Any unused veterinary medicinal product waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Controlled Medicine

Country Reg. Nr.:

Distributed by:

FE VENTURE Sdn Bhd

Lot 3 & 5, Jalan Pis 11/8
Bandar Sunway 46150 Petaling Jaya
Selangor Darul Ehsan - MALAYSIA
Phone: 60 3 563 33 493 Fax: 60 3 563 23 410

Registration Holder:

LABORATORIOS REVEEX ASIA Sdn Bhd

Lot 3 & 5, Jalan Pis 11/8
Bandar Sunway 46150 Petaling Jaya
Selangor Darul Ehsan - MALAYSIA
Phone: 60 3 563 33 493 Fax: 60 3 563 23 410



Manufactured by:

s.p. veterinaria, s.a.

Ctra. Reus-Vinyols, km. 4.1 • 43330 RIUDOMS (Tarragona) - Spain
Tel. +34 977 850 170 • Fax +34 977 850 405 • P.O.Box. 60