

TILCOSIN 100mg/ml Oral Solution

FOR ANIMAL USE ONLY

Brand or Product Name :

TILCOSIN 100mg/ml Oral Solution

Name and Strength of Active Substance(s)

Each ml contains 100mg Tilmicosin Phosphate and 0.1 %w/v Potassium Sorbate as preservative.

Product Description

TILCOSIN 100mg/ml Oral Solution is a light yellowish to brownish liquid.

Pharmacodynamics

Tilmicosin is a mainly bactericidal semi-synthetic antibiotic of the macrolide group. It is believed to affect bacterial protein synthesis.

Tilmicosin has a wide spectrum of activity against Gram-positive organisms and is particularly active against *Pasteurella*, *Actinobacillus* (*Haemophilus*) and *Mycoplasma* organisms of bovine, porcine and avian origin. Tilmicosin has some activity against certain Gram-negative micro-organisms. Cross resistance between tilmicosin and other macrolide antibiotics has been observed.

Pharmacokinetics

When administered orally to chickens or turkeys via the drinking water at recommended concentrations, tilmicosin is absorbed and moves rapidly out of the serum into areas of low pH. This results in very low serum concentrations, but detectable levels are found in lung tissue as early as 6 hours after commencing treatment. In chickens or turkeys, tilmicosin is also detected in pooled airsac tissue as early as 6 hours after commencing treatment.

Indication

For the treatment of respiratory infections in chicken caused by *Mycoplasma gallisepticum*, *M. synoviae* and other organisms sensitive to Tilmicosin.

Recommended Dosage

Chicken (Broiler) : 15 – 20 mg Tilmicosin per kg body weight for 3 days or 0.15 – 0.2 ml TILCOSIN 100mg/ml Oral Solution per kg body weight daily for 3 days.

Route of Administration

To be administered orally via animal drinking water.

Contraindications

The use of this drug is contraindicated in animals with a history of an allergic reaction to Tilmicosin.

Warnings and Precautions

Special Warnings:

Important: Must be diluted before administration to animals.

Water consumption should be monitored in order to guarantee adequate dosing. In case water consumption does not match quantities for which recommended concentrations were calculated, the concentration of the product has to be adapted in a way that the recommended dosage will be taken up by the animals or different medication should be considered.

Special precautions for use

(i) Special precautions for use in animals

For oral use only. Do not inject. Inappropriate use of the product may increase the prevalence of bacteria resistant to tilmicosin and may decrease the effectiveness of treatment with tilmicosin-related substances. The use of the product should be based on susceptibility tests. Medicated drinking water or milk replacer should be prepared fresh every 24 hours. Official, national and regional antimicrobial policies should be taken into account when the product is used.

(ii) Special precautions to be taken by the person administering the veterinary medicinal product to animals

- Tilmicosin may induce irritation. Macrolides, such as tilmicosin, may also cause hypersensitivity (allergy) following injection, inhalation, ingestion or contact with skin or eye. Hypersensitivity to tilmicosin may lead to cross reactions to other macrolides and vice versa. Allergic reactions to these substances may occasionally be serious and therefore direct contact should be avoided.

- To avoid exposure during preparation of the medicated drinking water, wear overalls, safety glasses, and impervious gloves. Do not eat, drink or smoke when handling this product. Wash hands after use.

- In the case of accidental ingestion, wash out mouth immediately with water and seek medical advice. In the event of accidental skin contact, wash thoroughly with soap and water. In case of accidental eye contact, flush the eyes with plenty of clean, running water.

- Do not handle the product if you are allergic to ingredients in the product.

- If you develop symptoms following exposure, such as skin rash, you should seek medical advice and show the physician this warning. Swelling of the face, lips and eyes or difficulty in breathing are more serious symptoms and require urgent medical attention.

Interactions with Other Medicaments

Cross resistance between tilmicosin and other macrolide antibiotics and lincosamides has been observed. Tilmicosin may lessen the antibacterial activity of β -lactam antibiotics.

Statement on usage during pregnancy and lactation

Pregnancy : The safety of tilmicosin has not been established in animals used for breeding purposes. When considering treatment for life-threatening infection and/or prolonged duration of therapy, the potential risk to the fetus must be balanced against the severity of the potential illness.

Lactation : The safety of tilmicosin has not been established.

Side Effects

In very rare cases, a decrease in water intake has been observed.

Overdose and Treatment

No symptoms of overdose have been seen in chickens given drinking water containing levels of tilmicosin up to 375 mg/litre for 5 days. No symptoms of overdose have been seen in turkeys given drinking water containing levels of tilmicosin up to 375 mg/litre for 3 days or 75 mg/litre for 6 days.

Withdrawal Period(s)

Poultry meat : 12 days. Eggs: Do not use. Not for use in animals from which eggs are produced for human consumption.

Shelf-life

As packaged for sale: 24 months

After first opening the immediate packaging: 24 hours

After reconstitution or dilution : Medicated water must be prepared fresh daily. Any unused medicated water is to be discarded after 24 hours.

Maximum Residual Limit (MRL)

Muscle, fat (poultry) = 100 μ g/kg, [Ref. REGOVP, Version 2, dated Dec 2009]

Storage Conditions

Do not store above 25°C. Keep out of reach of children. Jauhi Ubat daripada kanak-kanak.

Disposal of Containers

Empty bottle must be disposed of in accordance to local government regulations.

Dosage Forms and Packaging Available

Available in 1 litre per bottle, pack in 12 bottles or 20 bottles per carton.

Name and Address of Manufacturer

Life Biopharma Sdn. Bhd. (company no. 546163-D)

No. 250-251, Jalan Industri Galla 13, Kawasan Perindustrian Galla, 70200 Seremban, Negeri Sembilan, Malaysia.

Date of Revision of Package Insert

2014-11-04