

VETFLOR 50g/kg PREMIX POWDER

FOR ANIMAL USE ONLY

Brand or Product Name :

VETFLOR 50g/kg PREMIX POWDER

Name and Strength of Active Substance(s)

Each kg contains 50g Florfenicol.

Product Description

VETFLOR 50g/kg PREMIX POWDER is a white to off-white powder.

Pharmacodynamics Properties

Florfenicol is a broad-spectrum synthetic antibiotic in the phenicol group that is active against most Gram-positive and Gram-negative bacteria isolated from domestic animals. Florfenicol acts by inhibition of protein synthesis at the ribosomal level and is bacteriostatic.

Pharmacokinetics Properties

After administration to pigs by gavage at 10 mg/kg under experimental conditions, absorption of florfenicol was variable but peak serum concentrations of approximately 5 µg/ml were reached approximately 3 hours after dosing. The terminal half-life was between 3 and 4 hours. After a single treatment the serum concentrations of florfenicol remained above 1 µg/ml for 12 to 18 hours. After absorption and distribution, florfenicol is extensively metabolised by pigs and rapidly eliminated, primarily in urine and the faeces in a ratio of 3:1.

A fraction is secreted unchanged, the rest is metabolised to 5 metabolites. After parenteral dosing of florfenicol to pigs, it has been shown that lung concentrations are similar to serum concentrations.

Indications

For the control of swine respiratory disease (SRD) associated with *Actinobacillus pleuropneumoniae*, *Pasteurella multocida*, *Streptococcus suis*, and *Bordetella bronchiseptica* in groups of swine in buildings experiencing an outbreak of SRD.

Recommended Dosage

Swine : Use 3.64 kg VETFLOR 50g/kg PREMIX POWDER per ton of feed (equivalent to 182g florfenicol per ton), feed continuously as the sole ration for 5 days.

Mixing Directions :

To ensure thorough dispersion of the product it should first be mixed with a suitable quantity of feed before incorporation into the final mix.

Mode of Administration

To be mixed thoroughly with animal feed.

Contraindications

Do not use in boars intended for breeding purposes. Do not administer in cases of previous allergic reactions to florfenicol.

Warnings and Precautions

Warnings : Avoid inhalation, oral exposure, and direct contact with skin or eyes. Operators mixing and handling this product should use protective clothing, gloves, goggles and a NIOSH-approved duct mask. Wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water. If irritation persists, seek medical attention. Not for human consumption. Keep out of reach of children.

Precautions : Feed containing florfenicol shall not be fed to pigs for more than 5 days. Following administration, pigs should be re-evaluated by a licensed veterinarian before reinitiating further course of therapy.

Interactions with Other Medicaments

There is inadequate data on adverse drug interactions produced by florfenicol.

Statement on usage during pregnancy and lactation

The safety of the product during pregnancy and lactation has not been studied in sows. Use of the product during pregnancy and lactation is therefore not recommended.

Side Effects

Commonly observed adverse effects are diarrhoea, perianal inflammation and rectal eversion. These effects are transient, resolving on cessation of treatment.

Overdose and Treatment

In case of overdosing, a decrease in weight gain, food and water consumption, peri-anal erythema and oedema and modification of some haematological and biochemical parameters indicative of dehydration may be observed.

Storage Conditions

Store airtight and below 30°C. Keep out of reach of children. Jauhi ubat dari kanak-kanak.

Shelf-life

3 years from date of manufacture. Opened bag and reconstituted feed is recommended to be used within 24 hours.

Withdrawal Period(s)

Feeds containing florfenicol must be withdrawn 14 days prior to slaughter.

Allowable Maximum Residual Limit (MRL)

Porcine : Muscle = 300 µg/kg, Skin + fat = 500 µg/kg, Liver = 2000 µg/kg, Kidney = 500 µg/kg [Ref. EMEA/MRL/822/02-FINAL, January 2002]

Disposal of Containers

Emptied bag can be safely disposed in trash according to local government regulations.

Dosage Forms and packaging available

25 kg in PP lined paper bags.

Name and Address of manufacturer

Life Biopharma Sdn. Bhd. (company no. 546163-D)

No. 250-251, Jalan Industri Galla 13, Kawasan Perindustrian Galla, 70200 Seremban, Negeri Sembilan, Malaysia.

Date of Revision of Package Insert

2014-12-31