


PRODUCT INFO			
BLUE #:	213311AP	Product Name:	Tylan 200 Injection
Item Code:	PA213311X	Component:	Package Inserts and Leaflets
Product Code:	AH0206	Pack Size:	100ml
Previous Item Code	PA280512X		

ARTWORK INFO			
Template:	Pl_148x210mm_Norbrook_v1_1	Packaging Spec(s):	Base Spec MXG410 Primary Spec N/A
Barcodes/Type:	Pharmacode: 304	Add. Info:	Please note new pharmacode assigned
		Minimum Core Data Point Size:	8 point
GTIN:	GTIN Not Required	Proof #	P1a
		By/Date	SG 03-MAR-2022
NON PRINTING BLACK			

ELANCO ARTWORK LEGEND v16



148mm

Pharma code bar height = 12 mm

210mm

75 mm

12 mm

12 mm

BARCODE READ DIRECTION

Elanco™

AH0206

PA213311X

Tylan™ 200

Tylosin 200 mg/ml

PRODUCT NAME: Tylan 200 mg/ml Injection

ACTIVE INGREDIENT: 200 mg/ml Tylosin

PRESERVATIVE:

518 mg/ml Propylene Glycol (50% V/V)

41.8 mg/ml Benzyl Alcohol (4% V/V)

FOR INTRAMUSCULAR USE IN CATTLE AND PIGS

PRESENTATION:

It is a clear solution and is essentially free from visual particles. A parenteral solution of Tylosin Base in 50% Propylene Glycol, 4% Benzyl Alcohol. Each ml contains 200 mg of Tylosin Activity.

PHARMACODYNAMICS:

Tylosin is a macrolide antibiotic produced by a strain of *Streptomyces fradiae*. It exerts its antimicrobial effect by inhibiting protein synthesis of susceptible micro-organisms. The tylosin spectrum of activity includes Gram-positive bacteria, some Gram-negative strains such as *Pasteurella*, and *Mycoplasma* spp at concentrations of 16 µg/ml or less.

PHARMACOKINETICS:

Absorption: Following intramuscular injection, tylosin blood levels peak 1-2 hours post-injection. Duration of activity is approximately 12 hours.

Distribution, Biotransformation and Elimination:

Tylosin levels of 1.4 to 1.6 and 2.2 to 6.7 µg/ml were recorded in serum and lung tissue respectively following intramuscular injection of 8.8 mg/kg bodyweight in pigs. Measurable amounts of tylosin were still present in both serum and lung tissue at 12 hours post-injection. Tylosin concentrations were greater in lung tissue than serum at all sample times.

INDICATIONS:

Tylan 200 mg/ml Injection is indicated in all conditions associated with bacteria sensitive to tylosin which includes organisms in the following genera:

<i>Streptococcus</i>	<i>Campylobacter</i>
<i>Bacillus</i>	<i>Spirochaetes</i>
<i>Staphylococcus</i>	<i>Mycoplasma</i>
<i>Corynebacterium</i>	<i>Fusiformis</i>
<i>Clostridium</i>	<i>Pasteurella</i>
<i>Erysipelothrix</i>	<i>Chlamydia</i>

Tylan 200 mg/ml Injection has been successfully used in respiratory and genito-urinary tract infections, otitis, cellulitis and secondary bacterial conditions associated with virus disease or post operative infections. Specific disease entities treated successfully with Tylan 200 mg/ml Injection include: Swine Dysentery, Erysipelas and Enzootic Pneumonia in pigs, foul in the foot, mastitis and calf pneumonia in cattle.

DOSAGE AND ADMINISTRATION:

Tylan 200 mg/ml Injection should be given by intramuscular injection at the following dose rates:

Cattle and calves: 4 to 10 mg per kg bodyweight daily.

Pigs: 2 to 10 mg per kg bodyweight daily.

If there is no response to treatment in 3 days, diagnosis and treatment should be reassessed. The maximum injection volume for cattle is limited to 15 ml per injection site. To ensure the correct dosage, bodyweight should be determined as accurately as possible to avoid under dosing.

CONTRA-INDICATIONS:

Tylan 200 mg/ml Injection should not be given to chickens or turkeys. Do not use in animals with known hypersensitivity to the active ingredients. Do not use in equine animals.

INTERACTION WITH OTHER MEDICATION:

None.

INCOMPATIBILITIES:

Do not mix with other solutions, since this may cause precipitation of the active ingredient.

PREGNANCY AND LACTATION:

No adverse effects to tylosin have been seen in fertility, multi-generation or teratology studies.

SYMPTOMS AND TREATMENT OF OVERDOSE:

Pigs and calves: Intramuscular injection of 30 mg/kg bodyweight per day (three times maximum recommended dose) for five days produced no adverse effects. The LD₅₀ for subcutaneous injection of tylosin in mice is estimated to be >2500 mg/kg bodyweight.

Shaded Areas = No text areas

148mm

210mm

WARNINGS AND PRECAUTIONS:

Special warnings for each target species

None.

Special precautions for use

(i) Special precautions for use in animals

Use of the product should be based on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local epidemiological information. For administration by the intramuscular route only. Use different injection sites for repeat injections.

(ii) Special precautions to be taken by the person administering the veterinary medicinal product to animals

Care should be taken to avoid accidental self-injection. If accidental self-injection occurs, seek medical attention immediately. In the event of accidental skin contact, wash thoroughly with soap and water. In case of accidental eye contact, flush the eyes with plenty of clean, running water. Wash hands after use. Tylosin may induce irritation. Macrolides, such as tylosin, may also cause hypersensitivity (allergy) following injection, inhalation, ingestion or contact with skin or eye. Hypersensitivity to tylosin may lead to cross reactions to other macrolides and *vice versa*. Allergic reactions to these substances may occasionally be serious and therefore direct contact should be avoided. Do not handle the product if you are allergic to ingredients in the product. If you develop symptoms following exposure, such as skin rash, you should seek medical advice and show the physician this warning. Swelling of the face, lips and eyes or difficulty in breathing are more serious symptoms and require urgent medical attention.

(iii) Other precautions

Blemishes may occur at the site of injection and can persist for up to 21 days following administration.

ADVERSE REACTIONS:

Possible adverse reactions attributed to the product when used as recommended and their frequency are: In very rare cases the following have been observed;

- swelling/inflammation at the site of injection.
- vulvular swelling in cattle.
- oedema of the rectal mucosa, partial anal protrusion ("rosebudding"), erythema and pruritus in pigs.
- Anaphylactic shock and death.

WITHDRAWAL PERIODS:

Swine: Meat - 9 days

Cattle: Meat - 28 days

Milk - 108 hours.

With cows milked twice daily, milk for human consumption may be taken only from 108 hours

(i.e. at the 9th milking) after the last treatment. With other dosing routines, the basis of the veterinary surgeons advice should be that milk may be taken for human consumption only after the same period from the last treatment (i.e. with three times a day milking, milk for human consumption may be taken only from 108 hours i.e. at the 14th milking).

STORAGE CONDITION:

Store at 25°C or below (Air conditioning). Use within 24 hours once opened.

DISPOSAL OF CONTAINER AND NEEDLES:

Dispose of empty container by wrapping with paper and putting in garbage. Discarded needles should immediately be placed in a designated and appropriately labelled "sharps" container.

PACK SIZE:

100 ml.

MANUFACTURED BY:

NORBROOK LABORATORIES LIMITED

105 Armagh Road, Newry

Co. Down BT35 6PU

Northern Ireland.

FOR:

Elanco Malaysia Sdn Bhd

Unit 5.04, Level 5 & 6, Tower Block, The Bousteador,

No. 10, Jalan PJU 7/6, Mutiara Damansara,

47800 Petaling Jaya, Selangor, Malaysia.

CONTROLLED MEDICINE

MAL15010009HA

For animal use only.

Keep out of reach of children.

Jauhi ubat dari kanak-kanak.

Date of revision of package insert: March 2022

Tylan, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

© 2022 Elanco or its affiliates

Shaded Areas = No text areas