

SAME SIZE ARTWORK
LEAFLET SIZE: 276 mm x 135 mm
FOLDING SIZE: 23 mm x 135 mm

25 mm

Treatment with higher than recommended doses may result in clinically significant adrenal suppression. If there is evidence for higher than recommended doses being used, then additional systemic corticosteroid cover should be considered during periods of stress or elective surgery.

In a placebo-controlled clinical trial in which paediatric patients were administered Mometasone Furoate Nasal Spray 100 micrograms daily for one year, no reduction in growth velocity was observed.

Safety and efficacy of Mometasone Furoate Nasal Spray in children and adolescents under 18 years of age have not been established.

During transfer from systemic corticosteroids to Mometasone Furoate Nasal Spray, some patients may experience symptoms of withdrawal from systemically active corticosteroid (e.g., joint and/or muscular pain, lassitude, and depression initially) despite relief from nasal symptoms and will require encouragement to continue Mometasone Furoate Nasal Spray therapy. Such transfer may also unmask pre-existing allergic conditions, such as allergic conjunctivitis and eczema, previously suppressed by systemic corticosteroid therapy.

Patients receiving corticosteroids who are potentially immunosuppressed should be warned of the risk of exposure to certain infections (e.g., chickenpox, measles) and of the importance of obtaining medical advice if such exposure occurs.

Following the use of intranasal aerosolized corticosteroids, instances of nasal septum perforation or increased intraocular pressure have been reported very rarely.

Acute Rhinosinusitis

If signs or symptoms of severe bacterial infection are observed (such as fever, persistent severe unilateral facial/tooth pain, orbital or peri-orbital facial swelling, or worsening of symptoms after all initial improvement). The patient should be advised to consult their physician immediately.

Safety and efficacy of Mometasone Furoate Aqueous Nasal Spray for the treatment of symptoms of rhinosinusitis in children under 12 years of age have not been studied.

General Nasal Corticosteroid Warning. Systemic effects of nasal corticosteroids may occur, particularly at high doses prescribed for prolonged periods. Growth retardation has been reported in children receiving corticosteroids at licensed doses.

It is recommended that the height of children receiving prolonged treatment with nasal corticosteroids is regularly monitored. If growth is slowed, therapy should be reviewed with the aim of reducing the dose of nasal corticosteroid, if possible, to the lowest dose at which effective control of symptoms is maintained. In addition, consideration should be given to referring patient to a paediatric specialist.

USE DURING PREGNANCY AND LACTATION:

There are no adequate or well controlled studies in pregnant women. As with other nasal corticosteroid preparations, Mometasone Furoate Aqueous Nasal Spray should be used in pregnant women, nursing mothers or women of childbearing age only if the potential benefit justifies the potential risk to the mother, fetus or infant. Infants born of mothers who received corticosteroids during pregnancy should be observed carefully for hypoadrenalism.

OVERDOSE

Because the systemic bioavailability of mometasone furoate nasal spray is <1% (using a sensitive assay with lower quantitation limit of 0.25 pg/ml), overdose is unlikely to require any therapy other than observation, followed by initiation of the appropriate prescribed dosage.

INCOMPATIBILITIES

Not applicable.

SHELF-LIFE

24 Months

STORAGE

Store below 30°C. Do not freeze. Protect from light. Shake well before use.

PRESENTATION

1) 60 Metered Doses

9.0 g suspension filled in HDPE bottle and crimped with a metered pump and fixed with adaptor suitable for nasal spray.

2) 120 & 140 Metered Doses

18.0 g suspension filled in HDPE bottle and crimped with a metered pump and fixed with adaptor suitable for nasal spray.

MAL NUMBER

MAL15015062AZ

INSTRUCTIONS FOR PATIENTS FOR USING MOMATE NASAL SPRAY

i) How to use your nasal spray

1. Remove the plastic cap (Figure 1).



(Figure 1)

2. Before you use *Momate* Nasal Spray for the first time prime the pump by pressing downward on the shoulders of the white nasal applicator using your index finger and middle finger while holding the base of the bottle with your thumb (Figure 2). Do not pierce the nasal applicator. Press down and release the pump 10 times or until a fine spray appears. Do not spray into eyes. The pump is now ready to use. The pump may be stored unused for upto 1 week without repriming. If unused for more than 1 week, reprime by spraying 2 times or until a fine spray appears.




(Figure 2)

3. Gently blow your nose to clear the nostrils. Close 1 nostril. Tilt your head forward slightly, keep the bottle upright, and carefully insert the nasal applicator into the other nostril (Figure 3). Do not spray directly onto the nasal septum (the wall between the two nostrils.).



(Figure 3)

4. Then breath out through the mouth.
5. Repeat in the other nostril.
6. Wipe the nasal applicator with a clean tissue and replace the plastic cap (Figure 4).




(Figure 4)

ii) Cleaning your nasal spray


It is important to clean your nasal spray regularly; otherwise it may not work properly.

1. To clean the nasal applicator, remove the plastic cap (Figure 1).




(Figure 1)

2. Pull gently upward on the white nasal applicator to remove (Figure 2).



(Figure 2)


3. Soak the Nozzle and Dust Cap in warm water for few minutes and then rinse thoroughly spray tip and dust cap in warm water and finally rinse under clean running water. Repeat the procedure if the mist is not coming out (Figure 3).



Note:

Incase during usage for the first time if the actuator is observed to be blocked condition due to accidental pressing during transit or during handling which prompts release of miniscule contents, never use a pin or any sharp objects to pierce to unblock the nozzle instead subject the actuator for cleaning.

4. Put the nasal applicator back together making sure the pump stem is reinserted into the applicator's center hole (Figure 4).



(Figure 4)

5. Reprime the pump by pressing downward on the shoulders of the white nasal applicator using your index and middle fingers while holding the base of the bottle with your thumb. Press down and release the pump 2 times or until a fine spray appears. Do not spray into eyes. The pump is now ready to use. The pump may be stored unused for upto 1 week without repriming. If unused for more than 1 week, reprime by spraying 2 times or until a fine spray appears (Figure 5).



(Figure 5)

6. Replace the plastic cap (Figure 6).



(Figure 6)

Date of Revision: 11 October 2022

Product Registration Holder:
Glenmark Pharmaceuticals
(Malaysia) Sdn Bhd (660397-W)
D-31-02, Menara Suezcap 1, No. 2, Jalan Kerinchi,
59200 Kuala Lumpur, Malaysia
Manufactured by :

glenmark
PHARMACEUTICALS LTD.

B/2, Mahalaxmi Chambers,
22, Bhuaibhai Desai Road, Mumbai - 400 026, India.
At: Unit III, Village Kishanpura Baddi Nalagarh Road,
Tehsil Baddi, Dist Solan (H.P.) -173 205, India
TM Trade Mark

MY
PE00000

ICONGRAPHICS CODE:

PANTONE SHADE



PANTONE
BLACK
PROCESS C

**Supersedes
Artwork Code:
PE50767**

PHARMACODE: