

	ANTIBIOTIC/ 抗生素製劑 FOR VETERINARY USE ONLY 動物用藥 10 KG JINPLUS 200 G/KG GRANULES 金順200 預混劑顆粒 Each kilogram contains 200g of Tilmicosin Phosphate and 5g of Propylparaben as preservative. Indication: For the prevention and treatment of respiratory disease caused by <i>Actinobacillus pleuropneumoniae</i>, <i>Mycoplasma hyopneumoniae</i>, <i>Pasteurella multocida</i> and other organisms sensitive to tilmicosin. Dosage and administration: Administer orally in the feed at a dose of 8 to 16 mg/kg body weight/day of tilmicosin (equivalent to 200 to 400 ppm in the feed) for a period of 15 to 21 days. Mix 1-2 kg of Jinplus 200G in 1 tonne of feed. Contraindication: Do not allow horses and other equines access to feed containing tilmicosin. Do not use in case of hypersensitivity to tilmicosin or to any of the excipients. Interactions with Other Medicament: None known. Storage condition: Store in a dry place at or below 30°C, and protect from direct sunlight. Shelf life after reconstitution: 24 hours. Shelf Life: 2 years Withdrawal Period: Withdrawal period in pig is 21 days. Disposal: Disposal must be in accordance with applicable federal, state/provincial, and/or local regulations. Incineration is the preferred method of disposal, when appropriate. Product Description: A yellowish brown powder with granules. Pharmacodynamics: Tilmicosin is a semi-synthetic antibiotic of the macrolide group and is believed to affect protein synthesis. It has bacteriostatic action but at high concentrations it may be bactericidal. This antibacterial activity is predominantly against Gram-positive microorganism with activity against certain gram-negative ones and Mycoplasma of a bovine, porcine, ovine and avian origin. In particular, its activity has been demonstrated against the following microorganism: <i>Mycoplasma hyopneumoniae</i>, <i>Pasteurella multocida</i> and <i>Actinobacillus pleuropneumoniae</i>. Scientific evidence suggests that macrolides act synergistically with the host immune system. Macrolides appear to enhance phagocyte killing of bacteria. Tilmicosin has been shown to inhibit in vitro the replication of the Porcine Reproductive and Respiratory Syndrome virus in alveolar macrophages in a dose dependent fashion. Cross resistance between tilmicosin and other macrolides and lincomycin has been observed. Pharmacokinetics: <u>Absorption:</u> When administered to pigs via the oral route at a dose of 400 mg tilmicosin/kg feed (equivalent to approximately 21.3 mg tilmicosin/kg bodyweight/day), tilmicosin moves rapidly out of the serum into areas of low pH. The highest concentration in the serum (0.23±0.08 µg/ml) was recorded on day 10 of medication, but concentrations above the limit of quantification (0.10 µg/ml) were not found in 3 out of 20 animals examined. Lung concentrations increased rapidly between days 2 and 4 but no significant changes were obtained following four days of dosing. The maximum concentration in lung tissue (2.59±1.01 µg/ml) was recorded on day 10 of medication. When administered at a dose of 200 mg tilmicosin/kg feed (equivalent to approximately 11.0 mg/kg/day), plasma concentrations above the limit of quantification (0.10 µg/ml) were found in 3 out of 20 animals examined. Quantifiable levels of tilmicosin were found in lung tissue with the maximum concentration (1.43±1.13 µg/ml) being recorded on day 10 of medication. <u>Distribution:</u> Following oral administration, tilmicosin is distributed throughout the body with especially high levels found in the lung and in lung tissue macrophages. It is also distributed in the liver and kidney tissues. <u>Biotransformation:</u> Several metabolites are formed, the predominant one being identified as T1. However the bulk of tilmicosin is excreted unchanged. <u>Elimination:</u> Following oral administration, tilmicosin is excreted mainly via the bile into the faeces, but a small proportion is excreted via the urine. Warning and Precautions: - Under practical conditions, the management of respiratory disease outbreaks recognises that acutely ill animals are inappetant and require parenteral therapy. - Inappropriate use of the product may increase the prevalence of bacteria resistant to tilmicosin and may decrease the effectiveness of treatment with tilmicosin related substances. - Official, national and regional antimicrobial policies should be taken into account when the product is used. - Due to the likely variability (time, geographical) in the occurrence of the resistance of bacteria for tilmicosin, bacteriological sampling and susceptibility testing are recommended. - Tilmicosin may induce irritation. Macrolides, such as tilmicosin, may also cause hypersensitivity (allergy) following injection, inhalation, ingestion or contact with skin or eye. Hypersensitivity to tilmicosin may lead to cross reactions to other macrolides and vice versa.	
	ANTIBIOTIC/ 抗生素製劑 FOR VETERINARY USE ONLY 動物用藥 10 KG JINPLUS 200 G/KG GRANULES 金順200 預混劑顆粒 Allergic reactions to these substances may occasionally be serious and therefore direct contact should be avoided. - To avoid exposure during preparation of the medicated feed, wear overalls, safety glasses, impervious gloves and wear either a disposable half mask respirator conforming to European Standard EN149 or a non-disposable respirator to European Standard EN140 with a filter to EN143. Do not eat, drink or smoke when handling this product. Wash hands after use. - In the case of accidental ingestion, wash out mouth immediately with water and seek medical advice. In the event of accidental skin contact, wash thoroughly with soap and water. In case of accidental eye contact, flush the eyes with plenty of clean, running water. - Do not handle the product if you are allergic to ingredients in the product. - If you develop symptoms following exposure, such as skin rash, you should seek medical advice and show the physician this warning. Swelling of the face, lips and eyes or difficulty in breathing are more serious symptoms and require urgent medical attention. Pregnancy and Lactation: The safety of Tilmicosin has not been established in pregnant sows and animals for breeding purposes. Side Effect: In very rare cases, feed intake may decrease (including feed refusal) in animals receiving medicated feed. This effect is transient. Symptoms and Treatment of Overdose: No symptoms of overdose have been seen in pigs fed a ration containing levels of tilmicosin up to 80 mg/kg bodyweight (equivalent to 2000 ppm in the feed or ten times the recommended dose) for 15 days. 每公斤含200克Tilmicosin Phosphate和 5克 Propylparaben 防腐劑 適應症：用於預防和治療由胸膜肺炎放線桿菌、肺炎黴漿菌、敗血性巴氏桿菌和其他對Tilmicosin具感受的微生物病原體所引起的呼吸道疾病 劑量和用法：在口服飼料中添加8至16毫克/公斤體重/天的Tilmicosin（相當於飼料中200至400 ppm）的劑量，持續15至21天。 在1噸飼料中混合1-2公斤Jinplus 200G. 儲存情況： 存放在30°C 或以下的干燥地方，并避免陽光直射。 保質期: 2 年 停藥期：停藥期豬為21天。 混合後保質期： 24小时 Keep medicine out of reach of children / 遠離兒童 /Jauhkan daripada kanak-kanak. MAL No.: 15025045HA Batch No.: Mfg Date: Exp Date: Product of YSP Manufacturer: Vetnostrum Animal Health Co. Ltd., Hsinchu Plant No. 290-3, Chung Lun, Chung Lun Village, Hsinfeng, Hsinchu, Taiwan (R.O.C) Product Registration Holder and Distributor: Y.S.P. INDUSTRIES (M) SDN. BHD. Lot 3, 5 & 7, Jalan P/7, Section 13, Kawasan Perindustrian Bandar Baru Bangi, 43000 Kajang, Selangor Darul Ehsan Malaysia. Ordering Line: 1 800 88 3027 Product Info Line: 1 800 88 3679 A subsidiary of Y.S.P.SAH® Revised on: Jan 2021 UBAT TERKAWAL CONTROLLED MEDICINE	