

PACKAGE INSERT

Diclazuril Premix 0.5% w/w Granular

Brand or product Name	Diclazuril Premix 0.5% w/w Granular
Name and strength of active substance(s)	Diclazuril 0.5% (w/w)
Product Description	Yellow to brown granular
Pharmacodynamics	Diclazuril is a chemical anticoccidial, a benzeneacetonitrile derivative-synthetic compound of the triazinone family. Diclazuril has a mode of action completely different from the existing ionophores and chemicals, available on the market. According to the anticoccidial properties of triazinones, diclazuril is active against intracellular development stages of coccidia, namely during schizogony and gametogony. After diclazuril treatment, the first- and second- generation schizonts show extensive degenerative changes which are characterised by the loss of internal structure, the appearance of many intracytoplasmic vacuoles and incomplete merogony. Ultrastructural findings indicate that diclazuril treatment primarily affects the normal differentiation of the respective endogenous stages during parasite development. This leads to complete degeneration of schizonts and gamonts.
Pharmacokinetics	The absorption of diclazuril in animal is poor after administration. Maximum concentrations in plasma are reached about 24 hours after dosing. The absorption decreases with the animals' age. The elimination half-life is about 30 hours. In-vitro studies on sheep hepatocytes demonstrated that metabolic transformation of diclazuril is limited. This was equally observed in other animal species. Excretion occurs almost completely via the faeces.
Environmental Properties	None known
Indication	Prevention of coccidiosis disease in broiler chickens
Recommended Dosage	Thoroughly mix 200 g of Diclazuril Premix 0.5 % per ton of feed obtain a concentration of 1 ppm Diclazuril (1 gram per ton).
Mode of Administration	Orally via feed mixing.
Contraindication	None known
Warning and precaution	Treated poultry must not be slaughtered for human consumption during treatment. As with all medicated feedstuffs, operators are advised to use protective clothing, gloves, dust mask and goggles when mixing or handling Diclazuril premix.

Interaction with other medicaments	Diclazuril Premix should not be mixed or used simultaneously with any other anticoccidial.
Statement on usage during pregnancy and lactation	Not for use in hens producing eggs for human foods.
Adverse effects/Undesirable Effects	Not known.
MRL	<u>Poultry:-</u> Muscle - 500 µg/kg Fat - 1000 µg/kg Kidney - 2000 µg/kg Liver - 3000 µg/kg
Overdose and treatment	None known
Withdrawal Periods	5 days.
Storage conditions	Store below 25 °C.
Dosage form and packaging available	Dosage form: Granules Packaging available: 25 kgs/bag
Name and Address of manufacturer	Shandong Luxi Animal Medicine Share Co., Ltd. No. 5, Waihuanxi Road, Qihe County, Shandong Province, P.R. China.
Name and Address of Registration holder	Weissen Company (M) Sdn Bhd, Lot 105, No. 84, Jalan Velox 23, Taman Velox, 48000 Rawang, Selangor, Malaysia.
Date of revision of package insert	01/07/2024
Shelf life of the veterinary product as packaged for sale	24 months
Shelf life after dilution or reconstitution according the directions	None.