

PACKAGE INSERT

Brand or product name	Nicarbazine Premix 25% w/w Powder
Name and Strength of Active Substance	Nicarbazine 25%
Product Description	Yellow Powder
Pharmacodynamics	The agents are absorbed separately from the digestive tract, and both are needed for anticoccidial activity. Dry crystals are strongly electrostatic and present some dry mixing problems. The precise mode of action of nicarbazin is not known.
Pharmacokinetics	Broiler chickens were fed a diet containing 15 mg/kg of nicarbazin, [¹⁴ C]-labeled in both phenylurea and pyrimidinone, for 7 days and groups of birds were sacrificed between day 2 and day 7. Concentrations of the [¹⁴ C]-labeled phenylurea portion of the complex were much higher in liver and kidney than in plasma and muscle. [¹⁴ C]-labeled pyrimidinone concentrations were highest in kidney, they were not significantly lower in muscle, plasma and liver. Liver and kidney concentrations of the pyrimidinone portion of the nicarbazin complex are about 10 times less than the concentrations of the phenylurea portion of the complex. The rapid elimination of the pyrimidinone portion of nicarbazin and the non-detection of metabolites has led to an almost exclusive focus on the phenylurea portion of the complex in subsequent metabolism and residue depletion studies.
Environmental Properties (for product used directly in the environment e.g. medicine for fish)	N/A
Indication	Aid in preventing outbreaks of cecal (<i>Elmeria tenella</i>) and intestinal (<i>E. acervulina</i> , <i>E. maxima</i> , <i>E. necatrix</i> and <i>E. brunetti</i>) coccidiosis.
Recommended Dosage	113.5 (0.0125 percent) grams Nicarbazin per ton of feed
Mode of Administration	Oral Administration
Contra-indications	Do not use in known cases of hypersensitivity to active ingredient.
Warnings and Precautions	It is forbidden for layers or breeders.
Interactions with Other Medicaments	In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other products.
Statement on usage during pregnancy and lactation	In the absence of toxicology study, this veterinary medicinal product must not be used during pregnancy and lactation. Do not use in laying birds producing eggs for human consumption.
Adverse Effects/Undesirable Effects	None known.

Overdose and Treatment	None known.
MRL	<u>Nicarbazine</u> Kidney, liver, muscle, and skin/fat (chicken) is 4 mg/kg.
Incompatibilities(for injections only)	N/A
Withdrawal period(s)	Chicken (breeders, broilers and roasters): 4 day withdrawal at doses =0.0125% and the 5 day withdrawal at doses >0.0125%
Storage conditions (may be omitted if the information is stated on the label or outer carton labels)/(should be supported by stability data)	Store in a cool and dry place below 30°C.
Dosage forms and packaging available	Dosage form: Powder Packaging available: 25kgs/bag
Name and address of manufacturer	Shandong Luxi Animal Medicine Share Co., Ltd No. 5, Waihuanxi Road, Qihe County, Shandong Province, P.R. China
Name and Address registration holder	Weissen Company (M) Sdn Bhd Lot 105, No. 84, Jalan Velox 23, Taman Velox, 48000 Rawang, Selangor, Malaysia.
Date of revision of Package insert	01.07.2024
Shelf-life of the veterinary product as packaged for sale	24 months at the temperature below 30 °C
Shelf-life after first opening the container (where relevant)	None known.
Shelf-life after first dilution or reconstitution according the directions (where relevant)	None