



維山(馬)有限公司

WEISSEN COMPANY (M) SDN. BHD. (75684-M)

No.84, Jalan Velox 23, Taman Industri Velox, 48000 Rawang, Selangor Darul Ehsan.

Tel: 016-216 6092 (HR) 016-218 1197 (ACC) Email: inquiry@weissen.com.my Web: www.weissen.com.my

PACKAGE INSERT

Brand / product name	PAN-CURE Dewormer 4% Premix Powder
Name and strength of active substance	Fenbendazole, 4 %
Product Description	Fenbendazole is sold as PAN-CURE Dewormer 4% Premix Powder which has an active ingredient concentration of 50 mg fenbendazole for each gram. This product appearance is white essentially odourless. Product form is premix powder. It is slightly soluble in water.
Pharmacodynamics	Fenbendazole is methylcarbamate benzimidazole antiparasitic agent and has a broad spectrum of activity against a variety of pathogenic internal parasites. In susceptible parasites, benzimidazole mechanism of action is believed due to disrupting intracellular microtubular transport systems by binding selectively and damage tubulin, preventing tubulin polymerization, and inhibiting microtubule formation. Benzimidazoles also act at higher concentration to disrupt metabolic enzymes, including malate dehydrogenase and fumarate reductase. Benzimidazoles may be considered time-dependent anti-parasitic agents.
Pharmacokinetics	Adsorption/ bioavailability Fenbendazole is only marginally absorbed after oral administration. The amount absorbed from the gut is apparently more associated with the solubility of the drug and not the dose given. Elimination Route In sheep, cattle and pigs, 44-50% of a dose of fenbendazole is excreted uncharged in the feces, and <1% in the urine.
Environmental Properties	N/A
Indication	A broad spectrum anthelmintic for the treatment of pigs infected with mature and immature forms of nematodes of the gastro-intestinal and respiratory tracts: Hyostrongylus rubidus (Red Stomach worm) Oesophagostomum sp. (Nodular worms) Ascaris suum (Eel worm) Trichuris suis (Whipworm) Also effective in the control of Metastrongylus apri (Lungworm). An ovicidal effect on nematode eggs.



維山(馬)有限公司

WEISSEN COMPANY (M) SDN. BHD. (75684-M)

No.84, Jalan Velox 23, Taman Industri Velox, 48000 Rawang, Selangor Darul Ehsan.

Tel: 016-216 6092 (HR) 016-218 1197 (ACC) Email: inquiry@weissen.com.my Web: www.weissen.com.my

Recommended Dosage	A manufacturer who is approved to incorporate directly at any concentration, veterinary medicinal products or premixtures containing such products must be responsible for mixing when incorporation is less than 2 kg per tonne for final feed. To be administered orally, mixed with feed. The dose rate for administration is 5 mg fenbendazole per kg bodyweight. The product may be administered to pigs either as a single dose or by divided dosage over 7 days. It may also be administered to sows over 14 days. For the treatment of infections with Trichuris and for control of Metastrongylus apri the dose of 5 mg/kg should be divided and spread over 7 days.
Mode of Administration	Mixed with the normal feed either for individual treatment or for mass treatment via a medicated feed.
Contraindication	No contraindications for the use of fenbendazole.
Warning and Precaution	Special precautions for use (i) Special precautions for use in animals When preparing medicated feed for pigs which are to be administered over a number of days, to ensure thorough dispersion a pre-mix should be prepared first. Once mixed, this should then be incorporated with the remaining feed. The product can be incorporated into pelleted feed preconditioned with steam at a temperature not greater than 45°C. Only to be incorporated by a manufacturer who are approved to mix at a rate of below 2 kg per tonne of final feed. (ii) Special precautions to be taken by the person administering the veterinary medicinal product to animals Incorporating into the feed must be performed by a suitably approved manufacturer. Impervious gloves must be worn when handling this product or feed containing this product.



維山(馬)有限公司

WEISSEN COMPANY (M) SDN. BHD. (75684-M)

No.84, Jalan Velox 23, Taman Industri Velox, 48000 Rawang, Selangor Darul Ehsan.

Tel: 016-216 6092 (HR) 016-218 1197 (ACC) Email: inquiry@weissen.com.my Web: www.weissen.com.my

	When handling or mixing, care should be taken to avoid direct contact with the skin and inhalation of any dust by wearing protective clothing, rubber gloves and a disposable half-mask respirator conforming to European Standard EN 149 or a non-disposable respirator to European Standard EN 140 with a filter to EN 143. Wash hands and exposed skin after handling medicated feedstuffs.
Interaction with other Medicaments	Fenbendazole should not be given concurrently with the bromsalan flukicides (dibromsalan, tribromsalan); abortions in cattle and death in sheep have been reported after using these compounds together.
Treatment on usage during pregnancy and lactation	Can be used during pregnancy and lactation. Administration to breeding and pregnant ruminants at 2 to 22 times the recommended dose has no apparent adverse effect. Fenbendazole is considered safe to use in pregnant bitches and is generally considered safe to use in pregnancy for all species. It is the drug of choice for treating giardia in pregnant animals (Tams 2007). In a system evaluating the safety of the drugs is categorized as in class: A (probably safe. Although specific studies may not have proved the safety of all drugs in dogs and cats, there are no reports of adverse effects in laboratory animal or women).
Adverse Effects/ undesirable Effects	At usual doses, fenbendazole generally does not cause any side effects. Side effects are very rare. Vomiting and diarrhea may occur.
Overdose and treatment	Fenbendazole is apparently well tolerated at doses up to 100 X recommended. The LD ₅₀ in laboratory animals exceeds 10 grams/kg when administered PO. It is unlikely an acute overdosage would lead to clinical signs.
Incompatibilities (for injection only)	N/A
Withdrawal period(s)	Meat and offal: 4 days.
Storage conditions	This product should be stored below 30°C (room temperature) in the closed container, dry, well-ventilated area and protect from sunlight.
Dosage form and packaging available	Dosage form: Powder Packaging available: 5 kgs



維山(馬)有限公司

WEISSEN COMPANY (M) SDN. BHD. (75684-M)

No.84, Jalan Velox 23, Taman Industri Velox, 48000 Rawang, Selangor Darul Ehsan.

Tel: 016-216 6092 (HR) 016-218 1197 (ACC) Email: inquiry@weissen.com.my Web: www.weissen.com.my

Name and address of manufacturer	Weissen Company (M) Sdn. Bhd. 15 & 17, Jalan Metro Perdana Barat 12, Kepong Sri Edaran Ind. Park, 52100 Kuala Lumpur, Malaysia.
Name and address of registration holder	Weissen Company (M) Sdn. Bhd. Lot 105, No. 84, Jalan Velox 23, Taman Velox, 48000 Rawang, Selangor, Malaysia.
Date of revision of package insert	02.10.2024
Shelf life (after first opening of container, where relevant)	None.
Shelf life of veterinary products as packaged for sales	24 months
Maximum Residual Limit (MRL)	Fenbendazole : All ruminant, Equidae (kidney, muscle, fat) is 50 µg/kg Porcine (Liver) is 500 µg/kg All ruminant (milk) is 10 µg/kg