

[Draft of Package Insert]

For Animal Use Only

TYLOSIN-100

Water Soluble Powder

CONTENT: Each sachet (120 g) contains,
Tylosin (as tartrate)100 g

PRODUCT DESCRIPTION: A light yellow powder.

PHARMACODYNAMICS:

Tylosin is a macrolide antibiotic produced by a strain of *Streptomyces fradiae*. It exerts its antimicrobial effect by inhibiting protein synthesis of susceptible micro-organisms. The tylosin spectrum of activity includes amongst others Gram-positive bacteria, some Gram – negative strains such as *Pasteurella*, and *Mycoplasma spp.*

PHARMACOKINETICS:

In most species peak plasma concentrations have been attained 1 to 2 hours after administration of tylosin. Compared to plasma levels clearly higher tissue concentrations have been observed. Tylosin was extensively metabolised.

Most of the residues are excreted in faeces predominantly consisting of tylosin (factor A), relomycin (factor D) and dihydrodesmycosin.

INDICATION:

Calves

- Treatment and prevention of pneumonia caused by *Mycoplasma spp* when the disease has been established in the herd.

Pigs

- Treatment and prevention of enzootic pneumonia caused by *Mycoplasma hyopneumoniae* and *Mycoplasma hyorhinis* when the disease has been established in the herd.

Chickens

- Treatment and prevention of chronic respiratory diseases (CRD) caused by *Mycoplasma gallisepticum* and *Mycoplasma synoviae* when the disease has been established in the flock.
- Treatment and prevention of necrotic enteritis caused by *Clostridium perfringens* when the disease has been established in the flock.

Turkeys

- Treatment and prevention of infectious sinusitis caused by *Mycoplasma gallisepticum*. When the disease has been established in the flock.

DOSAGE & ADMINISTRATION:

Oral administration through the drinking water.

In calves the product can also be administered through milk or milk replacer.

1 gram of the veterinary medicinal product corresponds to 0.833 gram of tylosin. The dosages are as follows:

Calves:

10 – 20 mg tylosin per kg BW (corresponding to 12 – 24 mg of the veterinary medicinal product per kg BW), twice daily (corresponding to a daily dose of 20 – 40 mg tylosin per kg BW), for 7 - 14 days.

Turkeys:

75 – 100 mg tylosin per kg BW per day (corresponding to 90 – 120 mg of the veterinary medicinal product per kg BW) for 3 – 5 days.

Chickens:

For the treatment of chronic respiratory disease:

75 – 100 mg tylosin per kg BW per day (corresponding to 90 - 120 mg of the veterinary medicinal product per kg BW) for 3 – 5 days.

For the treatment of necrotic enteritis:

20 mg tylosin per kg BW per day (corresponding to 24 mg of the veterinary medicinal product) for 3 days.

Pigs:

For the treatment of enzootic pneumonia:

20 mg tylosin per kg BW per day (corresponding to 24 mg of the veterinary medicinal product per kg BW) for 10 days.

For the preparation of the medicated water/milk/milk-replacer the body weight of the animals to be treated and their actual daily water/milk/milk-replacer consumption should be taken into due account. Consumption may vary depending on factors like age, state of health, breed, husbandry system. To provide the required amount of active substance in mg per litre drinking water/milk/milk-replacer the following calculation should be made:

$$\frac{\text{..... mg tylosin per kg bodyweight per day} \times \text{Average bodyweight (kg) of the animals to be treated}}{\text{Average amount of drinking water or milk / animal (l)}} = \frac{\text{.....mg tylosin}}{\text{l of drinking water}}$$

Sufficient access to the system of water supply should be available for the animals to be treated to ensure adequate water consumption. No other source of drinking water should be available during the medication period.

Should there be no clear response to treatment within 3 days the treatment approach should be reconsidered. After the end of the medication period the water supply system should be cleaned appropriately to avoid intake of sub-therapeutic amounts of the active substance which might support development of resistance.

Medicated water, milk or milk replacer should be replaced every 24 hours.

CONTRAINDICATIONS:

Do not leave or dispose of water containing tylosin tartrate where it may be accessible to either animals not under treatment or to wildlife.

WARNINGS / PRECAUTIONS:

Avoid direct contact with skin or conjunctivae.

DRUGS INTERACTIONS: None known.

PREGNANCY AND LACTATION:

Laboratory studies in mice and rats have not produced any evidence of teratogenic, foetotoxic or maternotoxic effects. No studies have been conducted in the target species. Use only according to the benefit/risk assessment by the responsible veterinarian.

ADVERSE EFFECTS:

If suspected adverse reactions occur, treatment should be discontinued and to inform veterinarian/local authority.

OVERDOSE AND TREATMENT:

Stop medication and to seek for veterinarian advice if there is sign of over dosage.

STORAGE CONDITIONS: Store below 30°C.

WITHDRAWAL PERIOD(S):

Calves (meat and offal): 12 days

Pigs (meat and offal): 1 day

Turkeys (meat and offal): 2 days

Turkey (eggs): Zero days

Chickens (meat and offal): 1 day

Chicken (eggs): Zero days

DOSAGE FORMS AND PACKAGING AVAILABLE: 120gm water soluble powder packed into aluminium sachet.

DATE OF REVISION OF PACKAGE INSERT: 12 February 2015.

Manufactured by:

PAHANG PHARMACY SDN. BHD. (10826-K)

LOT 5979, JALAN TERATAI,

5½ MILE OFF JALAN MERU, 41050 KLANG,

SELANGOR D.E., MALAYSIA.

Product Registration Holder:

GENPHARMA SDN. BHD. (268066-W)

LOT 5979, JALAN TERATAI,

5½ MILE OFF JALAN MERU, 41050 KLANG,

SELANGOR D.E., MALAYSIA.