

Biomutin 200 mg/ml Solution for Injection

Ingredients:

Each ml contains:

Tiamulin Hydrogen Fumarate 200 mg

Product Description:

Colourless, clear solution, acceptable slightly yellow tint. Solution for Injection.

Pharmacodynamics

The following organisms show sensitivity to tiamulin in vitro:

Brachyspira: *Brachyspira hyodysenteriae*, *Brachyspira pilosicoli*

Mycoplasmas: *Mycoplasma hyopneumoniae*, *M. hyorhinis*, *M. hyosynoviae*, *M. dispar*, *Ureaplasma spp.*, Gram-positive: *Staphylococcus spp.*, *Streptococcus spp.*, *Corynebacterium pyogenes*.

Gram-negative: *Pasteurella spp.*, *Klebsiella pneumoniae*, *Actinobacillus (Haemophilus)spp.*, *Fusobacterium necrophorum*, *Bacteroides spp.*, *Campylobacter coli*, *Lawsonia intracellularis*.

Pharmacokinetics

Following oral administration the active ingredient, tiamulin, is rapidly absorbed, peak serum levels being achieved within two hours. Tiamulin distributes widely in tissues, including the colon. High tiamulin concentrations are found in the lung of up to approximately 20 times the value found in the serum. Two hours post-injection, tiamulin lung levels of 14.5 – 16.7 mg/g have been found.

Indication(S):

Swine dysentery: For the treatment of swine dysentery caused by *Brachyspira hyodysenteriae* and complicated by *Bacteroides spp* and *Fusobacterium spp*.

Enzootic pneumonia complex of pigs: For the treatment of enzootic pneumonia complex, caused by *M. hyopneumoniae* and where the secondary bacteria are sensitive to tiamulin.

Mycoplasmal arthritis: For the treatment of arthritis caused by *Mycoplasma hyosynoviae*, to reduce lameness and restore growth performance.

Recommended Dosage and Mode of Administration

Solution for injections, intramuscular administration.

Swine dysentery Treatment: The dose is 1 ml/20 kg bodyweight (equivalent to 10 mg tiamulin hydrogen fumarate/kg body weight) intramuscularly administered as a single treatment to pigs showing clinical signs of the disease.

Enzootic Pneumonia complex of pigs treatment: The dose is 1.5 ml/20 kg bodyweight (equivalent to 15 mg of tiamulin hydrogen fumarate/kg body weight) administered intramuscularly for a period of 3 consecutive days to pigs showing clinical signs of the disease.

Mycoplasmal arthritis Treatment: The dose is 1.5 ml/20 kg bodyweight (equivalent to 15 mg tiamulin hydrogen fumarate/kg body weight) administered intramuscularly for a period of 3 consecutive days to pigs showing clinical signs of the disease.

Contraindications:

Not to be used in the case of over-sensitivity to tiamulin, in the case of bacterial resistance.

Adverse Reactions / Undesirable Effects

In rare cases, hypersensitivity to tiamulin is reported in terms of acute dermatitis with cutaneous erythema and intense pruritus. The adverse reactions are usually mild and transient but in very rare cases may be serious. Symptomatic treatment such as electrolyte therapy and an anti-inflammatory therapy may be useful.

Warnings and Precautions

Avoid self-injection. In the event of self-injection seek immediate medical advice. The product may cause local irritation, swelling and/ or pain reactions at the place of self-injection. In the event of the product coming into contact with skin, the affected area should be immediately rinsed under running water. In the event of the product coming into contact with eyes, it must be immediately rinsed under running water and medical advice must be sought.

Interactions with other medications

Do not combine Tiamulin with ionophore antibiotics (monensin, salinomycin, maduramicin, narasin, lasalocid) during or for at least seven days before and after treatment. Severe growth depression or death may result.

Statement On Usage During Pregnancy and Lactation

Can be used during pregnancy and lactation.

Overdose and Treatment

A single oral dose of 100 mg tiamulin/kg bodyweight caused hyperpnoea and abdominal discomfort in pigs. At a dose of 150 mg tiamulin/kg the only effect on the central nervous system was lethargy. A dose of 55 mg tiamulin/kg for 14 days caused increased salivation and a mild irritation of the stomach. Tiamulin hydrogen fumarate has a relatively high therapeutic index in pigs. The minimal lethal dose has not been established.

Withdrawal Period

Edible tissues - 21 days.

Incompatibilities

None known.

Storage Conditions

Store in a dry place at a temperature below 25°C. Protect from light.

Dosage Forms and Packaging Available

Solution for Injection. Packaging available: 100mL/unit



Product of YSP

Manufactured by:

The Bioveterinarian Industry Plant in Drwalew Joint Company

6 Grojecka Str., 05-651 Drwalew, Poland

Product Registration Holder:

Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)

Lot 3, 5, 7 & 12, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.

Ordering Line: 1 800 88 3027

Product Info: 1 800 88 3679