

VMT 250 mg/ml

Oral Solution

(FOR VETERINARY USE ONLY)

COMPOSITION

Each 1000 ml contents:

Tilmicosin (as Phosphate)	250 g
Benzyl alcohol	10 ml

PRODUCT DESCRIPTION

This product is brownish color clear solution.

PHARMACODYNAMICS

Tilmicosin is a semi-synthetic antibiotic of the macrolide group, and is believed to affect protein synthesis. It has bacteriostatic action but at high concentrations it may be bactericidal.

This antibacterial activity is predominantly against Gram-positive microorganism with activity against certain gram-negative ones and Mycoplasma of a bovine, porcine, ovine and avian origin. In particular, its activity has been demonstrated against the following microorganism:

Chickens: Mycoplasma gallisepticum and Mycoplasma synoviae

Scientific evidence suggests that macrolides act synergistically with the host immune system. Macrolides appear to enhance phagocyte killing of bacteria. Tilmicosin has been shown to inhibit in vitro the replication of the Porcine Reproductive and Respiratory Syndrome virus in alveolar macrophages in a dose dependent fashion.

Cross-resistance between tilmicosin and other macrolides and lincomycin has been observed.

PHARMACOKINETICS

Whilst blood concentrations of tilmicosin are low, there is pH-dependent macrophage accumulation of tilmicosin in inflamed tissue.

Poultry: As early as 6 hours after oral administration of 75 mg tilmicosin/litre drinking water, the average active substance concentrations detected in lung and alveolar tissue were 0.63 µg/g and 0.30 µg/g respectively. 48 hours after the start of the treatment, the tilmicosin concentrations in lung and alveolar tissue were 2.3 µg/g and 3.29 µg/g respectively.

TARGET SPECIES

Chickens (except hens producing eggs for human consumption)

INDICATIONS

Chickens: For the treatment and prevention of respiratory disease in chicken flocks, associated with mycoplasma gallisepticum and Mycoplasma synoviae.

ADMINISTRATION & DOSAGE

Oral administration through drinking water.

Chickens (except hens producing eggs for human consumption):

To be included in the drinking water at a daily dose of 15-20 mg/kg bodyweight in chickens for 3 days, which may be achieved by the inclusion of 75 mg tilmicosin per litre (30 ml VMT 250 mg/ml Oral Solution per 100 litres of drinking water).

The uptake of medicated water/milk replacer depends on the clinical condition of the animals. In order to obtain the correct dosage, the concentration of VMT 250 mg/ml Oral Solution should be adjusted accordingly.

CONTRA-INDICATIONS

Do not use in cases of hypersensitivity to the active substance or in cases of known resistance to tilmicosin.

WITHDRAWAL PERIODS

Meat and offal of Chicken: 12 days

Not authorized for use in laying birds producing eggs for human consumption. Do not use within 14 days of onset of the laying.

WARNING AND PRECAUTIONS

Warning: The safety of tilmicosin has not been established in animal used for breeding purposes.

Precautions: VMT 250 must be diluted before administration

- Keep medicine out of reach of children.
- Jauhkan ubat dari kanak-kanak.

INTERACTION WITH OTHER MEDICAMENTS

Cross resistance between tilmicosin and other macrolide antibiotics and lincosamide has been observed.

Tilmicosin may reduce the antimicrobial activity of β -lactam antibiotics.

PREGNANCY AND LACTATION

The safety of tilmicosin has not been established in animals used for breeding purposes.

SIDE EFFECT

In very rare cases, a decrease in water intake has been observed.

OVERDOSE AND TREATMENT

No symptoms of overdose have been seen in chickens given drinking water containing levels of tilmicosin up to 375 mg/litre (equivalent to 75-100 mg/kg bodyweight or 5 times the recommended dose) for 5 days, daily treatment with 75 mg/litre (equivalent to the maximum recommended dose) for 10 days resulted in a decrease in faecal consistency.

SHELF LIFE: 36 months

SHELF LIFE after first opening: Keep tightly after open. Use within 3 months

SHELF LIFE after dilution: Finish treatment within 12 hours after dilution

STORAGE

Store in cool and dry room ($\leq 25^{\circ}\text{C}$). Strictly avoid direct sunlight and heat exposure.

PACKAGING

The product filled 1000 ml in PE plastic bottle, by automatic filling machine and sealing, and packed in carton box.

DATE OF REVISION: 14 Dec 2018

Manufactured & Distributed By:

SHENNONG ANIMAL HEALTH (M) SDN BHD

神農動物保健(馬)有限公司

No 2, 4, 6, 8 & 10, Jalan Industri USJ 1/19,

Taman Perindustrian USJ 1,

47610 Subang Jaya, Selangor.

Tel: 03-8011 4646

Fax: 03-8024 1166

