

**SAME SIZE ARTWORK**  
**80 mm x 150 mm**

**SEBCA SHAMPOO 2.5% W/V**  
(Selenium Sulfide 2.5% Topical Suspension USP)

**COMPOSITION**

Selenium Sulfide USP 2.5% w/v

**DESCRIPTION**

Bright orange colored smooth suspension, free from foreign matter.

**PHARMACODYNAMICS**

Selenium sulfide is an antifungal agent often used in shampoos for the treatment of dandruff and seborrheic dermatitis. Selenium sulfide is highly active in inhibiting the growth of *P. ovale*. It is also a proven cytostatic agent, slowing the growth of both hyperproliferative and normal cells in dandruff and seborrheic dermatitis.

Topical selenium sulfide may act by an antimitotic action, resulting in a reduction in the turnover of epidermal cells. It also has antifungal activity. An antimitotic mechanism of action is suggested by data showing that selenium sulfide decreases the rate of incorporation of radioactively labeled thymidine into the DNA of dermal epithelial cells.

**PHARMACOKINETICS**

Selenium sulfide is not absorbed through normal skin, but can be absorbed through inflamed or damaged epithelium, leading to systemic toxicity. Selenium is then detectable in urine and plasma.

Urinary coproporphyrins and uroporphyrins may be elevated after systemic absorption. Selenium is excreted in humans in three elimination phases. Phase I lasts approximately 1 day, phase II lasts 20 days and phase III lasts 65-116 days. Selenium is able to bind to lipoprotein in plasma. Since selenium is an essential trace element it must cross the placenta but in physiological concentrations would cause no problems. No information is available on excretion in breast milk.

**Metabolism:**

Selenium sulfide is practically insoluble and can be rinsed off the skin. Following systemic absorption it is excreted mainly via the urine but also via the faeces.

Conversion to methyl selenite occurs, probably in the liver. Selenium is very rapidly oxidized to methyl selenite and none appears in the urine as selenium.

**INDICATIONS**

For the treatment of pityriasis capitis (dandruff), seborrheic dermatitis of the scalp and tinea versicolor.

**DOSAGE REGIMEN AND DIRECTION FOR USE**

For the treatment of dandruff and seborrheic dermatitis of the scalp: apply 5-10 ml to wet hair, lather and leave on the scalp for 3 minutes. Rinse and repeat application. The hair should be rinsed thoroughly after treatment and all traces of the lotion should be removed from the hands and nails. Use twice weekly for 2 weeks then once weekly for 2 weeks, and then only when necessary, or as directed by the physician.

For the treatment of tinea versicolor: apply the lotion undiluted to affected areas of the skin together with a small amount of water to form lather. Allow the lotion/lather to remain on the skin for 10 minutes, then rinse thoroughly. Repeat daily for 1 week or as directed by the physician.

**MODE OF ADMINISTRATION**

Topical

**CONTRAINDICATIONS**

Not to be used by patients allergic to any of its ingredients.

**PRECAUTIONS AND WARNING**

For external use only. Shake well before use. To minimize absorption Sebca (Selenium Sulfide 2.5%) should not be applied to mucous membranes, inflamed or broken skin, or to extensive areas of the skin. Contact with the eyes, genital area, should be avoided.

150 mm



BLACK

DATE: 27-10-2025  
VERSION-02

80 mm

**SAME SIZE ARTWORK**  
**80 mm x 150 mm**

Sebca (Selenium Sulfide 2.5%) shampoos should not be used within 48 hours of applying hair colours or permanent waving preparations. Remove ornaments before use.

**INTERACTIONS WITH OTHER MEDICAMENTS**

Your doctor or pharmacist may already be aware of any possible drug interactions and may be monitoring you for them. Do not start, stop, or change the dosage of any medicine before checking with your doctor or pharmacist first. Before using this medication, tell your doctor or pharmacist of all prescription and nonprescription/herbal products you may use. Keep a list of all your medications with you, and share the list with your doctor and pharmacist.

**PREGNANCY AND LACTATION**

There are no adequate data from the use of Selenium Sulfide in pregnant women. Animal studies are insufficient with respect to effects on pregnancy, foetal development and postnatal development. Hence avoid SEBCA Shampoo (Selenium Sulfide Topical Suspension USP 2.5%) during pregnancy & breastfeeding.

If required to be used during pregnancy & breastfeeding, consult doctor and use it as advised by doctor.

**SIDE EFFECTS**

Topical application of Sebca (Selenium Sulfide 2.5%) can produce irritation of the conjunctiva, scalp and skin, especially in the genital area and skin folds. Oiliness or dryness of scalp or hair, hair discoloration and hair loss have been reported. Sebca (Selenium Sulfide 2.5%) can be highly toxic when taken by mouth. Only traces of Sebca (Selenium Sulfide 2.5%) are absorbed through intact skin but prolonged use on broken skin has resulted in systemic toxicity.

Treatment of poisoning is symptomatic.

**SYMPTOMS AND TREATMENT OF OVERDOSE**

Selenium sulphide is highly toxic when taken by mouth. Symptoms of poisoning include an odour of garlic in the breath, a metallic taste, anorexia, vomiting. Treatment of poisoning is symptomatic.

**STORAGE CONDITION**

Store below 30°C in dry place.

**PACKAGING AND PRESENTATION**

Sebca (Selenium Sulfide 2.5%) is available in 100ml packs.

**PRODUCT REGISTRATION HOLDER**

Apex Pharmacy Marketing Sdn Bhd  
No. 2, Jalan SS13/5, 47500 Subang Jaya, Selangor, Malaysia.

**UNDER LICENCE FROM AND MANUFACTURED FOR:**

ZENTA HEALTHCARE PRIVATE LIMITED  
C 78, DDASHED,  
OKHLA INDUSTRIAL AREA PHASE 1,  
NEW DELHI 110 020, INDIA.

**MANUFACTURER**

Galentic Pharma (India) Pvt. Ltd.  
R-673, MIDC, TTC, Rabale,  
Thane-Belapur Road, Navi Mumbai, India.  
e mail: info@galentic.com

**DATE OF REVISION**

November 2025

ML 0188 PI-0

80 mm

150 mm