

[Package insert information will be printed on the product label]

PACKAGE INSERTS

PRODUCT NAME: HINAFLOR 300g/kg Granules

COMPOSITION: Each 100g premix contains FLORFENICOL 30g.

PRODUCT DESCRIPTION: White or almost white granules.

PHARMACODYNAMIC:

Florfenicol is a broad-spectrum synthetic antibiotic in the phenicol group that is active against most Gram-positive and Gram-negative bacteria isolated from domestic animals. Florfenicol acts by inhibition of protein synthesis at the ribosomal level and is bacteriostatic.

PHARMACOKINETIC:

Administration to pigs by gavage at 10 mg/kg under experimental conditions, absorption of florfenicol was variable but peak serum concentrations of approximately 5µg/ml were reached approximately 3 hours after dosing. The terminal half-life was between 3 and 4 hours. When pigs were given free access, for 5 days, to feed medicated with florfenicol (premix for medicated feeding stuff) at the recommended dose of 10 mg/kg serum florfenicol concentrations exceeded 1 µg/ml for more than 16 hours each day of treatment.

Florfenicol is well absorbed when administered orally and following distribution it is rapidly eliminated in the urine and faeces in a ratio of 3:1. A fraction is excreted unchanged and the rest is metabolised into 5 major metabolites. After parenteral dosing of florfenicol to pigs, it has been shown that lung concentrations are similar to serum concentrations. After a single dose of 10 mg florfenicol/kg b.w. mixed with feed to fasted pigs, maximum plasma concentration of approximately 7.4 µg/ml was reached up to 1.0 hour after dosing. The terminal half-life was approximately 2.8 hours.

INDICATION:

Pigs (Fattening pigs): For the treatment and prevention of swine respiratory disease caused by *Pasteurella multocida* susceptible to florfenicol in infected herds. The presence of the disease should be established in the herd before initiating preventive treatment.

USAGE & DOSAGE:

Mix with the feed basing on effective ingredient Florfenicol.

Dosage: 10 mg of florfenicol per kg body weight (bw) (equivalent to 33.33 mg HINAFLOR 300g/kg Granules) per day administered for 5 consecutive days. For a daily feed intake of 50 g/kg bodyweight, this dosage corresponds to a rate of incorporation of 0.667 kg of HINAFLOR 300g/kg Granules per ton of feed, i.e. 200 ppm of florfenicol. In all cases the recommended dose of 10 mg of florfenicol per kg of body weight per day, for 5 consecutive days has to be respected.

CONTRAINDICATIONS:

Do not administer to boars intended for breeding.

Do not use in case of hypersensitivity to the active substance or any of the excipients.

WARNING & PRECAUTIONS:

Animals showing a decreased appetite and/or a poor general condition should be treated by the parenteral route.

Special precautions to be taken by the person administering the medicinal product to animals.

Skin sensitisation may occur.

Avoid skin contact.

Special precautions for use in animals

Do not handle this product in case of known sensitization to active substance or any of the excipients.

The product should be used in conjunction with susceptibility testing and take into account official and local policy relating to the use of antimicrobials. - This premix is intended for the manufacturing of solid medicated feed and cannot be used as is; the incorporation rate of the premix in feed cannot be lower than 0.667kg/tonne. - Treatment should not exceed 5 days

DRUGS INTERACTIONS: None Known

PREGNANCY AND LACTATION:

The safety of the product during pregnancy and lactation has not been studied in sows. Use of the product during pregnancy and lactation is therefore not recommended.

ADVERSE EFFECTS:

Commonly observed adverse effects are diarrhoea perianal inflammation and rectal eversion. Increased serum calcium may also be observed. These effects are transient, resolving on cessation of treatment.

OVERDOSE AND TREATMENT:

In the event of overdose, a reduction in food and water consumption, together with a decrease in bodyweight may be observed. There may be an increase in refused feed.

WITHDRAWAL PERIODS: Meat and offal: 14 days

DISPOSAL OF CONTAINER: Unused material and packaging should be disposed of in accordance with local and national legislation

STORAGE CONDITIONS: Keep sealed and stored in cool and dry place below 30°C

DOSAGE FORMS AND PACKAGING AVAILABLE: 25kg/drum

MANUFACTURED BY:

HINAPHARM PHARMACEUTICAL CO., LTD. FOSHAN

Add: Shashuihe West Road, Cangjiang industrial zone, gaoming district, Foshan, Guangdong.

Office Add: Room 910&911, Tian An Hi-Tech Innovation Plaza, Ecological Hi-Tech Park, Yinbing Road, Panyu District, Guangzhou, Guangdong, 511400, P.R.China

PRODUCT REGISTRATION HOLDER:

PAHANG PHARMACY SDN. BHD. (10826-K)

Add: Lot 5979 Jalan Teratai, 5 1/2 Mile Off Jalan Meru, 41050 Klang, Selangor D.E., Malaysia.

DATE OF REVISION: 06 June 2015.