

VETFLOR 100mg/ml Oral Solution

FOR ANIMAL USE ONLY

Brand or Product Name :

VETFLOR 100mg/ml Oral Solution.

Name and Strength of Active Substance(s)

Each ml contains 100 mg florfenicol and 10 mg benzyl alcohol (as preservative).

Product Description

VETFLOR 100mg/ml Oral Solution is a light yellowish liquid with characteristic odour and lightly sweet taste.

Pharmacodynamics

Florfenicol is a broad spectrum antibiotic that has activity against many bacteria. It acts by binding to the 50S ribosome, thereby inhibiting bacterial protein synthesis. Florfenicol has a wide spectrum of activity against many Gram-positive and negative bacteria. Susceptible bacteria include *Haemophilus* spp., *Pasteurella* spp., *Actinobacillus pleuropneumoniae*, and *Streptococcus* spp.

Pharmacokinetics

Broiler chicken : After the first oral gavage administration of florfenicol of 42mg/kg bw/day for 3 days to broiler chicken, the highest concentration of florfenicol in serum was measured after 30 minutes then declined to reach 810 µg/ml, 8 hours later. Florfenicol could not be detected in the serum, 60 hours after the last administration. During a continuous treatment of broiler chicken with florfenicol in drinking water at a daily dose of 26mg/kg bw for 18 hours for 3 days, the estimated mean serum concentration of florfenicol was 711 µg/ml. No florfenicol was detected in the serum, 72 hours after the terminal dose. After repeated oral administration of ¹⁴C-florfenicol, given twice daily 12 hours apart at 20mg/kg bw/dose for three days, 93.7% and 98.2% of total radioactivity administered were excreted within 1 day and 7 days after the last dose. In excreta, at 7 days, the parent compound represented the major fraction of the radioactivity (42%), florfenicol amine (25%), florfenicol oxamic acid (5%) and florfenicol alcohol (10%) and the remaining part of radioactivity being represented by a small percentage of three unknown compounds. No monochloroflorfenicol was detected in excreta. In a radiometric study, chicken received ¹⁴C-florfenicol by oral gavage twice daily 12 hour apart at 20 mg/kg bw/dose for three days (i.e. 40 mg/kg bw/day for three days). Twenty-four hour after the end of oral administration, the levels of radioactivity in edible tissues were 146 µg equivalents florfenicol/kg in muscle, 475 µg/kg in skin+ fat, 11148 µg/kg in liver and 3125 µg/kg in kidney. Then declined to reach approximately 50 µg/kg in muscle and in skin + fat 5 days after the end of treatment, whereas in liver and kidney significant amounts of residues were still found : 2403 and 844 µg equivalents florfenicol/kg respectively. Florfenicol amine was the major metabolite measured in edible tissues of chicken, florfenicol, the only microbiologically active compounds, being only detected in skin + fat in low concentrations (4%) and in kidney (1%). In a non-radiometric study, broiler chickens received florfenicol according to the recommended regimen via drinking water at concentrations equivalent to 17 to 30 mg/kg bw/day for 3 days. At 0.5 days after the end of the treatment the concentrations of florfenicol amine were lower than the limit of quantification for muscle (less than 50 µg/kg) and for skin + fat (less than 109 µg/kg) whereas in liver and kidney, significant amounts of florfenicol-amine were measured 2862 and 1161 µg/kg respectively. Florfenicol amine concentrations declined to reach 2038 and 679 µg/kg in liver and kidney, 1 day after the end of the treatment and to 1215 and 484 µg/kg in liver and kidney, 3 days after the end of the treatment. Seven days after the end of the treatment, the concentrations of florfenicol amine were below the limit of quantification for liver (less than 461 µg/kg) and could be still measured in kidney (136 µg/kg).

Pigs : After administration to pigs by gavage at 15 mg/kg under experimental conditions, absorption of florfenicol was variable but peak serum concentrations of approximately 5 µg/mL were reached approximately 2 hours after dosing. The terminal half-life was between 2 and 3 hours. When pigs were given free access, for 5 days, to water medicated at a concentration of 100 mg florfenicol per litre of water, serum concentrations of florfenicol exceeded 1 µg/mL for the entire 5 day treatment period except for a couple of short excursions below 1 µg/mL. After absorption and distribution, florfenicol is extensively metabolised by pigs and rapidly eliminated, primarily in urine. After parenteral dosing of florfenicol to pigs, it has been shown that lung concentrations are similar to serum concentrations.

Indication

Broiler chickens : Treatment and control of air sacculitis associated with *E. coli* susceptible to florfenicol.

Swine: Treatment and prevention of swine respiratory disease associated with *Actinobacillus pleuropneumoniae* and *Pasteurella multocida* susceptible to florfenicol. The presence of the disease should be established in the herd before initiating preventive treatment.

Recommended Dose

To be administered via animal drinking water.

Broiler chicken : 1 ml per L drinking water daily for 3 consecutive days. This dosage is equivalent to 100 mg florfenicol per L or 35 mg florfenicol per kg body weight.

Swine : 1 ml per L drinking water daily for 5 consecutive days. This dosage is equivalent to 100 mg florfenicol per L or 10 mg florfenicol per kg bodyweight.

Administration / Direction of Use

The medicated water should be administered as the only source of drinking water. The appropriate quantity of medicated water or pre-diluted medicated water should be prepared based on the daily water consumption. Medication should be initiated promptly when disease is diagnosed. Prepare fresh stock solution and medicated water daily. If a positive clinical response is not noted within 48 hours of initiation of treatment, the diagnosis should be reevaluated.

Contraindications

Do not use in boars or birds intended for breeding purposes.

Do not administer in cases of previous allergic reactions to florfenicol.

Warnings and Precautions

Warnings :

Inappropriate use of the veterinary medicinal product may increase the prevalence of bacteria resistant to florfenicol. To limit the development of antimicrobial resistance, the choice of this drug as the most appropriate treatment should be confirmed by clinical experience supported where possible by pathogen culture and drug susceptibility testing. Not for human use.

Special warnings for each target species

None.

Special precautions for use

i. Special precautions for use in animals

The veterinary medicinal product should be used in conjunction with susceptibility testing. Inappropriate use of the veterinary medicinal product may increase the prevalence of bacteria resistant to florfenicol. Treatment should not exceed the recommended days.

ii. Special precautions to be taken by the person administering the veterinary medicinal product to animals

Avoid direct contact with skin, eyes and clothes. In case of accidental skin exposure, wash with soap and water. Remove contaminated clothing. Consult a physician if irritation persists.

ii. Other precautions

Manure from treated pigs should be stored for one month prior to spreading and incorporating into land

Interactions with Other Medicaments

There is inadequate data on adverse drug interactions produced by florfenicol. In some species, including the horse, florfenicol may be anticipated to inhibit hepatic microsomal enzymes and therefore like chloramphenicol to interact adversely with drugs metabolized in the liver, but details are lacking.

Pregnancy and lactation

The effects of florfenicol on treated porcine and chickens reproductive performance, pregnancy and lactation have not been assessed. The use is not recommended during pregnancy and lactation.

Adverse Effects

Chicken : None known. Pig : A slight reduction of water consumption by the animals, dark brown faeces and constipation may be observed during treatment. Commonly observed adverse effects are diarrhoea and/or peri-anal and rectal erythema/oedema which may affect approximately 40% of the animals. These effects are transient. In a few of the affected animals, prolapse of the rectum, that resolves without treatment, may be observed.

Overdose and Treatment

Chicken : None known.

Pig : In case of overdosing, a decrease in weight gain, food and water consumption, peri-anal erythema and oedema and modification of some haematological and biochemical parameters indicative of dehydration may be observed

Withdrawal Period(s)

Swine : 21 days, Broiler chicken: 5 days

Shelf-life

2 years from date of manufacture. Opened bottle and unused medicated water to be discarded after 24 hours.

Maximum Residual Limit (MRL)

Poultry : Muscle = 100 µg/kg, Skin & fat = 200 µg/kg, Liver = 2500 µg/kg, Kidney = 750 µg/kg. Not for use in animals from which eggs are produced for human consumption. Porcine : Muscle = 300 µg/kg, Skin + fat = 500 µg/kg, Liver = 2000 µg/kg, Kidney = 500 µg/kg [Ref. EMA]

Storage Conditions

Store in a dry place below 25°C. Jauhi ubat dari kanak-kanak. Opened bottle and unused medicated water to be discarded after 24 hours. Do not use/ store this product in galvanized metal watering systems or containers. Do not operate chlorinators while administering medication.

Disposal of Containers

Empty bottle must be safely disposed in trash according to local government regulations.

Dosage Forms and Packaging Available

1 litre in HDPE bottle, pack in 12 bottles or 20 bottles per carton.

Name and Address of Manufacturer

Life Biopharma Sdn. Bhd. (546163-D)

No. 250-251, Jalan Industri Galla 13, Kawasan Perindustrian Galla, 70200 Seremban, Negeri Sembilan, Malaysia.

Date of Revision of Package Insert

2015-02-10