

Aviax 50 gram/kg powder

BROAD SPECTRUM ANTICOCIDIAL PREMIX

FORMULA:

Each 100g of premix contains

Semduramicin.....	5g
Excipients up to	100g

PRODUCT DESCRIPTION:

It is a free flowing, uniform, tan to brown powder, free of foreign matter and agglomerates.

PHARMACODYNAMICS:

Semduramicin has a high affinity for the membranes of eimeria species, thus facilitating the transfer of cation across the membrane into the cytoplasm. This process will modifies inside the cell, this will rupture and disintegrate the coccidial cell wall.

PHARMACOKINETICS:

Unchanged Semduramicin Sodium is the major residue component in the chicken tissues. The liver is the organ where higher concentrations of semduramicin can be found.

ENVIRONMENTAL PROPERTIES:

Aviax when used as directed does not have any significant adverse effects on terrestrial organisms in the environment.

INDICATION:

For the prevention of coccidiosis caused by *Eimeria acervulina*, *E.brunetti*, *E. maxima*, *E. mivati*, *E. necatrix* and *E. tenella* sensitive to semduramicin.

RECOMMENDED DOSAGE:

Mix 500 g (0.5kg) of this premix per metric ton (1000kg) of feed, to provide 25ppm of semduramicin in the finished feed.

MODE OF ADMINISTRATION:

Route of Administration: ORAL

Use continuously as the sole ration in finished feed to broilers chickens.

CONTRAINDICATIONS:

There are no specific contraindications to semduramicin when used as directed in the feed of broiler chickens. This product is not intended for treatment of coccidiosis.

WARNINGS:

Handle with care. Wash hands thoroughly after preparation. Operators should wear protective clothing and mask during feed mixing.

PRECAUTIONS:

- For use in broiler chickens.
- Do not exceed recommended dosage.
- Ensure that feeds containing Aviax are not accidentally administered to animals other than broiler birds.

INTERACTIONS WITH OTHER MEDICAMENTS:

There are no known physico-chemical interactions which could restrict the use of this product.

STATEMENT ON USAGE DURING PREGNANCY AND LACTATION:

Not for use in laying chickens.

ADVERSE EFFECTS / UNDESIRABLE EFFECTS:

Semduramicin is well tolerated by broiler chickens but when used in excessive levels a drop in feed intake and weight depression will be observed.

OVERDOSE AND TREATMENT:

None known. If overdose is confirmed, remove the feed containing semduramicin and provide proper access to drinking water to the birds.

WITHDRAWAL PERIOD(S):

The withdrawal period in broilers is 0 (zero) days.

STORAGE CONDITIONS:

Store in a cool and dry place below 30°C.

DOSAGE FORMS AND PACKAGING AVAILABLE:

Medicated powder premix
25 kg bag

Manufactured by Phibro Saúde Animal Internacional LTDA - BRAZIL - Reg. MAPA 016/2000
Av. Presidente Tancredo de A. Neves, n° 1063, 07112-070, Guarulhos- SP- BRAZIL

Registration holder: Phibro Corporation (M) Sdn Bhd
311, Level 3, Menara Axis,
2, Jalan 51A/223, Seksyen 51A,
46100 Petaling Jaya, Selangor, Malaysia

Revision Date of Package Insert: Aug 15, 2025