

Apramycin changed to Apramycin

Veterinary Use Only

Kapramycin Apramycin 33.3% Soluble Powder

1. Brand or Product Name : Kapramycin Apramycin 33.3 Soluble Powder
2. Name and Strength of Active Substance(s)
Each kg contains : Apramycin (as Apramycin sulfate) ----- 333g
3. Product Description : Brown or pale brown granular powder
4. Pharmacodynamics/Pharmacokinetics/Environmental Properties
Pharmacodynamics :
Apramycin is bactericidal. It also acts against bacteria by inhibiting protein synthesis at the ribosome level. Like the aminoglycosides, it inhibits the translocation step of protein synthesis and induces translation errors.
Pharmacokinetics :
Absorption
The oral administration of apramycin is intended for antimicrobial activity within the gut; apramycin is poorly absorbed, particularly in older animals: Absorption: Pigs, Nine mg of apramycin/kg bodyweight administered orally is well absorbed in 2-day old pigs, slightly absorbed in 4-week old pigs and not absorbed in 8-week old pigs.
Distribution, Biotransformation and Excretion
Pigs. Very little metabolism of apramycin takes place in the animal. Dosing 10kg pigs with 14C apramycin resulted in approximately 83% being recovered from the faeces, and 4% from the urine, as 14C apramycin.
5. Indication
Pigs : For the treatment of bacterial enteritis caused by susceptible strains of *Escherichia coli* and other organisms susceptible to apramycin
Broiler : For use in the control and treatment of colisepticaemia (colibacillosis) in broilers caused by susceptible strains of *E.coli*
6. Recommended Dosage / Mode of Administration
Oral : Via drinking water
1)Pigs
Neonatal: Pig should receive 25mg apramycin activity per kg of body weight daily.
Start treatment on the day of birth and continue once daily for two additional days.
Weanling: For treatment of colibacillosis, pigs should consume sufficient medicated drinking water to obtain 12.5mg of apramycin activity per kg of body weight daily. Continue treatment for seven days. Prepare the fresh medicated solution daily to assure activity.
2)Broilers
250 to 500mg apramycin activity/litre of drinking water (0.5 to 1 packs of Kapramycin 33.3% per 200 litres) for 5 consecutive days. This is equivalent to an average of 40-80mg apramycin activity per kg body weight daily.
7. Contraindications
- Do not use in cases of known hypersensitivity to apramycin.
- Do not use in cats.

8. Warnings and Precautions
- Do not administer to cat.
- Read full directions before use.
- Do not exceed the recommended dosage.
- Keep the withdrawal period.
- Put on the protected wears and devices such as gloves and mask in order to client's safety.
- Use by veterinarian's instruction, and veterinary use only.
- Keep out of reach of children/ Jauhi ubat daripada kanak-kanak.
9. Interactions with Other Medicaments
This veterinary medicinal product must not be mixed with other products.
10. Statement on usage during pregnancy and lactation
This veterinary medicinal product must not be used during pregnancy and lactation. Do not use in laying birds producing eggs for human consumption.
11. Adverse Effects/Undesirable Effects
If suspected adverse reactions occur, treatment should be discontinued and to inform veterinarian/ local authority.
12. Overdose and Treatment
Stop medication and to seek for veterinarian advice if there is sign of overdosage.
13. Withdrawal Period(s)
Meat - Do not use less than 14 days before slaughter for human consumption.
14. Shelf life (after reconstitution or dilution)
24 hours
15. Storage
Store in a dry and cool place below 30°C.
Protect from sunlight. Store in well-tight container.
16. Dose Forms and packaging available
300g
17. Name and Address of manufacturer/marketing authorization holder
Nutri Pharmax Sdn Bhd
6, Lorong Serunai 1, Valdor Sungai Bakap, 14200 Sungai Jawi, Pulau Pinang, Malaysia.
18. Date of Revision of Package Insert : 12 JUL 2019

Distributed by
NP Nutri Pharmax Sdn.Bhd.
6, Lorong Serunai 1, Valdor Sungai Bakap,
14200 Sungai Jawi, Pulau Pinang, Malaysia
Tel: +604-582 9377

Manufactured by
KBNP, INC.
235-9 Chusaro, Sinam, Yesan,
Chungnam, Korea

4063015
190717

The address is changed.