

SULFACOX 50% POWDER

Description and Composition

SULFACOX 50% POWDER is a light yellowish beige water soluble powder. Each kg contains 500g Sulfachloropyrazine Sodium monohydrate.

Pharmacodynamics

Sulfachloropyrazine sodium is an antibiotic belonging to the group of sulfonamides. Like all sulfonamides, Sulfachloropyrazine is a competitive antagonist of para-aminobenzoic acid (PABA), a precursor of folic acid, in protozoa and bacteria. Folic acid is a coenzyme necessary for the synthesis of nucleic acid, Sulfonamides interfere with the biosynthesis of folic acid in bacterial cells; they compete with para-aminobenzoic acid (PABA) for incorporation in the folic acid molecule. By replacing the PABA molecule and preventing the folic acid formation required for DNA synthesis, the sulfonamides prevent multiplication of the bacterial cell. Hence sensitive species do not multiply in the presence of Sulfachloropyrazine sodium. Only organisms that synthesize their own folic acid are susceptible; mammalian cells use preformed folic acid and, therefore, are not susceptible. Cells that produce excess PABA or environments with PABA, such as necrotic tissues, allow for resistance by competition with the sulfonamide.

Pharmacokinetics

Most sulfonamides are well absorbed orally. Delays in absorption may occur in adult ruminants or when sulfonamides are administered with food to monogastric animals. Sulfonamides are widely distributed throughout the body. They cross the placenta, and a few penetrate into the cerebrospinal fluid. Sulfonamides are primarily metabolized in the liver but metabolism also occurs in other tissues.

Indication

Indicated for the treatment of coccidiosis.

Recommended Dose

0.03% in drinking water. Administer in drinking water for 3 days as sole source of drinking water and sulphonamide medication.

Route of Administration

For oral administration in drinking water.

Contraindication

None known

Warning and Precautions

For use in drinking water of poultry. Prepare fresh water daily.

Avoid underdosing. Sick birds may drink less water than normal. When such birds can be isolated the proportion of SULFACOX 50% POWDER in the water can be proportionally increased.

Avoid overdosing. At high ambient temperatures birds drink more water than usual, and thus the concentration of SULFACOX 50% POWDER in the water should be decreased.

Interaction with Other Medicinal Product

Drug interactions relating specifically to the use of sulfonamides in animals are rarely reported in veterinary literature.

Pregnancy and Lactation

Not to be administered to chickens producing eggs for human consumption

Side Effects

Crystallization in the urinary tract. Note: *Crystallization* of sulfonamides can occur in the kidneys or urine with high doses of sulfonamide or when an animal is dehydrated.

Symptoms and Treatment of Overdose

Toxicities secondary to acute overdose of sulfonamides are not typically reported.

Storage Condition

Store below 30°C

Shelf life

3 years. To be used within 24 hours after dilution.

Withdrawal Period

Meat: 4 days

Not to administered to chickens producing eggs for human consumption

Packing

100g, 108g, 200g, 500g, 1kg and 25kg

REVISED: 5.10.2023

	Manufactured by: Range Pharma Sdn Bhd 1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 E-mail: info@rangepharma.com
---	--