

Le-Stres 800 mg/g Water Soluble Powder

Le-Stres 800 mg/g Water Soluble Powder appears to be a pink color flakes.

Composition:

Each g contains

Sodium Salicylate 800 mg

Pharmacodynamics:

Sodium salicylate is a non-steroidal anti-inflammatory drug (NSAID) and exerts an anti-inflammatory, analgesic and anti-pyretic effect. The effects are linked to the inhibition of the enzyme cyclo-oxygenase by which the synthesis of prostaglandin (mediator for inflammation) decreases.

Pharmacokinetics:

Orally ingested salicylates are absorbed rapidly by passive diffusion, partly from the stomach but mostly from the upper small intestine.

After absorption, salicylate is distributed throughout most body tissues. Values of volume of distribution (Vd) are higher in the newborns. Half-lives are longer in the very young resulting in slower elimination of the substance. This is most prominent in animals up to 7-14 days of age.

The metabolism of salicylate takes mainly place in hepatic endoplasmic reticulum and mitochondria.

Excretion is mainly via the urine and is a pH-dependent process.

Indications:

Calves: For supportive treatment of pyrexia in acute respiratory disease, in combination with appropriate (anti-infective) therapy if necessary.

Pigs: For the treatment of inflammation, in combination with concurrent antibiotic therapy.

Directions for Use:

Calves: 40 mg sodium salicylate per kg of bodyweight once daily, (equivalent to 50 mg product per kg BW per day), for 1 - 3 days.

Pigs: 35 mg sodium salicylate per kg of bodyweight per day, (equivalent to 43.75 mg product per kg BW per day), for 3-5 days.

The product can be administered orally through the milk-replacer and/or the drinking water.

Mode of Administration: Oral Administration

Contraindication:

Do not administer to animals with severe hypoproteinaemia, liver and kidney affections.

Do not administer in case of gastrointestinal ulcerations and chronic gastrointestinal disorders.

Do not administer in case of malfunction of the hematopoietic system, coagulopathies, haemorrhagic diathesis.

Do not use sodium salicylates in neonates or calves less than 2 weeks of age.

Do not use in piglets less than 4 weeks of age.

Do not use in animals with known hypersensitivity to sodium salicylate or to excipient.

Warning and Precautions:

Special warnings for each target species

None.

Special precautions for use

i. Special precautions for use in animals

Given that sodium salicylate may inhibit clotting of blood, it is recommended that elective surgery should not be performed on animals within 7 days after the end of treatment.

ii. Special precautions to be taken by the person administering the veterinary medicinal product to animals

People with known hypersensitivity to sodium salicylate or related substances (e.g. aspirin) or excipients should avoid contact with the veterinary medicinal product.

Irritation of the skin, eye, and respiratory tract might occur. During preparation and mixing of the product, direct contact with the skin and eyes, and direct inhalation of the powder should be avoided. It is recommended to wear gloves, safety glasses, and a dust mask. Particular attention should be taken when opening the bucket.

In case of accidental dermal exposure wash skin immediately with water.

In the event of accidental eye contact, the user is advised to wash the eye with plenty of water for 15 minutes, and seek medical advice if irritation persists.

During administration of medicated drinking water or milk (replacer) to the animals skin contact should be prevented by wearing gloves. Wash accidentally exposed skin immediately with water.

Interaction with Other Medicines:

Concurrent administration of potentially nephrotoxic drugs (e.g. aminoglycosides) should be avoided.

Salicylic acid is highly plasma (albumin) bound and competes with a variety of compounds (e.g. ketoprofen) for plasma protein binding sites

Plasma clearance of salicylic acid has been reported to increase in combination with corticosteroids, possibly due to induction of metabolism of salicylic acid.

Concurrent use with other NSAIDs is not recommended, because of increased risk of gastro-intestinal ulcerations.

Do not use in combination with drugs known to have anticoagulant properties.

Statement on Usage During Pregnancy and Lactation:

Laboratory studies in rats have shown evidence of teratogenic and foetotoxic effects.

Salicylic acid crosses the placenta and is excreted with the milk. Half-life in the newborn is longer and thus toxicity symptoms may occur much sooner. Furthermore platelet aggregation is inhibited and bleeding time is increased, a situation which is not favourable during difficult parturition / caesarean section. Finally some studies indicate that delivery is postponed.

The product should not be used during pregnancy and lactation.

Adverse Effects / Undesirable Effects:

Gastrointestinal irritation may occur especially in animals with pre-existing gastrointestinal disease. Such irritation may clinically be manifested by production of black manure due to blood loss in the gastrointestinal tract.

Inhibition of normal blood clotting may occur incidentally. If this effect occurs it will be reversible and effects will diminish within approximately 7 days.

Overdose and Treatment::

Calves tolerate dosages up to 80 mg/kg for 5 days or 40 mg/kg for 10 days without any adverse effects.

Pigs tolerate dosages up to 175 mg/kg for up to 10 days without any significant adverse effects.

In case of an acute overdose intravenous bicarbonate infusion results in a higher clearance of salicylic acid by alkalinisation of the urine and may be beneficial in correcting (secondary metabolic) acidosis.

Withdrawal Period (s):

Calves and pigs:

Meat and offal: zero days.

Storage Condition:

Store below 30 °C. Protect from heat and sunlight exposure.

Keep medicine out of reach of children

(Jauhkan ubat dari kanak-kanak).

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Shelf-life after opening: 7 days

Shelf life after reconstitution: 24 hours

Packaging Available: 100 g & 500 g

Date of revision: 7-4-2015

Name and Address of Manufacturer:

NAM PHARMA SDN. BHD.

5, Lebuhr Perusahan Klebang 11,

Taman Perindustrian Antarabangsa IGB,

31200 Chemor, Perak, Malaysia.

Tel: +605-2931838, 1839 Fax: +605-2931833