

FLORCOL ORAL SOLUTION 250MG/ML

Ingredient(s):

Each ml contains:

Florfenicol 250mg

Pharmacodynamic:

Florfenicol is a broad-spectrum synthetic antibiotic in the phenicol group that is active against most Gram-positive and Gram-negative bacteria isolated from domestic animals. Florfenicol acts by inhibition of protein synthesis at the ribosomal level and is bacteriostatic. However, bactericidal activity has been demonstrated in-vitro against *Actinobacillus pleuropneumoniae* and *Pasteurella multocida* when florfenicol is present at concentrations above the MIC for up to 12 hours.

In-vitro testing has shown that florfenicol is active against the bacterial pathogens most commonly isolated in respiratory diseases in pigs, including *Actinobacillus pleuropneumoniae* and *Pasteurella multocida*.

Pharmacokinetic:

After administration to pigs by gavage at 15 mg/kg under experimental conditions, absorption of florfenicol was variable but peak serum concentrations of approximately 5 µg/mL were reached approximately 2 hours after dosing. The terminal half-life was between 2 and 3 hours. When pigs were given free access, for 5 days, to water medicated with florfenicol drinking water concentrate at a concentration of 100 mg florfenicol per litre of water, serum concentrations of florfenicol exceeded 1 µg/mL for the entire 5 day treatment period except for a couple of short excursions below 1 µg/mL.

After absorption and distribution, florfenicol is extensively metabolised by pigs and rapidly eliminated, primarily in urine.

After parenteral dosing of florfenicol to pigs, it has been shown that lung concentrations are similar to serum concentrations.

Indication:

Treatment and prevention at the group level where clinical signs are present of swine respiratory disease associated with *Actinobacillus pleuropneumoniae* and *Pasteurella multocida* susceptible to florfenicol. The presence of the disease should be established in the herd before initiating preventive treatment.

Target Species:

Swine.

Recommended Dose:

0.4ml of Florcol Oral Solution per 10 kg bodyweight per day. Use for 5 consecutive days.
(equivalent to 10 mg florfenicol per kg bodyweight per day).

The appropriate quantity of medicated water or pre-diluted medicated water should be prepared based on the daily water consumption.

Mode of Administration:

Oral. For use in swine drinking water.

Contraindication(s):

Do not use in boars intended for breeding purposes.

Do not administer in cases of previous allergic reactions to florfenicol.

Warning and Precautions:

To be used by or on the order of a licensed veterinarian.

Inappropriate use of the veterinary medicinal product may increase the prevalence of bacteria resistant to florfenicol.

Treatment should not exceed 5 days.

Interaction with Other Medicaments:

No data available.

1. To extend the shelf life from 2 years to 3 years.
2. To include the phrase "Made in Vietnam".
3. To revise the term "Marketing Authorization Holder" to "Product Registration Holder".
4. To standardize YSP Malaysia address with the Quest3+ system.
5. The variation for Change of Site (COS) from YSP Malaysia to YSP Vietnam was approved on 18 May 2016. However, the approved package insert in the Quest3+ system still mentioned the old manufacturing site (system error). Therefore, we have updated the manufacturer information in the package insert accordingly.
6. To update the date of revision.

Pregnancy and Lactation:

Studies in laboratory animals have not revealed any evidence of potential embryotoxic or foetotoxic effect of florfenicol.

The safety of the veterinary medicinal product in sows has not been established during pregnancy and lactation.

The use is not recommended during pregnancy and lactation.

Side Effect(s):

A slight reduction of water consumption by the animals, dark brown faeces and constipation may be observed during treatment.

Commonly observed adverse effects are diarrhoea and/or peri-anal and rectal erythema/oedema which may affect approximately 40% of the animals. These effects are transient. In a few of the affected animals, prolapse of the rectum, that resolves without treatment, may be observed.

Environmental Property:

Florfenicol is not readily biodegradable but there is evidence of inherent biodegradability.

Symptoms and Treatment for Overdosage, and Antidote(s):

In case of overdosing, a decrease in weight gain, food and water consumption, peri-anal erythema and oedema and modification of some haematological and biochemical parameters indicative of dehydration may be observed.

Storage Condition(s):

Store at temperature below 30°C. Protect from direct sun light and water.

Shelf-Life:

3 Years

Used within 24 hours after reconstitution.

Shelf life after first opening (3 years).

Withdrawal Periods

Swine: Meat and offal: 20 days

Maximum Residual Limit (MRL)

Swine:

Muscle - 300µg/kg

Skin & fat - 500µg/kg

Liver - 2000µg/kg

Kidney - 500µg/kg

Product Description

A clear, colourless to pale yellowish solution.

Plastic bottle 1 liter.

Made in Vietnam

Manufacturer:

Y.S.P INDUSTRIES VIETNAM CO., LTD.

Road No. 6, Nhon Trach Industrial Zone 3, Nhon Trach Dist.,
Dong Nai Province, Vietnam Tel: +84251 3569 993

Product Registration Holder

Y.S.P INDUSTRIES (M) SDN. BHD. (199601001034)

Lot 3, 5, 7 & 12, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor Darul Ehsan Malaysia.

Ordering Line: 1 800 88 3027 Product Info: 1 800 88 3679



Y. S. P.

218028-01

Date of revision 09Apr24

L=260mm

L=260mm

W= 110mm

W= 110mm