

TAFLO[®]R 20% W/V ORAL SOLUTION

Name and Strength of Active Substance(s)

Florfenicol 20%w/v

Product Description

Colorless to pale yellowish solution.

Pharmacodynamics

Florfenicol is a broad-spectrum synthetic antibiotic in the phenicol group that is active against most Gram-positive and Gram-negative bacteria isolated from domestic animals. Florfenicol acts by inhibition of protein synthesis at the ribosomal level and is bacteriostatic. However, bactericidal activity has been demonstrated in-vitro against *Actinobacillus pleuropneumoniae* and *Pasteurella multocida* when florfenicol is present at concentrations above the MIC for up to 12 hours.

In-vitro testing has shown that florfenicol is active against the bacterial pathogens most commonly isolated in respiratory diseases in pigs, including *Actinobacillus pleuropneumoniae* and *Pasteurella multocida*.

Pharmacokinetics

Swine

After administration to pigs by gavage at 15mg/kg under experimental conditions, absorption of florfenicol was variable but peak serum concentrations of approximately 5µg/mL were reached approximately 2 hours after dosing. The terminal half-life was between 2 and 3 hours. When pigs were given free access, for 5 days, to water medicated with florfenicol at a concentration of 100mg florfenicol per litre of water, serum concentrations of florfenicol exceeded 1µg/mL for the entire 5 day treatment period except for a couple of short excursions below 1µg/mL.

After absorption and distribution, florfenicol is extensively metabolised by pigs and rapidly eliminated, primarily in urine.

After parenteral dosing of florfenicol to pigs, it has been shown that lung concentrations are similar to serum concentrations.

Chicken

During a continuous treatment of broiler chicken with florfenicol in drinking water at a daily dose of 26mg/kg bw for 18 hours for 3 days, the estimated mean serum concentration of florfenicol was 711µg/L. No florfenicol was detected in the serum, 72 hours after terminal dose.

The major metabolite was florfenicol amine measured in edible tissues of chicken, and florfenicol is only detected in skin + fat in low concentrations (4%) and in kidney (1%).

Environmental Properties Not relevant

Indication

SWINE: For the treatment of swine respiratory disease associated with *Actinobacillus pleuropneumoniae* and *Pasteurella multocida*.

BROILER CHICKENS: For the indication in broiler chickens for the treatment and control of air sacculitis associated with *E. coli* susceptible to florfenicol.

Recommended Dosage

Swine: 100ppm(100mg/L) for 5 consecutive days in drinking water. This dosage rate is equivalent to 10mg/L body weight for pigs that are drinking 10% of their body weight.

Chicken: 100ppm(100mg/L) for 3 consecutive days in drinking water. This dosage rate is equivalent to 35mg/kg body weight of birds.

Mode of Administration Orally

Contraindications

Do not use in boars intended for breeding purposes.

Do not administer in cases of previous allergic reactions to florfenicol

Warnings and Precautions

i . Special precautions for use in animals

The veterinary medicinal product should be used in conjunction with susceptibility testing.

Inappropriate use of the veterinary medicinal product may increase the prevalence of bacteria resistant to florfenicol. Treatment should not exceed 5 days.

- ii. Special precautions to be taken by the person administering the veterinary medicinal product to animals
People with known hypersensitivity to polyethylene glycols should avoid contact with the veterinary medicinal product. In case of accidental spillage onto skin rinse with water.
- iii. Other precautions
Manure from treated pigs should be stored for one month prior to spreading and incorporating into land

Interactions with Other Medicaments No data available.

Statement on usage during pregnancy and lactation

Studies in laboratory animals have not revealed any evidence of potential embryotoxic or foetotoxic effect of florfenicol.

The safety of the veterinary medicinal product in sows has not been established during pregnancy and lactation.

The use is not recommended during pregnancy and lactation.

Adverse Effects/Undesirable Effects

A slight reduction of water consumption by the animals, dark brown faeces and constipation may be observed during treatment.

Commonly observed adverse effects are diarrhoea and/or peri-anal and rectal erythema/oedema which may affect approximately 40% of the animals. These effects are transient. In a few of the affected animals, prolapse of the rectum, that resolves without treatment, may be observed.

Overdose and Treatment

In case of overdosing, a decrease in weight gain, food and water consumption, peri-anal erythema and oedema and modification of some haematological and biochemical parameters indicative of dehydration may be observed.

Withdrawal Period(s)

Swine (Meat and offal): 20 days.

Broiler chicken: 5 days.

Storage Conditions Store below 30°C

Proposed Shelf Life (after first opening of container) 5 weeks.

Proposed Shelf Life (after dilution/reconstitution) 6 days.

Dosage Forms Oral solution

Packaging available 1Liter

Name and Address of Manufacturer

Samyang Anipharma Co., Ltd.

35, Songseon-ro 265 beon-gil, Pocheon-si, Gyeonggi-do, Korea

Date of Revision of Package Insert

5th March 2019 (1903-2TS)



SAMYANG ANIPHARM

35, SONGSEON-RO 265 BEON-GIL, POCHeon-SI, GYEONGGI-DO, KOREA

TEL : 82-2-388-2521 FAX : 82-2-357-8753