

Package Insert

TILMISEEN 25% W/V ORAL SOLUTION

Name and Strength of Active Substance(s)

Tilmicosin Phosphate 25%w/v

Product Description

Yellowish brown solution.

Pharmacodynamics

Tilmicosin is a semi-synthetic antibiotic of the macrolide group and is believed to affect protein synthesis. It has bacteriostatic action but at high concentrations it may be bactericidal.

This antibacterial activity is predominantly against Gram-positive microorganism with activity against certain gram-negative ones and *Mycoplasma* of a bovine, porcine, ovine and avian origin. In particular, its activity has been demonstrated against the following microorganism:

- *Pigs: Mycoplasma hyopneumoniae, Pasteurella multocida and Actinobacillus pleuropneumoniae*
- *Chickens and turkeys: Mycoplasma gallisepticum and Mycoplasma synoviae*
- *Calves: Mannheimia haemolytica, Pasteurella multocida, Mycoplasma bovis and M. dispar.*

Scientific evidence suggests that macrolides act synergistically with the host immune system. Macrolides appear to enhance phagocyte killing of bacteria. Tilmicosin has been shown to inhibit *in vitro* the replication of the Porcine Reproductive and Respiratory Syndrome virus in alveolar macrophages in a dose dependent fashion.

Cross-resistance between tilmicosin and other macrolides and lincomycin has been observed.

Pharmacokinetics

Whilst blood concentrations of tilmicosin are low, there is pH-dependent macrophage accumulation of tilmicosin in inflamed tissues.

Pigs: After oral administration of 200 mg tilmicosin/l drinking water, the average active substance concentrations detected in lung tissue, alveolar macrophages and bronchial epithelium 5 days after the start of treatment were 1.44 µg/ml, 3.8 µg/ml and 7.4 µg/g respectively.

Poultry: As early as 6 hours after oral administration of 75 mg tilmicosin/l drinking water, the average active substance concentrations detected in lung and alveolar tissue were 0.63 µg/g and 0.30 µg/g respectively. 48 hours after the start of treatment, the tilmicosin concentrations in lung and alveolar tissue were 2.3 µg/g and 3.29 µg/g respectively.

Environmental Properties

Not relevant

Indication

Pigs: For the treatment and prevention of respiratory disease in pig herds, associated with *Mycoplasma hyopneumoniae*, *Pasteurella multocida*, *Actinobacillus pleuropneumoniae* and other organisms susceptible to tilmicosin.

Chickens: For the treatment and prevention of respiratory disease in chicken flocks, associated with *Mycoplasma gallisepticum* and *M. synoviae*.

Recommended Dosage

Pigs: To be included in the drinking water to provide a daily dose of 15-20 mg/kg bodyweight for 5 days, which may be achieved by the inclusion of 200 mg tilmicosin per litre (80 ml Pulmotil AC per 100 litres).

Chickens and Turkeys (except hens producing eggs for human consumption): To be included in the drinking water at a daily dose of 15-20 mg/kg bodyweight in chickens and 10-27 mg/kg bodyweight in turkeys for 3 days, which may be achieved by the inclusion of 75 mg tilmicosin per litre (30 ml Pulmotil AC per 100 litres).

Mode of Administration

Orally

Contraindications

Do not allow horses and other equines access to drinking water containing tilmicosin.

Do not use in case of hypersensitivity to tilmicosin or to any of the excipients.

Warnings and Precautions

Important: Must be diluted before administration to animals.

Pigs, chickens and turkeys: Water consumption should be monitored in order to guarantee adequate dosing. In case water consumption does not match quantities for which recommended concentrations were calculated, the concentration of Pulmotil AC has to be adapted in a way that the recommended dosage will be taken up by the animals or different medication should be considered.

i. Special precautions for use in animals

For oral use only. Contains disodium edetate; do not inject

Severely ill individuals tend to drink less and may need simultaneous treatment, preferably by parenteral medication.

Inappropriate use of the product may increase the prevalence of bacteria resistant to tilmicosin and may decrease the effectiveness of treatment with tilmicosin-related substances. The use of the product should be based on susceptibility tests.

Medicated drinking water or milk replacer should be prepared fresh every 24 hours.

Official, national and regional antimicrobial policies should be taken into account when the product is used.

ii. Special precautions to be taken by the person administering the veterinary medicinal product to animals

- Tilmicosin may induce irritation. Macrolides, such as tilmicosin, may also cause hypersensitivity (allergy) following injection, inhalation, ingestion or contact with skin or eye. Hypersensitivity to tilmicosin may lead to cross reactions to other macrolides and

vice versa. Allergic reactions to these substances may occasionally be serious and therefore direct contact should be avoided.

- To avoid exposure during preparation of the medicated drinking water, wear overalls, safety glasses, and impervious gloves. Do not eat, drink or smoke when handling this product. Wash hands after use.
- In the case of accidental ingestion, wash out mouth immediately with water and seek medical advice. In the event of accidental skin contact, wash thoroughly with soap and water. In case of accidental eye contact, flush the eyes with plenty of clean, running water.
- Do not handle the product if you are allergic to ingredients in the product.
- If you develop symptoms following exposure, such as skin rash, you should seek medical advice and show the physician this warning. Swelling of the face, lips and eyes or difficulty in breathing are more serious symptoms and require urgent medical attention.

Interactions with Other Medicaments

None known.

Statement on usage during pregnancy and lactation

The safety of tilmicosin has not been established in animals used for breeding purposes.

Adverse Effects/Undesirable Effects

In very rare cases, a decrease in water intake has been observed.

Overdose and Treatment

When pigs are offered drinking water containing 300 or 400 mg/litre (equivalent to 22.5-40 mg/kg bodyweight or 1.5-2 times the recommended concentration) commonly animals exhibit a reduced water intake. Although this has a self-limiting effect on tilmicosin intake, it could, in extreme circumstances, result in dehydration. This can be corrected by the removal of the medicated drinking water and replacement with fresh unmedicated water.

No symptoms of overdose have been seen in chickens given drinking water containing levels of tilmicosin up to 375 mg/litre (equivalent to 75-100 mg/kg bodyweight or 5 times the recommended dose) for 5 days; daily treatment with 75 mg/litre (equivalent to the maximum recommended dose) for 10 days resulted in a reduction in faecal consistency.

No symptoms of overdose have been seen in turkeys given drinking water containing levels of tilmicosin up to 375 mg/litre (equivalent to 50-135 mg/kg bodyweight or 5 times the recommended dose) for 3 days; daily treatment with 75 mg/litre (equivalent to the maximum recommended dose) for 6 days also produced no symptoms of overdose.

No symptoms of overdose, with exception of a slight decrease in the milk consumption, have been seen in calves given twice daily doses 5 times the maximum recommended dose or for twice the maximum recommended duration of treatment.

Withdrawal Period(s)

Pigs – 14 days

Chickens – 12 days.

Not authorised for use in laying birds producing eggs for human consumption. Do not use within 14 days of onset of the laying

Not for use in lactating animals.

Storage Conditions

Store below 30°C.

Shelf life

3 years

Shelf Life (after first opening of container)

5 weeks

Shelf Life (after dilution/reconstitution)

6 days

Dosage Forms

Oral solution

Packaging available

500ml

Name and Address of manufacturer/marketing authorisation holder

Samyang Anipharm Co.Ltd.

35, Songseon-ro 265 beon-gil, Pocheon-si, Gyeonggi-do, Korea

Marketing Authorisation Holder

Taseen Trading Sdn. Bhd. No.8, Jalan Astana 1F/KU2,

Bandar Bukit Raja, 41050 Klang, Selangor, Malaysia.

Date of Revision of Package Insert

27th Nov 2018.



SAMYANG ANIPHARM

35, SONGSEON-RO 265 BEON-GIL, POCHON-SI, GYEONGGI-DO, KOREA

TEL : 82-2-388-2521 FAX : 82-2-357-8753