

**Tylong 50%**  
Water Soluble Powder  
(FOR VETERINARY USE ONLY)

**COMPOSITION**

Tylosin ..... 500 mg  
Exp q.s to ..... 1 g

**PRODUCT DESCRIPTION**

The finished product is a yellowish color powder.

**TARGET SPECIES**

Calves, Pigs, Chickens and Turkeys

**PHARMACODYNAMIC**

Tylosin is a macrolide antibiotic produced by a strain of *Streptomyces fradiae*. It exerts its antimicrobial effect by inhibiting protein synthesis of susceptible micro-organisms.

The tylosin spectrum of activity includes Gram-positive bacteria including *Clostridium perfringens* and some Gram-negative strains such as *Pasteurella* and *Mycoplasma spp.* at concentrations of 16µg/ml or less.

**PHARMACOKINETIC**

**Absorption:** Tylosin reaches maximal blood levels between 1 and 3 hours after an oral dose. Minimal or no blood levels remain 24 hours after an oral dose.

**Distribution:** After oral doses were given to pigs, tylosin was found in all tissues, between 30 minutes and two hours after administration, except for the brain and spinal cord.

**Biotransformation and Elimination:** It has been shown that most of the material which is excreted is to be found in the faeces and consists of tylosin (factor A), relomycin (factor D) and dihydrodesmycosin.

**INDICATIONS**

For the control of *Mycoplasma synoviae* airsacculitis in chickens and *Mycoplasma gallisepticum* S6 in chickens and turkeys. In the field tylosin has also proved useful in reducing the level of infection following stress associated with live vaccination.

As an aid in the control of outbreaks of necrotic enteritis in chickens caused by *Clostridium perfringens*.

For the prevention and control of enzootic pneumonia, and scours caused by organisms sensitive to tylosin, in pigs.

For the control of pneumonia in cattle associated with mycoplasmata and *Pasteurella multocida* sensitive to tylosin.

**WARNINGS AND PRECAUTIONS**

**Special warnings for each target species**

None

**Special precautions for use**

i) Special precautions for use in animals:

Use of the product should be based on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy

should be based on local (regional, farm level) epidemiological information about susceptibility of the target bacteria.

Official, national and regional antimicrobial policies should be taken into account when the product is used.

A high rate of in vitro resistance has been demonstrated in European strains of *Brachyspira hyodysenteriae* implying that the product will not be sufficiently efficacious against swine dysentery.

Use of the product deviating from the instructions given in the SPC may increase the prevalence of bacteria resistant to tylosin and may decrease the effectiveness of treatment with other macrolides, lincosamides and streptogramin B due to the potential for cross-resistance.

ii) Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Tylosin may induce irritation. Macrolides, such as tylosin, may also cause hypersensitivity (allergy) following injection, inhalation, ingestion or contact with skin or eye. Hypersensitivity to tylosin may lead to cross reactions to other macrolides and vice versa. Allergic reactions to these substances may occasionally be serious and therefore direct contact should be avoided.

To avoid exposure during preparation of the medicated drinking water, wear overalls, safety glasses, impervious gloves and wear either a disposable half mask respirator conforming to European Standard EN149 or a non-disposable respirator to European Standard EN140 with a filter to EN143. Wash hands after use.

In the event of accidental skin contact, wash thoroughly with soap and water. In case of accidental eye contact, flush the eyes with plenty of clean, running water.

Do not handle the product if you are allergic to ingredients in the product.

If you develop symptoms following exposure, such as skin rash, you should seek medical advice and show the physician this warning. Swelling of the face, lips and eyes or difficulty in breathing are more serious symptoms and require urgent medical attention.

- Keep medicine out of reach of children.
- Jauhkan ubat dari kanak-kanak

**CONTRA-INDICATIONS**

Do not leave or dispose of water containing tylosin tartrate where it may be accessible to either animals not under treatment or to wildlife.

Do not use in known cases of hypersensitivity to tylosin or other macrolides.

**ADMINISTRATION & DOSAGE**

For oral administration.

Recommendations for use in **chickens** and **turkeys**:

Tylong 50% is administered in the drinking water at a concentration of 1 g product per litre.

As an aid in the control of outbreaks of necrotic enteritis caused by *Clostridium perfringens* in chickens use Tylong 50% in the drinking water for 5 days at a concentration of 0.3 g product per litre water, to provide 20-50 mg/kg bw, depending on the age and the water consumption of the birds.

Recommendations for use in **pigs**:

Tylong 50% is administered in the drinking water to provide 25 mg tylosin/kg bodyweight for 3-10 days. This may normally be achieved by adding 0.50 g product per litre.

Recommendations for use in **calves**:

1 g tylosin/calf administered orally twice daily for 7-14 days. The tylosin tartrate may be incorporated into the milk or reconstituted milk replacer at the time of feeding.

**INTERACTION WITH OTHER MEDICAMENTS**

None known

**PREGNANCY AND LACTATION**

No adverse effects to tylosin have been seen in fertility, multi-generation or teratology studies.

**OVERDOSE AND TREATMENT**

There is no evidence of tylosin toxicity in animals, at dose rates of up to 1000 mg/kg.

**SIDE EFFECT**

None known

**WITHDRAWAL TIME**

Turkeys & Pigs (Meat): 0 day

Calves (Meat): 14 days

Chicken (Meat):

- Following a dose rate of 50-200 mg tylosin/kg bw: 1 days

- Following a dose rate of 20-50 mg tylosin/kg bw: 0 day

Chicken (Egg): 0 day

**SHELF LIFE:** 36 months

**SHELF LIFE after first opening & reconstitution:** 24 hours

**STORAGE**

Store in a cool dry place ( $\leq 25^{\circ}\text{C}$ ). Strictly avoid light & heat exposure.

**PACKAGING**

The powder filled 150 g & 750 g in aluminum sachets, by automatic filling machine and sealing, and packed in carton box.

**DATE OF REVISION:** 10 Apr 2019



Manufactured & Distributed By:

**SHENNONG ANIMAL HEALTH (M) SDN BHD**

神農動物保健(馬)有限公司

**No. 2, 4, 6, 8 & 10, Jalan Industri USJ 1/19,**

**Taman Perindustrian USJ 1,**

**47610 Subang Jaya, Selangor, Malaysia.**