



Osmosis Nutrition Sdn Bhd (594876-A)

Package Insert

Tilmicos 25% w/v Oral Solution

Active Ingredient

Each ml contains:

Tilmicosin (as phosphate) 250 mg

Description

A yellow to amber-colored solution containing Tilmicosin as an active ingredient in 250mg/ml. Disodium Edetate/Disodium Ethylenediaminetetraacetate (2 mg per mL) was added as stabilizer, Sodium Benzoate (10 mg per mL) was added as preservative while purified water (added to 1 mL) as its vehicle. Phosphoric Acid/Orthophosphoric was used as pH adjuster to adjust the solution in a pH range between 4.50 – 5.50.

Pharmacodynamics:

Tilmicosin is a mainly bactericidal semi synthetic antibiotic of the macrolide group. it is believed to affect bacterial protein synthesis.

Tilmicosin has a wide spectrum of activity against gram positive organisms and is particularly active against Pasteurella, Actinobacillus (Haemophilus) and Mycoplasma organisms. Tilmicosin has some activity against certain gram negative micro-organisms.

Pharmacokinetics:

Tilmicosin is absorbed and moves rapidly out of the serum into areas of low pH. this results in very low serum concentrations, but detectable levels of tilmicosin are found in lung tissue as early as 6 hours after starting the treatment.

Tilmicosin is also detected in pooled air sac tissue as early as 6 hours after starting the treatment.

Indication:

Chickens

For the treatment and prevention of respiratory disease in chicken flocks, associated with Mycoplasma gallisepticum and M. synoviae.

Turkeys

For the treatment and prevention of respiratory disease in turkey flocks, associated with Mycoplasma gallisepticum and M. synoviae.

Recommended Dose:

Important:

Must be diluted before administration.

To be included in the drinking water at 75mg tilmicosin per litre for 3 days.


A daily dose of 15-20 mg/kg bodyweight in chickens or 10-27 mg/kg in turkeys.

Medicated drinking water should be used within 24 hours, using only clean water, according to the following dilution table:

QUANTITY OF TILMICOS 25% TO BE ADDED	DRINKING WATER	
	Litres	Gallons
5 ml	17	3.7
10 ml	33	7.3
15 ml	50	11
20 ml	67	14.7
30 ml	100	22
60 ml	200	44
240 ml	800	176
960 ml	3200	704

Mode of Administration:

Oral Administration.



Osmosis Nutrition Sdn Bhd (594876-A)

Contraindication:

Not to be used in animals with known sensitivity to the active ingredients.

Warnings and Precautions:

Keep out of reach of children.

Jauhi dari kanak-kanak.

Special precautions for use in animals

Chickens and turkeys: Due to the administration route and as water consumption depends on the clinical condition of the animal, in order to ensure a correct dosage, the concentration of the product must be adjusted according to the water intake. If this is not possible, then an alternative medication may be required.

Animals with acute infections and severely reduced feed intake should be treated first with a suitable injectable product.

The medicated water should be prepared fresh every 24 hours.

Inappropriate use of the product may increase the prevalence of bacteria resistant to tilmicosin and may decrease the effectiveness of treatment with tilmicosin related substances. It is sound clinical practice to base treatment on susceptibility testing.

Do not allow horses or other equines access to drinking water containing tilmicosin.

Special precautions for the person administering the veterinary medicinal product to animals

People with known hypersensitivity to tilmicosin should avoid contact with the product. The veterinary medicinal product may cause irritation or sensitisation by skin contact.

Avoid skin and ocular contact. Wear protective gloves and protective clothes when handling the veterinary medicinal product.

In case of contact with skin or eyes, rinse abundantly with fresh water. If irritation persists and in case of incidental ingestion, seek immediately medical advice or call a poison center (dangers linked to disturbances in cardiac conduction).

Interactions with other Medicaments:

None known.

Statement on usage during pregnancy and lactation:

Do not use in pullets and laying hens producing eggs for human consumption.

Use only in accordance with risk/benefit assessment by the responsible veterinarian.

Side Effects:

None known.

Symptoms and Treatment of Overdose:

No symptoms were seen in poultry treated at 375 mg/litre during 5 days. A dose of 75 mg/litre during 10 days resulted in less consistent faeces.


Withdrawal Period:

Chickens - 12 days

Turkeys – 19 days

Not authorized for use in laying birds producing eggs for human consumption.

Do not use within 14 days of onset of the laying.



Osmosis Nutrition Sdn Bhd (594876-A)

Maximum Residual Limit (MRL):

Chicken & Turkey

Muscle: Not More Than 75 microgram/kg

Skin/fat: Not More Than 75 microgram/kg

Liver: Not More Than 1000 microgram/kg

Kidney: Not More Than 250 microgram/kg

Storage Conditions:

Store below 30°C.

Protect from direct sunlight.

Pack Size:

1 Liter in White HDPE Plastic bottle with white HDPE Plastic Cap

Manufactured by Contract Manufacturer:

Osmosis Nutrition Sdn. Bhd.,

Lot 16014 (PT No.24341), Jalan Nilam 3,

Bandar Nilai Utama,

71800 Nilai, Negeri Sembilan,

Malaysia.

Distributed by Product Registration Holder:

Vision Biotech Sdn. Bhd.,

PT 832, Jalan Teratai 3,

Taman Indah Site A,

43900 Sepang Pos,

Negeri Sembilan.

Date of Revision:

28-09-2015