

ALFASAN
DORMINAL 20% INJECTION
(PENTOBARBITAL SODIUM 200MG/ML)

Active Substance

Pentobarbital Sodium 200 mg/ml (Equivalent to 182.3mg/ml pentobarbital)

Product Description

A limped pale red coloured injectable solution packed in colourless vials, glass type 11, of 100mL with butyl rubber stoppers and non- reusable aluminium closures.

Pharmacodynamics

Pentobarbitone is an anaesthetic agent and belongs to the group of barbituric acid derivatives. The LD50 in dogs and cats is approximately 40 to 60mg/kg bodyweight when injected intravenously.

In endothermic animals, the immediate effect is the loss of consciousness followed by deep anaesthesia followed by death. Breathing stops and is quickly followed by cardiac arrest.

Pharmacokinetics

Pentobarbital is absorbed quite rapidly from the gut after oral or rectal administration with peak plasma concentrations occurring between 30- 60 minutes after oral dosing in humans. The onset of action usually occurs within 15-60 minutes after oral dosing and within 1 minute after IV administration.

Pentobarbital, like all barbiturates, distributes rapidly to all body tissues with highest concentrations found in the liver and brain. It is 35-45% bound to plasma proteins in humans. Although less lipophilic than the ultra-short acting barbiturates (e.g. thiopental), pentobarbital is highly lipid soluble and patient fat content may alter the distributive qualities of the drug. All barbiturates cross the placenta and enter milk (at concentrations far below those of plasma).

Pentobarbital is metabolized in the liver principally by oxidation. Excretion of the drug is not appreciably enhanced by increasing urine flow or alkalinizing the urine. Ruminants (especially sheep and goats) metabolize pentobarbital at a very rapid rate. The elimination half-life in the goat has been reported to be approximately 0.9 hrs. Conversely, the half-life in dogs is approximately 8 hours and ranges from 15-50 hours in man.

Indication

For euthanasia of domestic pets and smaller farm animals.

Recommended Dosage

0.7mL per kg bodyweight by rapid intravenous injection. The product solution is not intended for dilution with water or any other fluid.

Mode of Administration

By rapid intravenous injection.

Contraindications

Do not use in animals intended for human consumption. Do not use for anaesthetic purposes.

Warnings and Precautions

Special warnings for each target species

Carcasses of animals euthanized with this product should be disposed of in accordance with national legislation.

Carcasses of animals euthanized with this product should not be fed to other animals due to the risk of secondary intoxication.

The intraperitoneal route of administration may cause a prolonged onset of action with an increased risk of adverse effects noted in 4.6. Prior sedation is advisable.

The intrapulmonary route of administration may cause a prolonged onset of action with an increased risk of adverse effects noted in 4.6 and should be reserved for cases where other routes of administration are not possible. Prior sedation is mandatory before this route of administration is employed.

When euthanasia of poikilotherms is undertaken, the animal must be maintained at its preferred optimum temperature, otherwise efficacy may be unreliable. Species appropriate measures (e.g. pithing) should be taken to ensure that euthanasia is complete in order that spontaneous recovery sometime later, does not occur.

Special precautions for use***Special precautions for use in animals:***

Venomous snakes are best euthanized by intra-coelomic injections of sodium pentobarbitone solution with judicious use of prior sedation in order to minimize danger to humans.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

This is a potent drug which is toxic in man – particular care should be taken to avoid accidental ingestion and self- injection. Wear suitable protective gloves when handling this product - pentobarbital can be absorbed via skin and mucosa.

In the event of accident the following action should be taken:

Skin – Wash immediately with water and then thoroughly with soap and water.

Eyes – Wash immediately with cold water and obtain medical advice if irritation persists.

Ingestion – Obtain medical attention immediately. Wash out mouth. Keep warm and rest.

Accidental self-injection – Obtain URGENT medical attention (take the package insert with you), advising medical services of barbiturate poisoning. Do not leave patient unattended.

Advice to Doctor – Maintain airways and give symptomatic and supportive treatment.

Interactions with Other Medicaments

CNS depressants (narcotics, phenothiazines, antihistamines, etc.) may increase the effect of pentobarbital.

Pregnancy and Lactation

Not relevant.

Adverse Effects

Minor muscle twitching may occur after injection. In cattle, gasping may occur in rare cases if Pentobarbital-Sodium is administered below the recommended dose. Death may be delayed if the injection is administered perivascularly. Barbiturates can be irritating when administered subcutaneously or perivascularly.

Symptoms and Treatment of Overdose

In the event of accidental administration to an animal not presented for euthanasia, measures such as artificial respiration, administration of oxygen and the use of analeptics are appropriate.

Incompatibilities

None known.

Withdrawal Period

Not applicable.

Storage Conditions

Store below 25°C and protect from light.

Once a vial has been broached, the contents should be used within 1 month and stored in the refrigerator (2- 8°C). Keep out of reach of children. Jauhi dari kanak-kanak.

Dosage Forms and Packaging Available

Injectable solution available in packaging of 100ml.

Manufacturer

Alfasan International BV

Kuipersweg 9, 3449 JA Woerden, The Netherlands

Product Registration Holder

Delfri Marketing Sdn. Bhd. (15872-P)

Level 6, Block A, Sky Park One City, Jalan USJ 25/1, 47650 Subang Jaya, Selangor, Malaysia

Date of Revision

17/03/2015