

[Draft of Package Insert]

**For Animal Use Only**

# NEOSUL-242

**POWDER**

**CONTENT:** Each sachet (500g) contains:

Neomycin Sulphate.....400g (Equivalent to neomycin 242g)

**PRODUCT DESCRIPTION:** A yellowish white color powder.

**PHARMACODYNAMICS:**

Neomycin is an aminoglycoside antibiotic consisting of 3 components, A, B and C. Neomycin acts by interfering with bacterial protein synthesis by binding primarily the 30S subunit of bacterial ribosomes.

**PHARMACOKINETICS:**

Neomycin is poorly absorbed from the gastrointestinal tract of animals, and it has low absorption from the udder. Once it is absorbed it is rapidly excreted by the kidneys in active form. It has been reported to have a half life of 2 to 3 hours. About 97% of an oral dose being excreted uncharged in the faeces.

**INDICATION:** Antimicrobial

Chickens and turkeys: As an aid in the treatment of bluecomb or transmissible enteritis in turkeys, bacterial enteritis, enteritis associated with chronic respiratory disease (CRD) and other neomycin-sensitive infections in chickens and turkeys.

Swine, cattle, sheep and horses: As an aid in the treatment of scours (bacterial enteritis).

**DOSAGE & ADMINISTRATION:**

Chickens and turkeys: Flock treatment: As soon as symptoms are observed, give NEOSUL-242 POWDER for 3 to 5 days. If no improvement is noted after 5 days of treatment, the diagnosis should be redetermined.

Add NEOSUL-242 POWDER to the drinking water as follows:

Chickens:

- Day-old to 2 weeks: 20 g of powder per 223.4 L of drinking water or 100 g of powder per 1116.8 L of drinking water.
- 2 weeks to adulthood: 40 g of powder per 223.4 L of drinking water or 200 g of powder per 1116.8 L of drinking water.

Turkeys:

- Bacterial enteritis and enteritis associated with CRD: 40 g of powder per 223.4 L of drinking water or 200 g of powder per 1116.8 L of drinking water.
- Bluecomb or transmissible enteritis: 100 g of powder per 332.6 L of drinking water for 8 days medication, as per the following schedule:
  - a. 3 days - medication

- b. 2 days - no medication
- c. 3 days - medication
- d. 2 days - no medication
- e. 2 days - medication

Swine, cattle, sheep and horses:

- Individual treatment: Add to the drinking water consumed daily, 0.99 g of powder per 50 kg of bodyweight.
- Herd treatment: Add to the drinking water consumed daily, 99 g of powder per 5000 kg of bodyweight.

**CONTRAINDICATIONS:**

Do not use in animals with known hypersensitivity to active ingredients.

Do not use in calves to be processed for veal.

**WARNINGS / PRECAUTIONS:**

To administer the stated dosage, the concentration of neomycin required in medicated water must be adjusted to compensate for variation in age and weight of animal, the nature and severity of disease signs, and environmental temperature and humidity; each of which affects water consumption.

If symptoms persist after using this preparation for 2 to 3 days, consult a veterinarian. If symptoms such as fever, depression, or going off feed develop, oral neomycin is not indicated as the sole treatment since systemic levels of neomycin are not obtained due to low absorption from the gastrointestinal tract.

**DRUGS INTERACTIONS:** None known.

**PREGNANCY AND LACTATION:**

In the absence of toxicology study, this veterinary medicinal product must not be used during pregnancy and lactation. Do not use in laying birds producing eggs for human consumption.

**ADVERSE EFFECTS:** None known.

**OVERDOSE AND TREATMENT:**

When given orally in large doses, neomycin causes nausea, vomiting, and diarrhoea.

Prolonged oral use may cause a malabsorption syndrome with steatorrhoea and diarrhoea which can be very severe.

**STORAGE CONDITION:** Store below 30°C.

**WITHDRAWAL PERIOD(S):**

Treated animals must not be slaughtered for use in food for at least: broilers - 7 days, laying hens, turkeys, pigs, sheep - 14 days, cattle - 30 days after the latest treatment with this drug. This drug is not to be administered to horses that are to be slaughtered for use in food.

**DOSAGE FORMS AND PACKAGING AVAILABLE:** 500g powder packed into aluminium sachet.

**DATE OF REVISION OF PACKAGE INSERT:** 07 OCTOBER 2015.

Manufactured by:

**PAHANG PHARMACY SDN. BHD. (10826-K)**

LOT 5979, JALAN TERATAI,  
5½ MILE OFF JALAN MERU, 41050 KLANG,  
SELANGOR D.E., MALAYSIA.

Product Registration Holder:

**GENPHARMA SDN. BHD. (268066-W)**

LOT 5979, JALAN TERATAI,  
5½ MILE OFF JALAN MERU, 41050 KLANG,  
SELANGOR D.E., MALAYSIA.