

Salicyl 1000mg/g Soluble Powder

FOR ANIMAL USE ONLY

Brand or Product Name :

Salicyl 1000mg/g Soluble Powder

Name and Strength of Active Substance(s)

Each gram contains 1000 mg Sodium Salicylate

Product Description

Powder for oral solution. White or off white flakes.

Pharmacodynamic Properties

Sodium salicylate is a non-steroid anti-inflammatory drug (NSAID) and has an antiinflammatory, analgesic and antipyretic effect. The mode of action is based on inhibition of the enzyme cyclooxygenase, resulting in decreased production of prostaglandin (inflammation mediators).

Pharmacokinetics

Orally administered sodium salicylate is rapidly absorbed by passive diffusion, partially from the stomach, but mainly from the anterior part of the small intestine. Sodium salicylate distributes very well to the various tissues. Values of volume of distribution (Vd) are higher in the newborns. Half lives are longer in the very young resulting in slower elimination of the substance. This is most prominent in animals up to 7-14 days of age. Metabolism takes place mainly in the endoplasmatic reticulum and the mitochondria of the liver cells. Elimination occurs mainly via the urine and the pH of the urine can have a major effect on this elimination.

Indication

Calves: For supportive treatment of pyrexia in acute respiratory disease, in combination with appropriate (anti-infective) therapy if necessary.

Pigs: For the treatment of inflammation, in combination with concurrent antibiotic therapy.

Target Species

Calves and pigs.

Recommended Dosage

Calves: 40 mg sodium salicylate per kg of bodyweight once daily for 1 - 3 days.

Pigs: 35 mg sodium salicylate per kg of bodyweight per day for 3 - 5 days.

Can be administered orally in drinking water or milk.

Direction for Use

Prepare fresh medicated solution daily. Salicyl can be administered with the drinking water as pulse medication. Half of the calculated total daily amount of powder is mixed with 5 – 10 L of clean water and stirred until evenly dispersed. This solution is then added, while stirring, into an amount of drinking water that will be consumed within approximately 3 – 4 hours and administered twice daily.

Route of Administration

Administer orally in drinking water or milk.

Contraindications

Do not administer in case of severe hypoproteinaemia, liver and kidney disorder.

Do not administer in case of gastrointestinal ulcerations and chronic gastrointestinal disorders.

Do not administer in case of malfunction of the haemopoietic system, coagulopathy, haemorrhagic diathesis.

Do not use sodium salicylate in neonates or calves less than 2 weeks of age.

Do not use in piglets of less than 4 weeks of age.

Do not use in animals with known hypersensitivity to sodium salicylate.

Warnings and Precautions

Special warnings for each target species

None known.

Special precautions for use

i. **Special precautions for use in animals :** Given that sodium salicylate may inhibit clotting of blood, it is recommended that elective surgery should not be performed on animals within 7 days after the end of treatment.

ii. **Special precautions to be taken by the person administering the veterinary medicinal product to animals**

People with known hypersensitivity to sodium salicylate or correlated pharmaceuticals (e.g. aspirin) or excipients should avoid contact with the veterinary medicinal product. Irritation of the skin, eye, and respiratory tract might occur. During preparation and mixing of the product, direct contact with the skin and eyes, and direct inhalation of the powder should be avoided. It is recommended to wear gloves, safety glasses, and a dust mask. Particular attention should be taken when opening the bucket. Wash accidental exposure to skin immediately with water. In the event of accidental eye contact, the user is advised to wash the eye with plenty of water for 15 minutes, and seek medical advice if irritation persists. During administration of the medicine as reconstituted product in drinking water or milk (replacer) to the animals, skin contact should be prevented by wearing gloves. Wash accidentally exposed skin immediately with water.

Interaction with Other Medicaments

Concurrent administration of potentially nephrotoxic drugs (e.g. aminoglycosides) should be avoided. Salicylic acid is highly plasma (albumin) bound and competes with a variety of compounds (e.g. ketoprofen) for plasma protein binding sites. Plasma clearance of salicylic acid has been reported to increase in combination with corticosteroids possibly due to induction of metabolism of salicylic acid. Concurrent use with other NSAIDs is not recommended, because of increased risk of gastro-intestinal ulceration. Drugs which affect blood clotting should not be used in combination with sodium salicylate.

Statement on usage during pregnancy and lactation

The use is not recommended during pregnancy and lactation because laboratory studies in rats have shown evidence of teratogenic and foetotoxic effects. Salicylic acid crosses the placenta and is excreted with the milk. Half-life in the newborn is longer and thus toxicity symptoms may occur much sooner. Furthermore platelet aggregation is inhibited and bleeding time increased which is not favourable during difficult parturition / caesarean section. Finally some studies indicate that parturition is postponed.

Side Effects

Gastrointestinal irritation may occur especially in animals with pre-existing gastrointestinal disease. Such irritation may be clinically manifested by production of black faeces due to bleeding in the gastrointestinal tract. Inhibition of normal blood clotting may occur incidentally. This effect is reversible and diminishes within approximately 7 days.

Symptoms and Treatment of Overdose

Calves tolerate dosages up to 80mg/kg for 5 days or 40mg/kg for 10 days without any adverse effects. Pigs tolerate dosages up to 175mg/kg for up to 10 days without any significant adverse effects. In case of an acute overdose intravenous bicarbonate infusion results in a higher clearance of salicylic acid by alkalization of the urine and may be beneficial in correcting (secondary metabolic) acidosis.

Storage Conditions

Store in a dry place below 30°C. Keep out of reach of children. Jauhkan ubat daripada kanak-kanak.

Shelf-life

Shelf life as packaged for sale: 3 years.

Shelf life after first opening of the immediate packaging: 24 hours.

Shelf life after reconstitution in drinking water: 24 hours.

Shelf life after reconstitution in milk (replacer): 6 hours.

Withdrawal Period(s)

Meat and offal: Pigs: 3 days. Calves: zero days. Do not use in cows producing milk for human consumption

Maximum Residual Limit (MRL)

No MRL required for sodium salicylate regarding oral use in bovine and porcine species. (EMA/CVMP/71291/2012)

Disposal of containers

Empty aluminum sachet must be safely disposed of according to local government regulations.

Dosage Forms and packaging available

Available in 100g, 200g, 500g and 1000g powder per sachet, pack in 10, 20, 50 and 100 sachets per box.

Name and Address of manufacturer

Life Biopharma Sdn. Bhd. (company no. 546163-D)

No. 250-251, Jalan Industri Galla 13, Kawasan Perindustrian Galla, 70200 Seremban, Negeri Sembilan, Malaysia.

Date of Revision of Package Insert

2015-10-29