150mm

105mm

TIGER BALM®

ACTIVE

MUSCLE GEL

Non-sticky pain relief gel

Tiger Balm Active Muscle Gel

Name and Strength of Active Substance(s)
Methyl Salicylate 16% w/w
Menthol 8% w/w

Product Description
A clear transparent greenish-blue gel with characteristic smell.

Pharmacodynamics
Menthol
Menthol dilates the blood vessels causing a sensation of coldness, followed by an analgesic effect. Menthol also acts as a penetration enhancer, increasing the penetration of drugs when applied on the skin, to give faster onset of action.

Methyl salicylate
Methyl salicylate is a salicylic acid derivative.
Salicylates inhibit cyclooxygenase, thereby reducing the formation of prostaglandins, and cause platelet dysfunction.
Methyl salicylate is used topically as a counter-irritant. Upon application, it is absorbed through the skin and is applied for the relief of pain in rheumatic conditions and painful muscles or joints.

Pharmacokinetics
Menthol
- After absorption, menthol is excreted in the urine and bile as a glucuronide.

Methyl salicylate
- May be absorbed through intact skin.
- The absorption of topical salicylates is proportional to the surface area involved, duration of exposure, concentration and skin integrity.
- Absorption characteristics of salicylates vary with the dose, formulation and route of administration.
- Percutaneous absorption is enhanced by exercise, heat, occlusion, or disruption of the integrity of the skin or application to large areas of skin.
- Both the rate and extent of absorption increases after repeated application; increasing the bioavailability.
- Methyl salicylate is extensively metabolised to salicylic acid in the dermal and subcutaneous tissues after topical application.
- At therapeutic levels, the half life of salicylates is 2-4 hours. As salicylate level increases into the toxic range, the half-life can be greater than 18 hours.

Pharmacological Information
Tiger Balm Active Muscle Gel has mild analgesic and local anaesthetic effects. When massaged gently on the skin, a cooling effect is felt initially, followed by a sensation of warmth.

Indication
Tiger Balm Active Muscle Gel provides temporary pain relief from minor aches and pains of muscles and joints associated with arthritis, simple backache, strains and sprains.

Recommended Dosage
Adults & children above 2 years old.
Apply to the affected area not more than 3 to 4 times daily.
Rub/Massage gently.
For external application only.

Mode of Administration
Topical

Contraindications:
- Do not use on patients with aspirin or salicylate idiosyncrasy.
- Hypersensitivity to any of the ingredients in the product.
- Do not use during pregnancy or when breastfeeding.
- This product should not be used on children under 2 years of age.
- Use with caution on older children.

Warnings and Precautions
Under Caution:
This product contains methyl salicylate and when applied or rubbed on to the skin, can be absorbed through the skin into the blood. For patients taking warfarin, excessive application on to the skin for muscle or joint pains may increase the chances of bleeding. If you are taking warfarin, please consult your doctor before using this product.

Warning:
- Topical analgesic preparations containing methyl salicylate should be used with caution in patients at increased risk of developing salicylate adverse effects.
- Children suffering from flu, chickenpox, or fever should avoid using this product because salicylates may induce Reyes Syndrome.

MAL15125044X

Front

150mm

TIGER BALM®

ACTIVE

MUSCLE GEL

Non-sticky pain relief gel

Precautions:
- Initially apply a small amount to a limited skin area and observe whether there is discomfort or itch before applying to a larger skin area. Discontinue use if irritation develops.
- For external use only; not to be taken.
- Avoid contact with the eyes and mucous membranes.
- Do not apply to open wounds or damaged skin.
- Do not apply over large areas of the body.
- Do not use bandage or heating pad on the applied area.
- If symptoms persist, consult your doctor.

Effects on Ability to Drive and Use Machines
None known

Interactions with Other Medicaments
This product contains methyl salicylate and when applied or rubbed on to the skin, can be absorbed through the skin into the blood. For patients taking warfarin, excessive application on to the skin for muscle or joint pains may increase the chances of bleeding.

Statement on Usage during Pregnancy and Lactation
Do not use during pregnancy or when breastfeeding.

Adverse Effects/Undesirable Effects
Headache, dizziness, nausea and vomiting, skin irritation, contact dermatitis, rash, itching, redness or swelling, burning or stinging sensation may occur.

May cause hypersensitivity/allergic reactions on some individuals with sensitive skin.

Overdose and Treatment
Symptoms of methyl salicylate overdose
Salicylate intoxication can occur after ingestion or topical application of methyl salicylate.
Mild chronic salicylate intoxication, or salicylism, usually occurs only after repeated use of large doses. Salicylism can also occur following excessive topical application of salicylates.
Symptoms include dizziness, tinnitus, deafness, sweating, nausea and vomiting, headache, and confusion, and may be controlled by reducing the dosage.
Symptoms of more severe intoxication or of acute poisoning following overdose include hyperventilation, fever, restlessness, ketosis, and respiratory alkalosis and metabolic acidosis. Depression of the CNS may lead to coma; cardiovascular collapse and respiratory failure may also occur.

Symptoms of menthol overdose
Ingestion of significant quantities is reported to cause symptom: severe abdominal pain, nausea, vomiting, vertigo, ataxia, drowsiness, and coma; instant collapse in infants after the local application of menthol to their nostrils.

Treatment
The stomach should be emptied by gastric lavage or administration of oral activated charcoal.
Fluid and electrolyte management is the mainstay of treatment with the immediate aim of correction of acidosis, hyperpyrexia, hypokalaemia and dehydration if present.
Any convulsions must be controlled first through supportive care including anticonvulsant therapy.

Storage Conditions
Store below 30°C.

Dosage Forms and Packaging Available
5g & 60g

Name and Address of Manufacturer
Manufactured under license from Haw Par Corporation Ltd., Singapore by Tiger Balm (M) Sdn. Bhd. No. 53, Jalan I-Park SAC 7, Taman Perindustrian I-Park SAC, 81400 Senai, Johor, Malaysia.

Name and Address of Product Registration Holder
Tiger Balm (M) Sdn. Bhd.
No. 53, Jalan I-Park SAC 7, Taman Perindustrian I-Park SAC, 81400 Senai, Johor, Malaysia.

Date of P.I.L : 13/11/24

MP1M0106 2

Back

FONT SIZE: 5.25pt

Client	**Haw Par Healthcare Pte Ltd**	Person to contact	**Shermaine / Hui Hui**	Software Version	**Adobe Illustrator CS6**
Artwork	**Visibility Design Pte Ltd**	Job No.	**Q/HP/1024/6792**	Date In	**14 Nov 2024**
Revision Code	**VD-00**	File	**Malaysia TB Active Muscle Gel 60g Insert (TBM)**	Material Code	**MP1M0106 2**
Last date of amendment: **14 Nov 2024**					

Black

TYPE FACES:

ENGLISH: HELVETICA NEUE