

**KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY**

DECCOX[®] POWDER 6% W/W

ACTIVE CONSTITUENT: Each kilogram contains 60g of Decoquinatate.

PRODUCT DESCRIPTION: Deccox[®] Powder 6% W/W is yellow, fine granular powder feed additive.

INDICATION, DOSAGE & ADMINISTRATION

Oral administration by added into animal feed.

Species	Use level (Decoquinatate)	Amount of decoquinatate Powder	Indication for Use
Cattle	27.2 grams of decoquinatate per ton of feed.	0.5 kg per ton of feed	For the prevention of coccidiosis in cattle (including ruminating and non-ruminating calves and veal calves) caused by <i>Eimeria bovis</i> and <i>E.zuernii</i> .
Chicken (broiler)	27.2 grams of decoquinatate per ton of feed.	0.5 kg per ton of feed	For the prevention of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. mivati</i> , <i>E.acervulina</i> , <i>E. maxima</i> and <i>E. brunetti</i> .
Sheep	12.9 to 90.8 grams decoquinatate per ton of feed. Provide 22.7 milligrams per 100 pounds of body weight (0.5 milligrams per kilogram) per day. Feed for at least 28 days	0.2-1.5kg per ton of feed	For the prevention of coccidiosis caused by <i>Eimeria ovinoidalis</i> , <i>E. parva</i> , <i>E. bakuensis</i> , and <i>E. crandallis</i> .
	90.9 – 535.7 grams decoquinatate per ton of feed. Provide 22.7 milligrams per 100 pounds of body weight (0.5 milligrams per kilogram) per day. Feed for at least 28 days	1.5-8.9 kg per ton of feed	
Goats	12.9 to 90.8 grams decoquinatate per ton of feed. Provide 22.7 milligrams per 100 pounds of body weight (0.5 milligrams per kilogram) per day. Feed for at least 28 days	0.2-1.5kg per ton of feed	For the prevention of coccidiosis caused by <i>Eimeria christenseni</i> and <i>E.ninakohlyakimovae</i> .
	90.9 – 535.7 grams decoquinatate per ton of feed. Provide 22.7 milligrams per 100 pounds of body weight (0.5 milligrams per kilogram) per day. Feed for at least 28 days	1.5-8.9 kg per ton of feed	

PHARMACODYNAMIC (Mode of Action): Decoquinatate is a 4-hydroxyquinoline antiprotozoal compound active against *Eimeria* spp. and *Toxoplasma* spp. Decoquinatate inhibits the development of coccidia in the small intestine in the early part of the infective cycle, resulting in lower morbidity and mortality. The exact mode of action is unknown.

PHARMACOKINETICS: The drug is administered by the oral route, the main site of action being within the gastro-intestinal tract. It is poorly absorbed by the target species and is largely eliminated in faeces unchanged. Consequently, tissue residues are low and deplete rapidly with time. Recovery of the administered material via excretion is essentially complete.

CONTRAINDICATIONS: Do not feed to laying hens. Do not feed to cows, goats or sheep producing milk for human consumption.

WARNINGS AND PRECAUTIONS:

Not for human use, for animal feed only. Avoid inhaling dust during mixing. Avoid contact of product with skin, eyes and clothing. Wash thoroughly after handling.

INTERACTIONS WITH OTHER MEDICAMENTS: Do not use in feeds containing Bentonite.

PREGNANCY AND LACTATION: Do not feed to laying hens. Do not feed to cows, goats or sheep producing milk for human consumption. May be used in pregnant animals.

UNDESIRABLE EFFECTS: None known.

OVERDOSE AND TREATMENT: Overdose is unlikely with in-feed medication. Decoquinatate dosages of 4 mg/kg bodyweight in sheep and lambs and 6 mg/kg bodyweight in calves have been found to be well tolerated and without observable side effects.

STORAGE: Store below 25°C away from sunlight in dry conditions. The bag should be closed securely after use.

DOSAGE FORM: Oral Powder.

PACKAGING: Each lot is packaged into 25 kg multi-walled kraft bags.

DISPOSAL OF CONTAINER: Shake and empty contents into feed. Dispose of empty bag in the rubbish.

WITHHOLDING PERIODS: Cattle, Goats, Sheep and Chicken (broiler)-Zero days

PROPOSED SHELF LIFE AFTER RECONSTITUTION IN FEED: 24 hours

MANUFACTURED BY:

Zoetis Inc.
400 State Street
Chicago Heights,
Illinois, 60411 USA

Date of Revision: 29 April 2016

MARKETING AUTHORISATION HOLDER:

Pahang Pharmacy Sdn Bhd
Lot 5979, Jalan Teratai
5 1/2 Mile off Jalan Meru,
41050 Klang Selangor D.E. MALAYSIA

Malaysian Registration Number: MAL15125064HA

zoetis[™]

Batch:

Manufacturing Date:

Expiry: