



	Dogs	Cats	Pigs	Pigs	Cattle	Calves
Dose rate (mg/kg bw)	5	5	2.5	5	5	5
Route of administration	sc	sc	im	im	iv	sc
Tmax (h)	0.5	2	2	2	-	1.2
Cmax (µg/ml)	1.8	1.3	0.7	1.6	-	0.73
AUC (µg ·h/ml)	-	-	6.6	15.9	7.11	3.09
Terminal half-life (h)	-	-	13.12	8.10	-	2.34
Elimination half-life (h)	4.4	6.7	7.73	7.73	2.2	-
F (%)	-	-	95.6	-	-	-

Indication

Calves

Treatment of infections of the respiratory tract caused by enrofloxacin susceptible strains of *Pasteurella multocida*, *Mannheimia haemolytica* and *Mycoplasma* spp.

Treatment of infections of the alimentary tract caused by enrofloxacin susceptible strains of *Escherichia coli*.

Treatment of septicaemia caused by enrofloxacin susceptible strains of *Escherichia coli*

Treatment of acute mycoplasma-associated arthritis due to enrofloxacin susceptible strains of *Mycoplasma bovis*.

Pigs

Treatment of infections of the respiratory tract caused by enrofloxacin susceptible strains of *Pasteurella multocida*, *Mycoplasma* spp. and *Actinobacillus pleuropneumoniae*.

Treatment of infections of the alimentary tract caused by enrofloxacin susceptible strains of *Escherichia coli*.

Treatment of septicaemia caused by enrofloxacin susceptible strains of *Escherichia coli*

Dogs

Treatment of infections of the alimentary, respiratory and urogenital tracts (including prostatitis, adjunctive antibiotic therapy for pyometra), skin and wound infections, otitis (externa/media) caused by enrofloxacin susceptible strains of *Staphylococcus* spp., *Escherichia coli*, *Pasteurella* spp., *Klebsiella* spp., *Bordetella* spp., *Pseudomonas* spp. and *Proteus* spp.

Cats

Treatment of infections of the alimentary, respiratory and urogenital tracts (as adjunctive antibiotic therapy for pyometra), skin and wound infections, caused by enrofloxacin susceptible strains of, e.g.: *Staphylococcus* spp., *Escherichia coli*, *Pasteurella* spp., *Klebsiella* spp., *Bordetella* spp., *Pseudomonas* spp. and *Proteus* spp.

Recommended dose

Calves

5 mg of enrofloxacin/kg bw, corresponding to 1 ml/10 kg bw, once daily for 3-5 days

Acute mycoplasma-associated arthritis due to enrofloxacin susceptible strains of *Mycoplasma bovis*: 5 mg of enrofloxacin/kg bw, corresponding to 1 ml/10 kg bw, once daily for 5 days.

The product can be administered by slow intravenous or subcutaneous administration. No more than 10 ml should be administered at one subcutaneous injection site.

Pigs

2.5 mg of enrofloxacin/kg bw, corresponding to 0.5 ml/10 kg bw, once daily by intramuscular injection for 3 days.

Alimentary tract infection or septicaemia caused by *Escherichia coli*: 5 mg of enrofloxacin/kg bw, corresponding to 1 ml/10 kg bw, once daily by intramuscular injection for 3 days.

In pigs, the injection should be made in the neck at the ear base. Not more than 3 ml should be administered at one intramuscular injection site.

Dogs and cats

5 mg of enrofloxacin/kg bw, corresponding to 1 ml/10 kg bw, once daily by subcutaneous injection for up to 5 days.

Treatment may be initiated with injectable product and maintained with enrofloxacin tablets. Duration of treatment should be based on the duration of treatment approved for the appropriate indication in the product information of the tablet product.

PRODUCT INFO

BLUE #:	700928APB	Product Name:	Baytril 5% Injectable Solution
Item Code:	90208347	Component:	Package Inserts and Leaflets
Product Code:	N/A	Pack Size:	100 ml
Previous Item Code	90196831		

ARTWORK INFO

Template:	180(W) x 210(D) mm	Packaging Spec(s):	N/A
Barcodes/Type:	Code 128: 90208347	Add. Info:	N/A
		Minimum Core Data Point Size:	5.94 pt
GTIN:	GTIN Not Required	Proof #	P1b
		By/Date	TYA 31-MAR-2022
 			
NON PRINTING BLACK			

ELANCO ARTWORK LEGEND v16