

SIZE : 130X190mm

Controlled Medicine
Veterinary Use Only

Avante Ceftiofur 50mg/ml Sterile Powder

1. Brand or Product Name

2. Name and Strength of Active Substance(s)

Avante Ceftiofur Sodium Sterile Powder

Each mL of the resulting solution contains ceftiofur sodium equivalent to 50 mg ceftiofur.

3. Product Description

Yellow sterile powder.

4. Pharmacodynamics/Pharmacokinetics/Environmental Properties

Pharmacodynamics :

Ceftiofur is a 3rd generation cephalosporin. Cephalosporins have a bactericidal mechanism of action similar to that of other β -lactam antibiotics. They disrupt bacterial cell wall synthesis, targeting the penicillin-binding proteins (PBPs) located in the bacterial cell wall.

Ceftiofur and to a lesser extent its main metabolite, show activity against a variety of gram-negative and gram-positive bacteria in a time-dependent manner.

Pharmacokinetics :

Ceftiofur was poorly absorbed after oral administration and rapidly absorbed after intramuscular administration. Peak plasma concentrations of approximately 6 μ g equivalents/mL were attained approximately 30 minutes after intramuscular administration of 1 mg/kg bw/day to calves and pigs. In in vivo studies in rats and cattle and in in vitro studies with liver extracts from rats, pigs, cattle and chickens ceftiofur was rapidly metabolised to (initially) desfuroylceftiofur and furoic acid. Maximum blood concentrations of ceftiofur and its metabolites were achieved within 0.5 and 2 hours after dosing. No unmetabolised ceftiofur was detected in blood from 2 to 4 hours after dosing. Blood concentrations declined biphasically.

More than 95% of the intramuscularly administered dose was excreted within 24 hours of administration; 60 to 80% of this was in the urine and the remainder in the faeces. Most of the excreted material was desfuroylceftiofur and desfuroylceftiofur cysteine disulphide with small amounts of unmetabolised ceftiofur.

5. Indication

Cattle: Avante Sterile Powder is indicated for treatment of bovine respiratory disease (shipping fever, pneumonia) associated with Mannheimia haemolytica, Pasteurella multocida and Histophilus somni.

Avante Sterile Powder is also indicated for treatment of acute bovine interdigital necrobacillosis (foot rot, pododermatitis) associated with Fusobacterium necrophorum and Bacteroides melaninogenicus.

Swine: Avante Sterile Powder is indicated for treatment/control of swine bacterial respiratory disease (swine bacterial pneumonia) associated with Actinobacillus (Haemophilus) pleuropneumoniae, Pasteurella multocida, Salmonella choleraesuis and Streptococcus suis.

6. Recommended Dosage / Mode of Administration

Cattle: Administer to cattle by intramuscular or subcutaneous injection at the dosage of 0.5 to 1.0 mg ceftiofur per pound (1.1 to 2.2 mg/kg) of body weight (1-2 mL reconstituted sterile solution per 100 lbs body weight). Treatment should be repeated at 24-hour intervals for a total of three consecutive days.

Additional treatments may be given on days four and five for animals which do not show a satisfactory response (not recovered) after the initial three treatments. Selection of dosage (0.5 to 1.0 mg/lb) should be based on the practitioner's judgement of severity of disease (i.e., for respiratory disease, extent of elevated body temperature, depressed physical appearance, increased respiratory rate, coughing and/or loss of appetite; and for foot rot, extent of swelling, lesion and severity of lameness).

Swine: Administer to swine by intramuscular injection at the dosage of 1.36 to 2.27 mg ceftiofur per pound (3.0 to 5.0 mg/kg) of body weight (1 mL of reconstituted sterile solution per 22 to 37 lbs body weight). Treatment should be repeated at 24-hour intervals for a total of three consecutive days.

7. Contraindications

Do not use in animals previously found to be hypersensitive to ceftiofur and other β -lactam antibiotics. Do not use in case of known resistance to the active substance.

0000000/200915

8. Warnings and Precautions

Special precautions for use in animals:

Ceftiofur should be reserved for the treatment of clinical conditions which have responded poorly, or are expected to respond poorly, to more narrow spectrum antimicrobials.

Official and national antimicrobial policies should be taken into account when the product is used.

Use of the product should be based on susceptibility testing of the bacteria isolated from the animal.

If this is not possible, therapy should be based on local (regional, farm level) epidemiological information about susceptibility of the target bacteria.

Use of the product deviating from the instructions given in the SPC may increase the prevalence of bacteria resistant to ceftiofur and may decrease the effectiveness of treatment with other cephalosporins, due to the potential for cross resistance.

In the pig, particular care must be taken to avoid injection into the fat tissue.

Avoid repeated injection at the same site.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Penicillins and cephalosporins may cause hypersensitivity (allergy) following injection, inhalation, ingestion or skin contact. Hypersensitivity to penicillins may lead to cross reactions to cephalosporins and vice versa.

Allergic reactions to these substances may occasionally be serious.

Do not handle this product if you are sensitised or if you have been advised not to work with such preparations.

Handle this product with great care to avoid exposure, taking all recommended precautions.

If you develop symptoms following exposure such as a skin rash, you should seek medical advice and show the doctor this warning. Swelling of the face, lips or eyes or difficulty with breathing are more serious symptoms and require urgent medical attention.

Wash hands after use.

Keep out of children/ Jauhi ubat daripada kanak-kanak.

9. Interactions with Other Medicaments

None known

10. Statement on usage during pregnancy and lactation

Laboratory studies in the rat have not produced any evidence of teratogenic, foetotoxic or maternotoxic effects.

The safety of the veterinary medicinal product has not been assessed during pregnancy and lactation in cattle or pigs. Use only according to the benefit/risk assessment by the responsible veterinarian.

11. Adverse Effects/Undesirable Effects

Use of this product may cause transient local discomfort at the site of the injection.

Hypersensitivity reactions can occasionally occur. In the case of an allergic reaction, treatment should be stopped.

In pigs, local irritation at the injection site may occur and can persist for 5 days or more.

12. Overdose and Treatment

In cattle, no evidence of systemic toxicity was observed following administration of an overdose.

In pigs, no evidence of systemic toxicity was observed following the administration of doses of up to 8 times the recommended dose, administered daily by intramuscular injection for 15 days.

13. Withdrawal Period(s)

Swine (pig):

Meat: 4 days pre-slaughter withdrawal period when used according to label directions.

Cattle:

Meat: 4 days pre-slaughter withdrawal period when used according to label directions.

Milk: zero days

14. Incompatibility data

The injection is incompatible in alkaline pH, with oxidizing agents and heavy metal ions.

15. Dosage Forms and packaging available

Box containing 1 vial of injection powder 1g and 1 vial of 20ml sterile water for injection

Box containing 1 vial of injection powder 4g and 1 vial of 80ml sterile water for injection

16. Storage Condition

Store unconstituted product at controlled room temperature below 30°C.

Store reconstituted product at controlled room temperature below 30°C for up to 12 hours.

Protect from light.

17. Marketing authorization holder

Nutri Pharmax Sdn Bhd

6, Lorong Serunai 1, Valdor Sungai Bakap, 14200 Sungai Jawi, Pulau Pinang, Malaysia.

18. Date of Revision of Package Insert

23 November 2020

TO BE PRESCRIBED AND TREATED WITH BY REGISTERED VETERINARY SURGEONS ONLY

Manufactured by
 **KBNP, INC.**
235-9 Chusaro, Sinam, Yesan,
Chungnam, Korea

Distributed by
 **Nutri Pharmax Sdn.Bhd.**
6, Lorong Serunai 1, Valdor Sungai Bakap,
14200 Sungai Jawi, Pulau Pinang, Malaysia
Tel: +604-582 9377