

Linco Spectin™

44 g/kg Premix

Lincomycin 22 g/kg
Spectinomycin 22g/kg

For animal use only
Controlled Medicine/Ubat Terkawal
Keep out of reach of children
Jauhi ubat dari kanak-kanak

Product Description: A grayish white to light brown powder which has a cereal odour.

Pharmacology: Linco-Spectin 44g/kg contains the antibiotics lincomycin and spectinomycin. Lincomycin is a lincosamide antibiotic and is produced by *Streptomyces lincolnensis*. It is bacteriostatic and is primarily active against Gram-positive bacteria (both aerobic and anaerobic), Gram-negative anaerobic bacteria and mycoplasmas.

The mode of action is inhibition of protein synthesis at the ribosomal 50S sub-unit level. Lincomycin has about 50% systemic bioavailability by the oral route in pigs. It is primarily excreted in the faeces as both parent compound and metabolites with large biliary contribution. Spectinomycin is an aminocyclitol antibiotic and is produced by *Streptomyces spectabilis*. It is bacteriostatic and is primarily active against Gram negative bacteria. Its mode of action is inhibition of protein synthesis at the ribosomal 30S sub-unit level. Spectinomycin is not well absorbed by the oral route and the vast majority (>90%) is retained in the gut and excreted in the faeces. Lincomycin and spectinomycin have been shown to be synergistic *in vitro* against anaerobes implicated in the pathogenesis of swine dysentery.

Indication: For the control and treatment of swine dysentery caused by *Serpulina hyodysenteriae* associated with Fusobacterium, Bacteroides, Clostridium and/or Campylobacter spp. sensitive to the combination lincomycin and spectinomycin.

For the control and treatment of enteritis in pigs caused by *Escherichia coli* and Salmonella spp. sensitive to the combination lincomycin and spectinomycin. The product has been shown to be particularly effective against complicated or mixed enteric infections involving the above organisms.

For the control and treatment of enteritis associated with *Lawsonia intracellularis* (ileitis) in pigs.

As an aid in the control of mycoplasmal (enzootic) pneumonia in pigs.

For the treatment of Mastitis, Metritis, Agalactiae (MMA) syndrome of bacterial origin sensitive to the combination of lincomycin and spectinomycin in sows.

Recommended Dose: For incorporation into dry feed.

Treatment and control of enteric disease:

Treatment: 2 kg product/tonne feed for 3 weeks or until clinical signs disappear.

Control: 1-2 kg product/tonne feed over period of risk.

Treatment of MMA: 1-2 kg product/tonne feed for 5-10 days prior to farrowing and 2-3 weeks post-farrowing.

Aid in the control of mycoplasmal pneumonia : 1-2 kg product per tonne of feed and feed daily throughout the period of risk.

To ensure thorough dispersion of the product it should first be mixed with a suitable quantity of feed before incorporation into the final mix. The product can be incorporated into pelleted feed at a processing temperature not exceeding 70°C.

Route of Administration: For oral administration only.

Contraindication: Do not allow horses, ruminating animals, guinea pigs, hamsters or rabbits access to feeds containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects.

Warning and Precautions:

Special warnings for each target species:

Loosening of faeces and/or mild swelling of the anus may occur; this is usually transient. On rare occasions, mild irritability and reddening of skin may occur. These conditions are usually self-correcting within five to eight days without discontinuing therapy. If clinical signs are not improved during the first 10 days of medication, discontinue treatment and redetermine the diagnosis.

Special precautions for use:

(i) Special precautions for use in animals

When an antibiotic treatment programme is prescribed, consideration should be given to ensuring good management practices are also in place.

(ii) Special precautions to be taken by the person administering the veterinary medicinal product to animals

Persons known to be hypersensitive (allergic) to lincomycin, spectinomycin or soybean millfeed should not handle this product.

Care should be taken not to inhale any dust. The wearing of approved dust masks (either a disposable half-mask respirator conforming to European Standard EN 149 or a non-disposable respirator to European Standard EN 140 with a filter to EN 143) and safety glasses is recommended during the handling and mixing of this product.

Skin contact should be avoided.

Wash hands and any exposed skin with soap and water immediately after use.

Interaction with Other Medicinal Product: No negative interactions known. Linco-Spectin has been shown to be clinically compatible with salinomycin.

Pregnancy and Lactation: No restriction

Side Effects

Loosening of faeces and/or mild swelling of the anus may occur; this is usually transient. On rare occasions, mild irritability and reddening of skin may occur. These conditions are usually self-correcting within five to eight days without discontinuing therapy.

Symptoms and Treatment of Overdose: Stop medication and to seek for veterinarian advice if there is sign of overdosage.

Storage Condition: Store below 25 °C. Store opened bag in a dry place to prevent caking.

Withdrawal Period: Animals must not be slaughtered for human consumption during treatment. Pigs may be slaughtered for human consumption only after 2 days after last dose.

Packaging available: 20 kg per bag.

Disposal Of Containers: Any unused product or waste material should be disposed of in accordance with national requirements.

Phibro, Phibro logo and Linco-Spectin are trademarks owned by or licensed to Phibro Animal Health Corporation or its affiliates.

Manufactured by:

Phibro Suzhou Manufacturing Co., Ltd.
No 690, Jian Lin Road, Suzhou New District, Jiangsu 215151, P.R. China.

Registered by:

PHIBRO CORPORATION (M) SDN BHD
311, Level 3, Menara Axis
2, Jalan 51A/223, Seksyen 51A
46100 Petaling Jaya, Selangor, Malaysia

MAL16010007HA

LOT No.:

MFG. Date:

EXP. Date:

871-8033-02

Phibro
ANIMAL HEALTH CORPORATION®

Z Reference No.	FG Material No.	Description	Colors
NA	NA	LincoSpectin 44 20KG (MYS) Label	
Specification / DWG	Dimensions	Country	
NA	200 x 280 mm	MYS	
MPD No.	Date	Additional Information	
12916CR3	08/21/25		BLACK PMS 166 C