

Veterinary Use Only

Tolcoxin

Toltrazuril 2.5% w/v Oral Solution

1. Brand or Product Name / Name and Strength of Active Substance(s)

Tolcoxin Toltrazuril 2.5% w/v Oral Solution, Each L contains 25g toltrazuril

2. Product Description : Colorless or pale yellow liquid.**3. Pharmacodynamics/Pharmacokinetics/Environmental Properties****Pharmacodynamics :**

Toltrazuril is a triazinon derivative. It acts against coccidia of the genus Isospora. It is acting against all intracellular development stages of coccidia of the merogony (asexual multiplication) and gamogony (sexual phase). All stages are destroyed, thus the mode of action is coccidiocidal.

Pharmacokinetics :

In poultry, Toltrazuril is absorbed at a rate of at least 50%. The highest concentrations are found in the liver. The active substance is rapidly metabolised in poultry. The main metabolite is characterised as a sulfone derivative. About 1 week after the last dose this metabolite represents by far the most relevant residue in animals.

4. Indication

For the treatment of Coccidiosis in Broilers and Broiler Breeders.

Toltrazuril is effective against *Eimeria acervulina*, *E. brunetti*, *E. maxima*, *E. necatrix*, *E. tenella*, *E. mitis*.

5. Recommended Dosage / Mode of Administration

Oral: Via drinking water

1ml of Tolcoxin per 1L of the drinking water for 2 consecutive days. Alternatively, it can be added at the higher level (3ml per 1L) for 8 hours per day for 2 successive days.

6. Contraindications

Do not use in known cases of hypersensitivity to active ingredient.

7. Warnings and Precautions

- Keep the withdrawal period.
- Do not exceed the recommended dosage.
- Use by veterinarian's indication, and veterinary use only.
- Read full directions before use.
- Do not use in layers producing eggs for human consumption.
- In case of accidental eye exposure, flush immediately with water for 15 minutes and consult a physician if irritation persists.
- In case of accidental skin exposure, wash with soap and water.
- Keep out of reach of children/ Jauhi ubat daripada kanak-kanak

8. Interactions with Other Medicaments

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other products.

9. Statement on usage during pregnancy and lactation

- Must not be used during pregnancy and lactation.
- Do not use in laying birds producing eggs for human consumption.

10. Adverse Effects/Undesirable Effects

If suspected adverse reactions occur, treatment should be discontinued and to inform veterinarian/ local authority.

11. Overdose and Treatment

A three - five fold overdose is readily tolerated without any symptoms. If the recommended dose is exceeded beyond 3 - 5 times there is a decrease in water intake. Stop medication and to seek for veterinarian advice If there is sign of overdose.

12. Withdrawal Period(s)

Meat: 18 days.

Do not use in birds producing eggs for human consumption.

13. Storage conditions

Store in a cool and dark place with dry condition below 30°C.

14. Dosage Forms and packaging available

1L

15. Shelf Life (after first opening of container) 24 hours.**16. Shelf Life (after reconstitution or dilution) 24 hours.****17. Name and Address of manufacturer/marketing authorization holder**

Nutri Pharmax Sdn Bhd

6, Lorong Serunai 1, Valdor Sungai Bakap, 14200 Sungai Jawi, Pulau Pinang, Malaysia.

18. Date of Revision of Package Insert : 04 March 2019

Change to new address



Manufactured by

KBNP, INC.

235-9 Chusaro, Sinam, Yesan,
Chungnam, Korea



Distributed by

Nutri Pharmax Sdn.Bhd.

6, Lorong Serunai 1, Valdor Sungai Bakap,
14200 Sungai Jawi, Pulau Pinang, Malaysia
Tel: +604-582 9377