

SAHARA

Tilmicosin 10% w/w Powder

1. Brand or Product Name / Name and Strength of Active Substance(s)

Sahara Tilmicosin 10% w/w Powder

Each kg contains tilmicosin (as Tilmicosin phosphate) 100g

2. Product Description : Yellow to brown powder.

3. Pharmacodynamics/Pharmacokinetics/Environmental Properties

Pharmacodynamics :

Tilmicosin is a semi-synthetic antibiotic of the macrolide group, and is believed to affect bacterial protein synthesis through binding to 50S ribosomal subunits and inhibiting peptide formation. It is bacteriostatic but at high concentrations it may be bactericidal.

Tilmicosin has an in-vitro antibacterial spectrum that is predominantly Gram-positive with activity against certain Gram-negative micro-organisms. Activity against several mycoplasma species has also been demonstrated. Cross resistance between tilmicosin and other macrolide antibiotics has been observed.

Pharmacokinetics :

Absorption: When administered to pigs via the oral route at a dose of 400 ppm in the feed (equivalent to approximately 21.3 mg/kg/day), tilmicosin moves rapidly out of the serum into areas of low pH. The highest concentration in the serum ($0.23 \pm 0.08 \mu\text{g/ml}$) was recorded on day 10 of medication, but concentrations above the limit of quantification ($0.10 \mu\text{g/ml}$) were not found in 3 out of 20 animals examined.

Lung concentrations increased rapidly between days 2 and 4, but no significant changes were obtained following four days of dosing. The maximum concentration in lung tissue ($2.59 \pm 1.01 \mu\text{g/ml}$) was recorded on day 10 of medication.

Distribution: Following oral administration, tilmicosin is distributed throughout the body, but especially high levels are found in the lung and in lung tissue macrophages. It is also distributed in the liver and kidney tissues.

Biotransformation: Several metabolites are formed, the predominant one being identified as T1.

However the bulk of the tilmicosin is excreted unchanged.

Elimination: Following oral administration, tilmicosin is excreted mainly via the bile into the faeces, but a small proportion is excreted via the urine.

4. Indication

Pigs

Prevention and treatment of respiratory disease caused by *Actinobacillus pleuropneumoniae*, *Mycoplasma hyopneumoniae*, *Pasteurella multocida* and other organisms sensitive to tilmicosin.

5. Recommended Dosage / Mode of Administration

Administer in the feed at a dose of 8 to 16 mg/kg bodyweight/day of tilmicosin (equivalent to 200 to 400 ppm in the feed) for a period of 15 to 21 days.

Inclusion rate in feed : 2~4kg of Sahara (100g tilmicosin /kg) per ton of feed for 15-21 days.

6. Contraindications

- The safety of tilmicosin has not been established in boars used for breeding purposes.
- Horses must not be allowed access to feeds containing tilmicosin. Horses fed with tilmicosin medicated feeds may present signs of toxicity with lethargy, anorexia, reduction of feed consumption, loose stools, colic, distension of the abdomen and death.
- Do not use in case of hypersensitivity to tilmicosin or to any of the excipients.

7. Warnings and Precautions

- For animal treatment only.
- In very rare cases, feed intake may decrease (including feed refusal) in animals receiving medicated feed. This effect is transient.
- Keep out of reach of children/Jauhi ubat daripada kanak-kanak.

Operator warnings

- Tilmicosin may induce irritation. Macrolides, such as tilmicosin, may also cause hypersensitivity (allergy) following injection, inhalation, ingestion or contact with skin or eye. Hypersensitivity to tilmicosin may lead to cross reactions to other macrolides and vice versa. Allergic reactions to these substances may occasionally be serious and therefore direct contact should be avoided.
- To avoid exposure during preparation of the medicated feed, wear overalls, safety glasses, impervious gloves and wear either a disposable half mask respirator conforming to European Standard EN149 or a non-disposable respirator to European Standard EN140 with a filter to EN143. Do not eat, drink or smoke when handling this product. Wash hands after use.
- In the case of accidental ingestion, wash out mouth immediately with water and seek medical advice. In the event of accidental skin contact, wash thoroughly with soap and water. In case of accidental eye contact, flush the eyes with plenty of clean, running water.
- Do not handle the product if you are allergic to ingredients in the product.
- If you develop symptoms following exposure, such as skin rash, you should seek medical advice and show the physician this warning. Swelling of the face, lips and eyes or difficulty in breathing are more serious symptoms and require urgent medical attention.

8. Interactions with Other Medicaments

This veterinary medicinal product must not be mixed with other products.

9. Statement on usage during pregnancy and lactation

This veterinary medicinal product must not be used during pregnancy and lactation.

Do not use in laying birds producing eggs for human consumption.

Use only after a risk/benefit assessment by a veterinarian

10. Adverse Effects/Undesirable Effects

Occasionally, feed intake may decrease (including feed refusal) in animals receiving medicated feed.

This effect is transient.

11. Overdose and Treatment

No symptoms of overdose have been seen in pigs fed a ration containing levels of tilmicosin up to 80 mg/kg bodyweight (equivalent to 2000 ppm in the feed or ten times the recommended dose) for 15 days.

12. Withdrawal Period(s)

Pig - 21 days

13. Storage Conditions

Store in a room temperature below 30°C.

Protect from light.

14. Dosage Forms and packaging available

20kg

15. Name and Address of manufacturer/marketing authorization holder

Nutri Pharmax Sdn Bhd

6, Lorong Serunai 1, Valdor Sungai Bakap, 14200 Sungai Jawi, Pulau Pinang, Malaysia.

16. Date of Revision of Package Insert : 09 FEB 2021

Manufactured by
 **KBNP, INC.**
235-9 Chusaro, Sinam, Yesan,
Chungnam, Korea

Product Registration Holder
 **Nutri Pharmax Sdn Bhd**
6 Lorong Serunai 1, Valdor Sungai Bakap,
14200 Sungai Jawi, Pulau Pinang Malaysia
Tel/Fax: +604-5829377