

BALAMIN 0.1% INJECTION**Description and Composition**

BALAMIN 0.1% INJECTION is a red clear solution. Each ml contains 1000mcg Cyanocobalamine as active ingredient and 10mg benzyl alcohol as preservative.

Pharmacodynamics

B vitamins are not stored in the body to any great extent

Pharmacokinetics

The metabolism of cyanocobalamin is complex and is associated closely with that of folic acid and of ascorbic acid. It is important for maintenance of normal haemopoiesis, protection of the liver, maintenance of muscle tissue, healthy skin, brain and pancreatic metabolism.

Indication

Prevention and treatment of vitamin B12 deficiency in cattle and pigs.

Recommended dose:

Intravenous, intramuscular and subcutaneous.

Cattle and pigs: 0.01 to 0.02mg of cyanocobalamin per kg of body weight equivalent to 1 to 2ml of solution per 100kg body weight. These doses can be restarted each daily for 8 days.

Route of Administration

By intra-muscular injection or subcutaneous or intravenous injection.

Contraindication

Do not use in case of hypersensitivity to the active substance or to any of the excipients.

Warning and Precautions

Care should be taken to avoid accidental self-injection. If accidental self-injection occurs, seek medical advice immediately. In case of accidental ingestion, seek medical advice immediately. In case of accidental spillage onto skin or eye, wash the affected area immediately with clean running water. Seek medical attention if irritation persists.

Rinse containers thoroughly with water. Dispose of rinsings in slurry or dirty water. Dispose of rinsed containers in the farm refuse. Used containers should not be recycled.

When the container is broached for the first time, using the in-use-shelf-life which is specified on the package insert, the date on which any product remaining in the container should be discarded needs to be written in the space provided on the label.

Interaction with Other Medicinal Product

None known

Incompatibilities

None known

Pregnancy and Lactation

Not contra-indicated in pregnant or lactating animals.

Side Effects

None known

Symptoms and Treatment of Overdose

Overdosage is unlikely.

Storage Condition

Store below 25°C. Protect from light.

Shelf Life

3 years. Following withdrawal of the first dose, use the product within 24 hours. Any unused material should be discarded.

Withdrawal periods:

Cattle:

Meat and offal: zero days

Milk: zero days

Pigs:

Meat and offal: zero days

Packing

100ml

REVISED: 11.1.2018

	Manufactured by::
	Range Pharma Sdn Bhd 1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 E-mail:sales@rangepharma.com.my