

BACITRACIN ZINC 15% W/W GRANULE

ACTIVE CONSTITUENT: Each kilogram contains 150 grams bacitracin activity from bacitracin zinc.

PRODUCT DESCRIPTION: Light brown or tan granule.

MODE OF ADMINISTRATION: Oral administration by mixing BACITRACIN ZINC 15% W/W GRANULE with feed.

INDICATION:

Chicken: For the reduction of early mortality in chicks and prevention of necrotic enteritis caused by *Clostridium perfringens* susceptible to bacitracin in broiler chickens.

Swine: As an aid in the prevention and treatment of Bacterial Enteritis (Scours) (except Coliform Diarrhoea) in swine.

DOSAGE:

Chicken:

For the reduction of early mortality in chicks:

110 mg of bacitracin in to 1kg of complete feed and provide this as the sole ration for 5 -15 days.

For the prevention of necrotic enteritis in broiler chicken:

55 mg of bacitracin in to 1kg of complete feed and provide this as the sole ration continuously.

Swine:

As an aid in the treatment of Bacterial Enteritis (Scours):

110 mg of bacitracin in to 1kg of complete feed and provide this as the sole ration for 5 to 15 days.

As an aid in the prevention of Bacterial Enteritis (Scours):

55 mg of bacitracin in to 1kg of complete feed and provide this as the sole ration continuously.

PHARMACODYNAMICS&PHARMACOKINETIC:

Bacitracin Zinc has strong anti-bacteria activities to gram-positive bacteria and part of gram-negative bacteria. It has a strong effect to most G⁺, such as *Staphylococcus aureus* (including *Staphylococcus aureus* resistant to Penicillin), *streptococcus*, *streptococcus pneumoniae*, *enterococcus*, et al., also affect to some G⁻ such as *meningococcus*, influenza bacteria, still has some effect to helicoid and actinomycete. G⁻ bacillus resistant to it.

After oral application to rats, chickens and pigs, bacitracin is hardly absorbed from the gastrointestinal tract, and the distribution to organs and tissues is negligible. In rats, chickens and pigs, approximately 95% of an oral dose is excreted via faeces, and only 3% or less via urine. Bacitracin is metabolized to amino acids and smaller peptides via the main metabolite desamidobacitracin, which is microbiologically inactive. Main metabolites in faeces are bacitracin A, B1, B2, F, desamidobacitracin and catabolic peptides. In urine and bile only hydrolytic cleavage products (di – and tripeptides) are present.

WARNING AND PRECAUTION: Will irritate the eyes, nose and throat. Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. Do not inhale dust. When opening the container and mixing into feed, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves, goggles and disposable dust mask. If products in eyes, wash it out immediately with water. After each day's use, wash gloves, goggles and contaminated clothing. Pigs refusing to eat should be treated individually.

INTERACTION WITH OTHER MEDICINAL PRODUCT : Bacitracin zinc could combine use with penicillin, streptomycin, neomycin, polymyxin and etc. Do not use in feeds containing bentonite or other pellet binding agents.

SYMPTOMS AND TREATMENT OF OVERDOSE: If an overdose is suspected seek advice from a veterinarian.

PREGNANCY AND LACTATION: Cannot be use in the laying period.

STORAGE CONDITION: Store in cool (25°C) and dry place and avoid light. Close tightly.

CONTRAINDICATION: None Known

SIDE EFFECTS: None known.

WITHDRAWAL PERIOD: Meat Nil, Egg Nil.

PACKAGING: 25kg/bag.

REGISTRATION NUMBER:

MANUFACTURED BY:

LIFECOME BIOCHEMISTRY CO., LTD

Add.:No.19, Nanpu Ecological Industrial Park,
Pucheng, Fujian, P.R. China (353400)

MARKETING AUTHORISATION HOLDER:

PANFAST MARKETING (M) SDN BHD(Co No.:730518-D)

Add.:Penang Science Park, 1102, MK13, Lorong Perindustrian
Bukit Minyak 18, Kawasan Perindustrian Bukit Minyak, 14100
Simpang Ampat, Penang, Malaysia

DATE OF REVISION: JUN 2015