

# BACITRACIN METHYLENE DISALICYLATE 15% GRANULE

**ACTIVE CONSTITUENT:** Each kilogram contains 150 grams bacitracin activity from bacitracin methylene disalicylate.

**PRODUCT DESCRIPTION:** Light yellow to light brown granule.

**MODE OF ADMINISTRATION:** Oral administration by mixing BACITRACIN METHYLENE DISALICYLATE 15% GRANULE in feed.

## INDICATIONS AND RECOMMENDED DOSE:

| Species                          | Grams bacitracin per ton of feed | Indications for Use                                                                                                                                                                                                                                                                                                                                                                                                                                          |
|----------------------------------|----------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Growing/finishing swine          | 250                              | For control of swine dysentery associated with <i>Treponema hyodysenteriae</i> on premises with a history of swine dysentery but where signs of the disease have not yet occurred; or following an approved treatment of the disease condition.<br>Limitations: As the sole ration. Not for use in swine weighing more than 250 pounds. Diagnosis should be confirmed by a veterinarian when results are not satisfactory.                                   |
| Pregnant sows                    | 250                              | For control of clostridial enteritis caused by <i>Clostridium perfringens</i> in suckling piglets when fed to sows from 14 days through 21 days after farrowing on premises with a history of clostridial scours.<br>Limitations: Feed as the sole ration to sows from 14 days before through 21 days after farrowing. Diagnosis should be confirmed by a veterinarian when results are not satisfactory.                                                    |
| Broiler and replacement chickens | 50                               | As an aid in the prevention of necrotic enteritis caused by or complicated by <i>Clostridium</i> spp. or other organisms susceptible to bacitracin methylene disalicylate.<br>Limitations: Feed as the sole ration.                                                                                                                                                                                                                                          |
|                                  | 100-200                          | As an aid in the control of necrotic enteritis caused or complicated by <i>Clostridium</i> species or other organisms susceptible to bacitracin methylene disalicylate.<br>Limitations: Feed continuously as sole ration. Start at first clinical signs of disease, vary dosage based on severity of infection, administer continuously for 5 to 7 days or as long as clinical signs persist, then reduce medication to prevention level (50 grams per ton). |
| Growing turkeys                  | 200                              | As an aid in the control of transmissible enteritis in growing turkeys complicated by organisms susceptible to bacitracin methylene disalicylate.<br>Limitations: No limitations are included for this species when using this product in this amount.                                                                                                                                                                                                       |
| Quail                            | 200                              | For prevention of ulcerative enteritis in growing quail caused by <i>Clostridium colinum</i> susceptible to bacitracin methylene disalicylate.<br>Limitations: Feed continuously as the sole ration..                                                                                                                                                                                                                                                        |
| Feedlot beef cattle              | 70mg/head/day                    | Used as bacitracin methylene disalicylate in feed. For reduction in the number of liver condemnations due to abscesses.<br>Limitations: Administer continuously throughout the feeding period.                                                                                                                                                                                                                                                               |
|                                  | 250mg/head/day                   | Used as bacitracin methylene disalicylate in feed. For reduction in the number of liver condemnations due to abscesses.<br>Limitations: Administer continuously for 5 days to provide 250 mg bacitracin methylene per head per day, then discontinue for subsequent 25 days; repeat the pattern during the feeding period.                                                                                                                                   |

## PHARMACODYNAMICS/PHARMACOKINETICS:

Bacitracin is an antibiotic from the group of peptide antibacterial compounds. Bacitracin is produced by *Bacillus licheniformis*, and is a mixture of several closely related polypeptides, mainly consisting of bacitracin A, B1, B2, C and a small portion of bacitracin F. Bacitracin is mainly active against gram-positive microorganisms. Bacitracin inhibits peptidoglycan synthesis in bacteria by nonspecifically blocking phosphorylase reactions, some of which occur during cell wall synthesis. After oral application to rats, chickens and pigs, bacitracin is hardly absorbed from the gastrointestinal tract, and the distribution to organs and tissues is negligible. In rats, chickens and pigs, approximately 95% of an oral dose is excreted via faeces, and only 3% or less via urine. Bacitracin is metabolized to amino acids and smaller peptides via the main metabolite desamidobacitracin, which is microbiologically inactive.

**WARNINGS AND PRECAUTIONS:** Will irritate the eyes, nose and throat. Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. Do not inhale dust. When opening the container and mixing into feed, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves, goggles and disposable dust mask. If product in eyes, wash it out immediately with water. After each day's use, wash gloves, goggles and contaminated clothing.

**INTERACTION WITH OTHER MEDICINAL PRODUCT:** Do not use in feeds containing pellet binding agents.

**OVERDOSE AND TREATMENT:** If an overdose is suspected seek advice from a veterinarian.

**PACKAGING:** 25kg/bag.

**CONTRAINDICATIONS:** None known.

**SIDE EFFECTS:** None known.

**PREGNANCY AND LACTATION:** None known.

**WITHDRAWAL PERIOD:** NIL

**DOSAGE FORM:** In the form of granule.

**STORAGE CONDITION:** Store in a cool (25°C) and dry place. Avoid light and close tightly.

**DATE OF REVISION:** MAL DEC 2014

## MANUFACTURED BY:

Lifecome Biochemistry Co., Ltd

Add.:No.19, Nanpu Ecological Industrial Park, Pucheng, Fujian, P.R. China (353400)

## MARKETING AUTHORISATION HOLDER:

PANFAST MARKETING (M) SDN BHD (Co No.:730518-D)

Add.:Penang Science Park, 1102, MK13, Lorong Perindustrian Bukit Minyak 18, Kawasan Perindustrian Bukit Minyak, 14100 Simpang Ampat, Penang, Malaysia