

105mm

150mm

150mm



ACTIVE



MUSCLE RUB

Non-greasy muscular pain relief cream

Tiger Balm Active Muscle Rub Cream

Name and Strength of Active Substance(s)

Methyl Salicylate 15% w/w
Menthol 5% w/w
Camphor 3% w/w

Preservative
Germaben II 0.2%

Product Description
Tiger Balm Active Muscle Rub is a whitish cream with characteristic smell of Methyl Salicylate.

Pharmacodynamics
Camphor
Camphor is a stimulant, rubefacient, antipruritic and feebly antiseptic. When in combination with other ingredient like menthol, methyl salicylate, it becomes ideal for neuralgia and other painful areas.

Menthol
Menthol dilates the blood vessels causing a sensation of coldness, followed by an analgesic effect. Menthol also acts as a penetration enhancer, increasing the penetration of drugs when applied on the skin, to give a faster onset of action.

Methyl salicylate
Methyl salicylate is a salicylic acid derivative. Salicylates inhibit cyclooxygenase, thereby reducing the formation of prostaglandins, and cause platelet dysfunction. Methyl salicylate is used topically as a counter-irritant. Upon application, it is absorbed through the skin and is applied for the relief of pain in rheumatic conditions and painful muscles or joints.

Pharmacokinetics
Camphor
• Camphor is readily absorbed from all administration sites. Camphor is well absorbed after dermal exposure.
• It is hydroxylated in the liver to yield hydroxycamphor metabolites which are then conjugated with glucuronic acid and excreted in the urine.
• Camphor crosses the placenta

Menthol
• After absorption, menthol is excreted in the urine and bile as a glucuronide.

Methyl salicylate
• May be absorbed through intact skin.

- The absorption of topical salicylates is proportional to the surface area involved, duration of exposure, concentration and skin integrity.
- Absorption characteristics of salicylates vary with the dose, formulation and route of administration.
- Percutaneous absorption is enhanced by exercise, heat, occlusion, or disruption of the integrity of the skin or application to large areas of skin.
- Both the rate and extent of absorption increases after repeated application; increasing the bioavailability.
- Methyl salicylate is extensively metabolised to salicylic acid in the dermal and subcutaneous tissues after topical application.
- At therapeutic levels, the half life of salicylates is 2-4 hours. As salicylate level increases into the toxic range, the half-life can be greater than 18 hours.

Indication
Tiger Balm Active Muscle Rub is a specially formulated non-greasy cream. It provides temporary relief of aches and pains from sore muscles, sprains and strains for active people.

Recommended Dosage
Apply generously before exercise or whenever you have muscular pains. Gently massage on the affected area until completely absorbed. Repeat 3 to 4 times daily as necessary. For external application only.

Mode of Administration
Topical

Contraindications:

- Do not use on patients with aspirin or salicylate idiosyncrasy.
- Hypersensitivity to any of the ingredients in the product.
- Product can cause convulsions. This product should not be used on children below 2 years of age.
- Caution must be exercised when older children are treated. Avoid direct application into nostrils.
- Do not use during pregnancy or when breastfeeding.

Warnings and Precautions
WARNING:
• This product is contraindicated on children below 2 years of age.
• Caution must be exercised when older children are treated.
• If you are taking warfarin, please consult your doctor before using this product.

MAL16015047X



ACTIVE



MUSCLE RUB

Non-greasy muscular pain relief cream

- Can cause convulsion.
- Topical analgesic preparation containing methyl salicylate should be used with caution in patients at increased risk of developing salicylate adverse effects.
- Children suffering flu, chicken pox, or fever should avoid using this product because salicylates may induce Reyes Syndrome.

PRECAUTION:

- It is dangerous to place any camphor containing product into the nostril of children.
- A small amount applied this way may cause immediate collapse.
- Initially apply a small amount to a limited skin area and observe whether there is discomfort or itch before applying to a larger skin area. Discontinue use if irritation develops.
- For external use only; not to be taken.
- Avoid contact with the eyes and mucous membranes.
- Do not apply to open wound or damaged skin.
- Do not apply over large areas of the body.
- Do not use bandage or heating pad on the applied area.
- If symptoms persist, consult your doctor.

Under Caution:
This product contains methyl salicylate and when applied or rubbed on to the skin, can be absorbed through the skin into the blood. For patients taking warfarin, excessive application on to the skin for muscle or joint pains may increase the chances of bleeding.

Effects on Ability to Drive and Use Machines
None known

Interactions with Other Medicaments
This product contains methyl salicylate and when applied or rubbed on to the skin, can be absorbed through the skin into the blood. For patients taking warfarin, excessive application on to the skin for muscle or joint pains may increase the chances of bleeding.

Statement on Usage during Pregnancy and Lactation
This product can cause convulsions.

Adverse Effects/Undesirable Effects
This product can cause convulsions.

Headache, dizziness, nausea and vomiting, skin irritation, contact dermatitis, rash, itching, redness or swelling, burning or stinging sensation may occur.
May cause hypersensitivity/allergic reactions on some individuals with sensitive skin.

Overdose and Treatment
Symptoms of methyl salicylate overdose
Salicylate intoxication can occur after ingestion or topical application of methyl salicylate. Mild chronic salicylate intoxication, or salicylism, usually occurs only after repeated use of large doses. Salicylism can also occur following excessive topical application of salicylates. Symptoms include dizziness, tinnitus, deafness, sweating, nausea and vomiting, headache, and confusion, and may be controlled by reducing the dosage.
Symptoms of more severe intoxication or of acute poisoning following overdose include hyperventilation, fever, restlessness, ketosis, and respiratory alkalosis and metabolic acidosis. Depression of the CNS may lead to coma; cardiovascular collapse and respiratory failure may also occur.

Symptoms of camphor/menthol overdose
Ingestion of significant quantities is reported to cause symptom: severe abdominal pain, nausea, vomiting, vertigo, ataxia, drowsiness, and coma; instant collapse in infants after the local application of menthol/camphor to their nostrils.
Seizures may be the first clinical sign of severe toxicity of camphor; however, seizures are usually self-limited. Severe toxicity of camphor can result in delirium, visual hallucinations, cerebral edema, and status epilepticus. Systemic toxicity may include hypotension, tachycardia, respiratory failure, and death.

Treatment
The stomach should be emptied by gastric lavage or administration of oral activated charcoal. Fluid and electrolyte management is the mainstay of treatment with the immediate aim of correction of acidosis, hyperpyrexia, hypokalaemia and dehydration if present.
Any convulsions must be controlled first through supportive care including anticonvulsant therapy.

Storage Conditions
Store below 30°C.

Dosage Forms and Packaging Available
5g & 60g, Cream

Name and Address of Manufacturer
Manufactured under license from Haw Par Corporation Ltd., Singapore by Tiger Balm (M) Sdn. Bhd. No. 53, Jalan I-Park SAC 7, Taman Perindustrian I-Park SAC, 81400 Senai, Johor, Malaysia.

Name and Address of Product Registration Holder
Tiger Balm (M) Sdn. Bhd.
No. 53, Jalan I-Park SAC 7, Taman Perindustrian I-Park SAC, 81400 Senai, Johor, Malaysia.

Date of P.I.L : 13/11/24

MP1M0105 2

Front

Back

FONT SIZE: 5.25pt

Client	Haw Par Healthcare Pte Ltd			Person to contact	Shermaine / Hui Hui		Software Version	Adobe Illustrator CS6	
Artwork	Visibility Design Pte Ltd			Job No.	Q/HP/1024/6791		Date In	14 Nov 2024	
Revision Code	VD-01	File	Malaysia TB Active Muscle Rub 60g Insert (TBM)			Material Code	MP1M0105 2		
Last date of amendment: 8 Apr 2025									



TYPE FACES:

ENGLISH: HELVETICA NEUE