


APOTEX PRINTED PACKAGING MATERIAL MASTER

New Material Code: TBD	ECL Common Text #: N/A	Description: 163586 Deferiprone Film Coated Tablets 1000mg 50 Extended Content Label (ECL) Leaflet Malaysia Pharmforte Malaysia - PHM
SAP Ref #: N/A		
Old Material Code: N/A	C of A: PKGP-CA-LBL-LEAFLET	Change Control #: N/A
Pantone Colours: 032 C BLACK		DieLine ☒ COLOUR PERCENTAGE
Dimensions/Dieline #: 50.8 mm x 138.125 mm - 11 Panel	Minimum Font Size: 7 pt	Version No.: 1 Cycle No.: 2

Please note that customer approval signature represents full and complete regulatory review and responsibility for content, accuracy and completeness.

Page 1 of 1



STD 11 PANEL LEAFLET
MARCH/30/2010

FRONT LEAFLET

non-iron-loaded rats and rabbits have indicated that deferiprone is teratogenic and embryotoxic at doses at least as low as 25 mg/kg body weight. The potential risk for humans is unknown. Women of childbearing potential must be advised to avoid pregnancy due to the clastogenic and teratogenic properties of the medicinal product. These women should be counseled to take contraceptive measures and must be advised to immediately stop taking deferiprone if they become pregnant or plan to become pregnant.

Use in lactation
It is not known whether deferiprone is excreted in human milk. Deferiprone should not be used by nursing mothers.

Laboratory tests
It is recommended that serum ferritin concentrations or other assessments of body iron load be monitored regularly to assess the long-term effectiveness of the chelation regimen in controlling the body iron load. Interruption of therapy with deferiprone should be considered if serum ferritin measurements fall below 500 µg/L. Monitoring of plasma Zn²⁺ and supplementation in case of deficiency, is recommended.

HIV positive or other immune compromised patients
No data are available on the use of deferiprone in HIV positive or in other immune compromised patients. Given that deferiprone is associated with neutropenia and agranulocytosis, therapy in immune compromised patients should not be initiated unless potential benefits outweigh potential risks.

Neurological disorders
Neurological disorders have been observed in children treated with more than 2.5 times the maximum recommended dose for several years but have also been observed with standard doses of deferiprone. Prescribers are reminded that the use of doses above 100 mg/kg/day are not recommended. Deferiprone use should be discontinued if neurological disorders are observed.

Discoloration of urine
Patients should be informed that their urine may show a reddish/brown discoloration (chromaturia) due to the excretion of the iron-deferiprone complex.

CONTRAINDICATIONS
Ferriprox is contraindicated in patients who:
• Have demonstrated hypersensitivity to the active substance or any of the excipients.
• Have a history of recurrent episodes of neutropenia.
• Have a history of agranulocytosis.
• Are pregnant or breast-feeding.

DRUG INTERACTIONS
Interactions between deferiprone and other medicinal products have not been reported. However, since this compound binds to metallic cations, the potential exists for interactions between deferiprone and trivalent cation-dependent medicinal products such as aluminium-based antacids. Therefore, it is not recommended to concomitantly ingest aluminium-based antacids.

Due to the unknown mechanism of deferiprone-induced neutropenia, patients should not take medicinal products known to be associated with neutropenia or those that can cause agranulocytosis.

The safety of concurrent use of deferiprone and vitamin C has not been formally studied. Based on the reported adverse interaction that can occur between deferoxamine and vitamin C, caution should be used when administering deferiprone and vitamin C concurrently.

Body weight (kg)	Total daily dose (mg)	Number of 1000 mg tablets*		
		Morning	Midday	Evening
20	1500	0.5	0.5	0.5
30	2250	1.0	0.5	1.0
40	3000	1.0	1.0	1.0
50	3750	1.5	1.0	1.5
60	4500	1.5	1.5	1.5
70	5250	2.0	1.5	2.0

*number of tablets rounded to nearest half tablet

Doses above 100 mg/kg/day are not recommended because of the potentially increased risk of adverse reactions. Limited information is available on the use of deferiprone in children between 6 and 10 years of age, and very limited information is available on deferiprone use in children under 6 years of age.

It is not necessary to take deferiprone with food. However, it may be easier to remember to take doses with meals, and clinical experience suggests nausea may be minimized if deferiprone is taken with food.

SYMPTOMS AND TREATMENT OF OVERDOSAGE
No cases of acute overdose have been reported. However, neurological disorders (such as cerebellar symptoms, diplopia, lateral nystagmus, psychomotor slowdown, hand movements and axial hypotonia) have been observed in children who had been voluntarily prescribed more than 2.5 times the maximum recommended dose of 100 mg/kg/day for several years. The neurological disorders progressively regressed after deferiprone discontinuation.

In case of overdose, close clinical supervision of the patient is required.

STRENGTH AND DOSAGE FORMS AVAILABLE
Each tablet of Ferriprox contains 1000 mg deferiprone.

The tablets are white to off-white, capsule-shaped, scored, coated tablets. Engraved "APO" bisect "1000" on one side, plain on the other side. The tablets are scored and breakable in half.

Ferriprox is available in HDPE bottles of 50 tablets with child-resistant closures and a desiccant.

STORAGE CONDITIONS
Do not store above 30°C.
Keep out of the reach and sight of children.
Keep the bottle tightly closed in order to protect from moisture.

SHELF-LIFE
Do not use after the expiry date stated on the label.

MANUFACTURER
Apotex Inc. – Etobicoke Site
50 Steinerway Boulevard
Etobicoke, Ontario
Canada M9W 6Y3

MARKETED BY
Chiesi Farmaceutici S.p.A.
Via Palermo 26/A
43122 Parma
Italy

Revised 09/2019.

INSIDE LEAFLET

FERRIPROX® 1000 mg film-coated tablets
Deferiprone
Iron chelating agent
Oral Use

PHARMACOLOGY
Deferiprone is a bidentate ligand which binds to iron in a 3:1 molar ratio and removes iron from the body. Clinical studies have demonstrated that deferiprone is effective in promoting iron excretion and can lower serum ferritin levels and tissue iron stores in patients with transfusion-dependent thalassemia. The precise mechanism by which deferiprone is effective in promoting iron excretion and preventing the progress of iron accumulation is unknown.

Pharmacokinetics
Deferiprone is rapidly absorbed from the upper part of the gastro-intestinal tract. Peak serum concentration is reported to occur 45 to 60 minutes following a single dose in fasted

Ferriprox® 1000 mg Film-coated tablets Deferiprone

Oral use 50 tablets

Controlled Medicine
Ubat Terkawal

Chiesi
MAL16065019ACZ

One tablet contains 1000 mg deferiprone.
Read the package leaflet before use.

JAUHKAN UBAT DARI KANAK-KANAK
Do not store above 30°C.
Keep the bottle tightly closed in order to protect from moisture.
Medicinal product subject to medical prescription.

patients. This may be extended to 2 hours in fed patients. Following a dose of 25 mg/kg, lower peak serum concentrations have been detected in patients in the fed state (85 µmol/l) than in the fasting state (126 µmol/l), although there was no decrease in the amount of deferiprone absorbed in patients with food. Deferiprone is metabolised predominantly to a glucuronide conjugate. This metabolite lacks iron-binding capability because of inactivation of the 3-hydroxy group of deferiprone. Peak concentrations of the glucuronide occur 2 to 3 hours after administration of deferiprone.

In humans, deferiprone is eliminated mainly via the kidneys; 75% to 90% of the ingested dose is reported as being recovered in the urine in the first 24 hours, in the form of free deferiprone, the glucuronide metabolite and the iron-deferiprone complex. A variable amount of elimination into the feces has been reported. The elimination half-life in most patients is 2 to 3 hours.

INDICATION
Ferriprox is indicated for the treatment of iron overload in patients with thalassemia major when deferoxamine therapy is contra-indicated or inadequate.

ADVERSE REACTIONS
The following side effects were reported in patients taking Ferriprox in clinical trials:
Very Common (≥1/10): Nausea, vomiting, abdominal pain, joint disease.
Common (≥1/100 to <1/10): Low white blood cell count (agranulocytosis and neutropenia), Stomach upset, increased appetite, diarrhea, Pain, headache, back pain, fatigue, flu syndrome, swelling, Elevated liver enzyme (ALT), low number of blood platelets, decline in other blood cell counts.
Uncommon (≥1/1,000 to <1/100): Fever, bacterial infections, malaise, cyst.

Elevated liver enzymes (AST, GGT), depletion of minerals, Dizziness, sleepiness, decreased muscle activity, Taste loss, deafness, ear pain, Sore throat, Liver tenderness, jaundice, hepatitis, Anorexia (loss of appetite), excess gas, gastritis, Skin reactions (rash, itchiness, redness), Bone pain, muscle pain, leg cramps, epicondylitis, Loss of menstruation, frequent urination.

If the patient notices any side effects not mentioned in this leaflet, they should inform their doctor, pharmacist, or the local distributor immediately.

The most serious adverse reaction with deferiprone is agranulocytosis, which occurred in 1.1% of people participating in clinical trials (0.6 cases per 100 patient-years of treatment). The incidence of the less severe form of neutropenia is 4.9% (2.5 cases per 100 patient-years). This rate should be considered in the context of the underlying elevated incidence of neutropenia in thalassemia patients, particularly in those with hypersplenism.

Episodes of diarrhea, mostly mild and transient, have been reported in patients treated with Ferriprox. Gastrointestinal effects are more frequent at the beginning of therapy and in most patients, resolve within a few weeks without the need for discontinuation of treatment. Arthropathy events, which ranged from mild pain in one or more joints to severe arthritis with effusion and significant disability, have also been reported. Mild arthropathies are generally transient.

Neurological disorders (such as cerebellar symptoms, diplopia, lateral nystagmus, psychomotor slowdown, hand movements and axial hypotonia) have been observed in children who had been voluntarily prescribed more than 2.5 times the maximum recommended dose of 100 mg/kg/day for several years. Episodes of hypotonia, instability, inability to walk, and hypertonia with inability of limb movement, have been reported in children in the post-marketing setting with standard doses of deferiprone. The neurological disorders progressively regressed after deferiprone discontinuation.

PRECAUTIONS/WARNINGS
Neutropenia/Agranulocytosis
Deferiprone has been shown to cause neutropenia, including agranulocytosis. It is recommended that a patient's neutrophil count be monitored every week. In clinical trials this has been effective in identifying cases of neutropenia and agranulocytosis, and in allowing early interruption of deferiprone therapy. Under those circumstances, neutropenia and agranulocytosis resolved once therapy was withdrawn. If the patient develops an infection while on deferiprone, therapy should be interrupted and the neutrophil count monitored more frequently. Patients should be advised to report immediately to their physician any symptoms indicative of infection such as fever, sore throat and flu-like symptoms.

It is recommended that a management protocol as outlined below be in place prior to initiating any patient on deferiprone treatment.

Treatment with deferiprone should not be initiated if the patient is neutropenic. The

risk of agranulocytosis and neutropenia is higher, if the baseline absolute neutrophil count (ANC) is less than 1.5x10⁹/L.

In the event of neutropenia:
Instruct the patient to immediately discontinue deferiprone and all other medicinal products with a potential to cause neutropenia. The patient should be advised to limit contact with other individuals in order to reduce the risk of infection. Obtain a complete blood cell (CBC) count, with a white blood cell (WBC) count, corrected for the presence of nucleated red blood cells, a neutrophil count, and a platelet count immediately upon diagnosing the event and then repeat daily. It is recommended that following recovery from neutropenia, weekly CBC, WBC, neutrophil and platelet counts continue to be obtained for three consecutive weeks, to ensure that the patient recovers fully. Should any evidence of infection develop concurrently with the neutropenia, the appropriate cultures and diagnostic procedures should be performed and an appropriate therapeutic regimen instituted.

In the event of severe neutropenia or agranulocytosis:
Follow the guidelines above and administer appropriate therapy such as granulocyte colony stimulating factor, beginning the same day that the event is identified; administer daily until the condition resolves. Provide protective isolation and if clinically indicated, admit patient to the hospital.

Limited information is available regarding rechallenge. Therefore, in the event of neutropenia, rechallenge is not recommended. In the event of agranulocytosis, a rechallenge is contraindicated.

Renal or hepatic impairment and liver fibrosis
Currently there is no available information on patients with renal impairment and limited information on patients with hepatic impairment. Since deferiprone is eliminated mainly via the kidneys, there may be an increased risk of complications in patients with impaired renal function. Similarly, as deferiprone is metabolized in the liver, caution must be exercised in patients with hepatic dysfunction. Renal and

hepatic function should be monitored in this patient population during deferiprone therapy. If there is persistent increase in serum ALT, interruption of deferiprone therapy should be considered.

In thalassemia patients there is an association between liver fibrosis and iron overload and/or hepatitis C. Special care must be taken to ensure that iron chelation in patients with hepatitis C is optimal. In these patients careful monitoring of liver histology is recommended.

Cardiac function
A comparative study indicates that deferiprone has a cardio-protective effect at least as good as, if not better than, the standard treatment with subcutaneous deferoxamine.

Carcinogenicity/mutagenicity/effects on fertility
Deferiprone displayed clastogenic characteristics in *in vitro* and *in vivo*

non-iron-loaded and iron-loaded systems. Deferiprone was non-mutagenic in the bacterial reverse mutation assay. A comparative study of clastogenicity in the lymphocytes of patients with thalassemia treated with deferiprone or with deferoxamine indicated that deferiprone is not associated with a greater frequency of chromosomal aberrations than that observed during therapy with deferoxamine. No carcinogenicity studies have been conducted with deferiprone. In view of the genotoxicity results, a carcinogenic potential of deferiprone cannot be excluded.

Use in pregnancy
There is no adequate information on the use of deferiprone in pregnant women. Studies in animals have shown reproductive toxicity. Reproductive studies in