

SANBEACETIN® 600

Acetylcysteine

Effervescent Tablet

COMPOSITION

Each effervescent tablet contains:

Acetylcysteine 600 mg

PRODUCT DESCRIPTION

SANBEACETIN® 600 (Acetylcysteine Effervescent Tablet) is a round flat tablet, the colour is white with yellow spot and has orange odour. After dissolved in 200 mL, the solution is rather turbid, colourless and has orange odour with sweet-sour taste.

PHARMACODYNAMICS

Acetylcysteine has strong fluidification activity to mucus secretion and mucopurulent by depolymerization of mucoprotein acid complex and nucleic acid were as act as viscosity agent from mucoid component, sputum purulent and others secretions, it acts as anti flogistic and increase mucose regeneration speed.

PHARMACOKINETICS

Absorption

Bioavailability

Absorbed following oral administration, with peak plasma concentration achieved within 1 - 2 hours. Absorbed from the pulmonary epithelium.

Distribution

Extent: Crosses the placenta following oral or IV administration.

Plasma protein binding: 50 - 83%.

Metabolism

Deacetylated in the liver to cysteine.

Elimination

Elimination route: Principally (70%) nonrenal.

Half - life: 6.25 hours after oral administration.

INDICATIONS

SANBEACETIN® 600 (Acetylcysteine Effervescent Tablet) is indicated in the treatment of respiratory affections characterized by thick and viscous hypersecretions:

- acute bronchitis
- chronic bronchitis and its exacerbations
- pulmonary emphysema
- mucoviscidosis
- bronchieactasis.

RECOMMENDED DOSAGE

Adults:

1 effervescent tablet of **SANBEACETIN® 600** a day (preferably in the evening).

DIRECTIONS FOR USE

Dissolve 1 effervescent tablet into a glass of water (± 200 mL).

ROUTE OF ADMINISTRATION

Oral.

CONTRA-INDICATIONS

Patients with hypersensitivity to Acetylcysteine.

Contraindicated in children below two (2) years of age.

WARNING AND PRECAUTIONS

Patient with gastric, should be administered after food. Several animal and human researches showed no teratogenic and dangerous adverse effect to pregnant and nursing patient, however administer only at several condition needs and always controlled by physician.

INTERACTIONS WITH OTHER MEDICAMENTS

No cases of drug-drug interaction are described in literature for Acetylcysteine by oral use.

PREGNANCY AND LACTATION

Pregnancy: Category B

Lactation: Not known whether acetylcysteine is distributed into milk; caution advised.

ADVERSE REACTIONS

Hypersensitivity reactions have been reported in patients receiving Acetylcysteine, including bronchospasm, angiodema, rashes and pruritus. Other adverse effects reported with Acetylcysteine include nausea and vomiting, fever, syncope, sweating, arthralgia, blurred vision, disturbances or liver function.

Immune System Disorders:

Anaphylactic/anaphylactoid reaction

Skin and Subcutaneous Tissue Disorders:

Severe cutaneous adverse reactions (SCAR) e.g. erythema multiforme, Stevens-Johnson syndrom (SJS) and toxic epidermal necrolysis (TEN). In most of these cases reported at least one other drug was administered at the same time, which may have possibly enhanced the described mucocutaneous effects.

OVERDOSE AND TREATMENT

In the cases of accidental or non-accidental overdose, it may cause irritation, pyrosis, stomachache, nausea, vomiting and diarrhea. There are no serious side effects or toxic symptoms reported, even in patients treated with very high doses of acetylcysteine. However, it is advisable to consult the physician in any case of overdoses.

DOSAGE FORM AND PACKAGING AVAILABLE

TABLET in form of EFFERVESCENT TABLET

Box of 1 tube @ 10 effervescent tablets.

STORAGE

Keep in a tightly closed container at temperature below 30°C in a dry place.

Manufactured by:

PT CAPRIFARMINDO LABS.

Jl. Industri Cimoreme No. 8

Desa Cipeundeuy, Kecamatan Padalarang,

Kabupaten Bandung Barat, Indonesia

For product owner:

PT SANBE FARMA

Jl. Taman Sari No.10

Bandung, Indonesia

Date of Revision : May 06th, 2023

* BF 352 - EMY3

Country:
Malaysia

Packaging:
Insert

Dimension:
70 x 184 mm

Material:
HVS paper 60 g

Color: 
BLACK

Scale:
100%

REV. DATE : 26/05/23