

Package Insert

Noripharma Calcium Lactate 300mg Tablet

COMPOSITION

Calcium Lactate 300mg/tablet
Equivalent to Calcium 39mg

PRODUCT DESCRIPTION

White to off white, round shaped film coated tablet with breakline on one side.

PHARMACOLOGY

Calcium is absorbed mainly from the small intestine by active transport and passive diffusion. About one-third of ingested calcium is absorbed although this can vary depending upon dietary factors and the state of the small intestine, also absorption is increased in calcium deficiency during periods of high physiological requirement such as during childhood or pregnancy and lactation. 1,25-Dihydroxycholecalciferol (calcitriol), a metabolite of Vitamin D, enhances the active phase of absorption. Excess calcium is mainly excreted renally. Unabsorbed calcium is eliminated in the faeces, together with that secreted in the bile and pancreatic juice. Minor amounts are lost in the sweat, skin, hair and nails. Calcium crosses the placenta and is distributed into breast milk.

INDICATION

As a health supplement.

RECOMMENDED DOSAGE

Adults: Take 1-2 tablets each time, 2 times a day, with or after meal, or as directed by your pharmacist or doctor.

ROUTE OF ADMINISTRATION

Oral.

Package Insert

CONTRAINDICATIONS

Severe hypercalcaemia and hypercalciuria (eg hypervitaminosis D, hyperparathyroidism and severe renal failure) must not be given calcium supplements.

WARNING AND PRECAUTIONS

Calcium salts should be administered with caution to patients with impaired renal function, cardiac disease, sarcoidosis and history of renal stone formation.

DRUG INTERACTIONS

1. Calcium enhances the effects of digitalis on the heart and may precipitate digitalis intoxication.
2. Calcium salts reduce the absorption of tetracyclines.

ADVERSE EFFECTS

Calcium salts may cause constipation.

OVERDOSE AND TREATMENT

Symptoms of overdosage of calcium may lead to hypercalcaemia usually in patients with renal failure which include anorexia, nausea, vomiting, constipation, abdominal pain, muscle weakness, mental disturbance, polydipsia, polyuria, nephrocalcinosis, renal calculi and in severe cases cardiac arrhythmias and coma.

Calcium intake should be reduced to a minimum and any dehydration and electrolyte imbalance corrected immediately by giving large amount of fluid.

STORAGE CONDITIONS

Store below 30°C.

Protect from light and moisture.

PRESENTATION

PVC/Alu blister of 10 tablets.

Pack Size: 10's, 20's, 30's, 60's, 90's, 100's, 150's, 200's, 300's and 500's.

Package Insert

MANUFACTURER & PRODUCT REGISTRATION HOLDER

Noripharma Sdn Bhd.,
Lot 5030 Jalan Teratai,
5½ Mile Off Jalan Meru,
41050 Klang, Selangor.

DATE OF REVISION

March 2022