

OPTISHINE® VITREO EYE DROPS

Each ml contains:

Active Ingredient(s):

Potassium Iodide 3.0mg
Sodium Iodide 3.0mg

Preservative:

Chlorhexidine Acetate 0.1mg

Pharmacology (Summary of Pharmacodynamic and Pharmacokinetic):

Pharmacodynamic:

Systemic as well as topical application of iodides can activate the metabolism and occasionally counteract cloudiness of the vitreous body.

The effect of iodine on early senile cataract has not yet been established. Occasionally it is possible to slow down the progression of cloudiness and the degradation of visual acuity.

Pharmacokinetic:

a) local administration into the eye:

Potassium iodide diffuses rapidly into the vitreous humour of the isolated rabbit's eye. Within 60 minutes 0.25 µmoles of potassium iodide accumulate in 1 ml of aqueous humour. 6.4% of the iodide present in the treated eye was found in the lens and 3.7% in the vitreous body.

b) systemic administration:

The degree of systemic resorption after local administration into the eye has not been investigated.

Indication(s):

Therapeutic uses:

Opacity and bleeding through the vitreous body of any cause (age, myopia, hypertonia, diabetes, periphlebitis), clouding of the lens as a first sign of senile cataract.

Dosage and Administration(s):

Apply 1 drop into the conjunctival sac 1-3 times a day.

Mode of Administration(s):

Ophthalmic.

Contraindication(s):

People allergic to any ingredient in this product should not use it.

This product is contraindicated with functional disturbances of the thyroid or a bland nodular struma.

Warning and Precautions(s):

Carriers of contact lenses should apply the product during the time when the lenses are not worn. Wait at least 30 minutes after the administration of the preparation before placing the lenses back into the eye. In order to minimize systemic resorption the patient should gently compress the lacrimal sac with his finger for 1-2 minutes after the instillation of the eye drops.

Interaction with Other Medicaments(s):

No interactions with systemically applied drugs have been reported. In cases of concomitant ophthalmic medication with other drugs, an interval of at least 5 minutes should be observed between the two medications.

Pregnancy and Lactation(s):

As there is no controlled studies in animals or in pregnant women are available, this product should only be prescribed in those cases where the potential benefit exceeds the risk for the fetus. The use of this product during lactation has not been investigated (Pregnancy category C).

Side Effects(s):

- 1) Systemic effects: This product contains iodides which can cause hyperthyroidism and acne-type effects in predisposed individuals.
- 2) Local effects on the eye: This product may cause slight burning or pricking sensation for short time after application. Occasionally increased tear flow may happen.

Symptoms and Treatment of Overdose(s):

The risk of adverse effects due to topical overdosage or accidental oral ingestion is very limited due to the relatively low total content of iodides per dropper bottle.

Storage Condition(s):

Store at temperature below 30°C (before and after opening).

Shelf Life(s):

2 years from the date of manufacture.

Do not use for more than 30 days after opening.

Close the bottle immediately after opening.

Product description & Packing(s):

A clear and colorless to slightly yellowish solution.

Plastic bottle of 10ml, 10ml x 10, 10ml x 12, 10ml x 20 & 10ml x 24.



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679