

OPTICLATE EYE DROPS 2.0% W/V

(Sodium Cromoglicate Eye drops, Solution 2.0% w/v)

Name and Strength of active ingredient:

Each ml contains:

Sodium cromoglicate... 20mg

Benzalkonium Chloride (as preservative)... 0.1mg

water for injection ... q.s. to 1ml

Dosage Form : EYE PREPARATIONS in form of EYE DROPS

Product Description

Clear colorless to pale yellow solution.

Pharmacodynamics

Cromoglicate appears to act mainly through the local effect on the eyes. Cromoglicate prevents release of mediator of type I allergic reactions, including histamine and cyteinyl leukotrienes (slow-reacting substance of anaphylaxis [SRS- A]) , from sensitized mast cells after the antigen-antibody union has taken place.

Cromoglicate suppresses IgE mediated release of substance (e.g.,histamine,cyteinyl leukotrienes)from mast cells, inhibits type III (late allergic, Arthus) reactions to a lesser extent, also inhibit secretion induced by other mast cell secretagogues (e.g.,theployamine 48/80).

Cromoglicate appears to inhibit phospho lipase A-induced mast cell degranulation and subsequent release of chemical mediators invitro. It has been suggested that the drug may block calcium channels in the mast cell membranes. It has also been suggested that cromoglicate may inhibit histamine release from mast cells by regulating phosphorylation of a specific mast cell protein involved in secretory mechanisms.

Pharmacokinetics

Sodium cromoglicate is only minimally absorbed from the eye following topical administration. Drug excretion studies in healthy individuals indicate that ~ 0.03% of on ophthalmic dose of cromoglicate as solution is absorbed systemically. Trace amounts (less than 0.01%) of the sodium cromoglicate dose penetrate in to the aqueous humor and clearance from this chamber is virtually complete within 24 hours after treatment is stopped. Sodium cromoglicate does not cross most of the biologic membranes. It is not known if the drug is distributed in human milk. The elimination half-life of sodium cromoglicate is reported to be 81 minutes.

Indication

For prevention and treatment of allergic conjunctivitis including seasonal perennial allergic conjunctivitis and vernal kerato conjunctivitis.

Recommended Dose

Instill one or two drops in each eye 4 times a day.

The number of daily applications and length of treatment can be modified in accordance with physician's criteria.

Route of Administration : OPTHALMIC

Contraindication

Sodium cromoglicate is contraindicated in patients with known hypersensitivity to cromoglicate or to any of the ingredients

Warning and Precautions

This formulation of Sodium Cromoglicate Eye Drops contains benzalkonium chloride as a preservative. Benzalkonium chloride may be deposited in soft contact lenses. Hence, Sodium Cromoglicate Eye Drops should not be used while wearing these lenses. The lenses should be removed before instillation of the drops and not reinserted earlier than 15 minutes after use.

Patients should also be instructed that ocular solutions, if handled improperly can become contaminated by common bacteria known to cause ocular infections. Serious damage to the eye and subsequent loss of vision may result from using contaminated solutions. Patients should also be advised that if they develop any intercurrent ocular condition (e.g. trauma, ocular surgery or infection), they should immediately seek their physician's advice concerning the continued use of present multi-dose container. There have been reports of bacterial keratitis associated with the use of topical ophthalmic products.

Interactions With Other Medicaments

None Known

Pregnancy and Lactation

There are no adequate and well-controlled studies of Sodium Cromoglicate eye drops in pregnant women. Therefore, use during pregnancy is not recommended unless the benefit outweighs the potential risk.

It is not known whether Sodium Cromoglicate is excreted in human milk. Therefore caution should be exercised when the eye drops are administered to nursing mothers.

Side Effects

Transient stinging and burning on instillation of the drops. Rarely, other symptoms of local irritation.

Symptoms and Treatment of Overdose

Medical observation is recommended in cases of overdosage

Sodium cromoglicate is poorly absorbed both from the eye and from the gastro intestinal tract

Storage Condition

Unopened & after opening: Store below 30°C. Protect from light.

Throw the bottle away 1month after opening, even if there is solution remaining.

Shelf Life

36 months from the date of manufacture

After Opening: Throw the bottle away 1 month after opening, even if there is solution remaining.

PACKAGING:

For 5ml pack: 5ml of solution in 5ml pack of LDPE container formed by BFS (Blow fill seal) is packed in a carton along with pack insert.

For 10ml pack: 10ml ml of solution in 10ml pack of LDPE container formed by BFS (Blow fill seal) is packed in a carton along with pack insert.

Manufactured by:

MICRO LABS LIMITED

No.113 to 116, 4th Phase, K.I.A.D.B.,

Bommasandra Industrial Area

BANGALORE – 560 099. INDIA.

Date of information: July 2016