



Front Label - 195x120mm



PRODUCT INFO

BLUE #:	502167APK	Product Name:	Baytril 10% w/v Oral Solution
Item Code:	90208654	Component:	Label
Product Code:	LBAH	Pack Size:	1000 ml
Previous Item Code	90204327		

ARTWORK INFO

Template:	YL_221x120mm_5PAGE_v2a	Packaging Spec(s):	TPM_016649		
Barcodes/Type:	EAN 13: 4007221054603 Laetus: 0021	Add. Info:	N/A		
		Minimum Core Data Point Size:	9pt		
GTIN:	N/A	Proof #	P1a	By/Date	MJ 10-MAY-2022
					
NON PRINTING	BLACK	PMS Process Blue C	PMS 368 C	PMS 285 C	

ELANCO ARTWORK LEGEND v16

Inside Front Label - 195x120mm

9mm

186mm

120mm

Pharmacodynamics

Mechanism of action: Two enzymes with an essential role in DNA replication and transcription, DNA gyrase and topoisomerase IV, have been identified as molecular target structures for fluoroquinolones. Both enzymes control the topological state of DNA by cleaving the DNA strands and then resealing them. Initially, both of the strands of the DNA are cleaved. A distant segment of the DNA is then passed through the break before the strands are resealed.

Target structure inhibition is brought about by noncovalent binding of a fluoroquinolone molecule in an intermediate state of the sequence reactions, in which the DNA is still cleaved and both strands are bound covalently to the enzymes. Replication forks and translational complexes are not able to proceed beyond such enzyme-DNA-fluoroquinolone complexes. The resulting inhibition of DNA and mRNA synthesis results in the rapid and concentration-dependent death of the pathogenic bacteria.

Antibacterial spectrum: Enrofloxacin acts against numerous gram-negative and gram-positive bacteria along with *Mycoplasma* spp. *In vitro* susceptibility has been demonstrated in the following strains: (i) Gram-negative species such as *Escherichia coli*, *Pasteurella multocida* and *Avibacterium* (*Haemophilus*) *paragallinarum* and (ii) *Mycoplasma gallisepticum* and *Mycoplasma synoviae* (see section Warning and Precaution).

Types and mechanisms of resistance: There are five causes of fluoroquinolone resistance: (i) Point mutations within the genes that code for DNA gyrase and or topoisomerase IV and so lead to changes in the individual enzymes, (ii) changes in permeability to the active ingredient in Gram-negative bacteria, (iii) efflux mechanisms, (iv) plasmid-mediated resistance and (v) gyrase-protecting proteins. All mechanisms lead to reduced bacteria susceptibility to fluoroquinolones. Cross-resistance within the fluoroquinolone class of antimicrobial active ingredients is common.

Pharmacokinetics

Enrofloxacin is absorbed rapidly and very well after administration to poultry via the drinking water, with a bioavailability of approx. 90%. The peak plasma concentration of 2 mg/l is reached within 1.5 hours of administration of a single bolus at a dosage of 10 mg/kg body weight, with a total systemic availability of 14.4 mg hr/l. The total body clearance of enrofloxacin is 10.3 ml/min*.kg. If the product is administered continuously via the drinking water, steady state concentrations of 0.5 mg (turkeys) and 0.8 mg (chickens) enrofloxacin per liter are achieved. A high average volume of distribution (5 l/kg) indicates good tissue penetration of enrofloxacin. The concentrations in target tissues such as the lungs, liver, kidneys, bowel and muscles significantly exceed plasma concentrations. In poultry, enrofloxacin is only poorly metabolized to its active metabolite, ciprofloxacin (approx. 5%). Enrofloxacin has an elimination half-life of 6 hours. Protein binding is approx. 25% in poultry.

182mm

Indications

Treatment of infections caused by the following enrofloxacin-susceptible bacteria:

Chickens: *Mycoplasma gallisepticum*, *Mycoplasma synoviae*, *Avibacterium paragallinarum*, *Pasteurella multocida*, *Escherichia coli*.

Turkeys: *Mycoplasma gallisepticum*, *Mycoplasma synoviae*, *Pasteurella multocida*, *Escherichia coli*.

Rabbits: For the treatment of infectious diseases caused by *Pasteurella multocida* and for the treatment of bacterial enteritis caused by *E. coli*.

Enrofloxacin should be used if it is indicated as the agent of choice on the basis of the available clinical experience, which should be supported where possible by susceptibility testing of the causative pathogen.

Recommended Dose

Chickens and turkeys

10 mg/enrofloxacin/kg body weight per day on 3-5 consecutive days.

Treatment for 3-5 consecutive days; for 5 consecutive days in the case of mixed infections and chronic disease forms. In the absence of any clinical improvement within 2-3 days, a change in the antibiotic therapy should be considered on the basis of susceptibility testing.

For use in the drinking water. Always make sure that the entire dose offered has been consumed. The medicated drinking water should be made up fresh each day just before being offered to the birds. The drinking water must be medicated throughout the treatment period, and no other source of water should be available. In order to avoid underdosing, the body weight of the birds should be determined as accurately as possible before treatment.

Only use presolutions made up fresh every day just before the start of treatment. Dispensers should be monitored continuously to ensure appropriate incorporation. Before treatment, the water dispenser system must be emptied and filled with the medicated water.

The daily dose (ml) of Baytril 10% oral solution must be calculated as follows for the period of treatment:

Number of birds x average body weight in kg x 0.1 = total volume (ml) of Baytril 10% oral solution in the drinking water per day.

Baytril 10% oral solution can be placed directly in the container or introduced via a water dispenser/metering pump system.

Rabbits

10 mg/kg body weight per day on 5 consecutive days.

The daily dose (ml) of Baytril 10% oral solution must be calculated as follows for the period of treatment:

Number of rabbits x average body weight in kg x 0.1 = total volume (ml) of Baytril 10% oral solution in the drinking water per day.

Administration Route

For use in drinking water.

120mm

182mm

120mm

Contraindication
Do not use for prophylaxis. Do not use in the presence of known resistance/ cross-resistance to (fluoro-)quinolones in the flocks/ populations to be treated. Do not use in case of hypersensitivity to the active substance, to other (fluoro-) quinolones or to any of the excipients.

Warning and Precaution
Special warnings for each target species
In some cases, the treatment of infections involving *Mycoplasma* spp. does not lead to elimination of the pathogen.
Special precautions for use in animals
The official and local guidelines on antibiotics must be observed when this veterinary medicinal product is used.

Fluoroquinolones should be reserved for treating clinical conditions which have failed to respond or are not expected to respond adequately to other classes of antibiotic. Since enrofloxacin was first authorized for use in poultry, there has been a widespread

reduction in the susceptibility of *E. coli* to fluoroquinolones and emergence of resistant organisms. The emergence of resistance in *Mycoplasma synoviae* has also been reported in the EU.

Whenever possible, fluoroquinolones should be used only on the basis of susceptibility testing.

Failure to use the product in accordance with the instructions in the Package Leaflet may promote the spread of fluoroquinolone-resistant bacteria and reduce the efficacy of treatment with other quinolones because of possible cross-resistance.

Special precautions to be taken by the person administering the veterinary medicinal product to animals
People with known hypersensitivity to fluoroquinolones or to any of the other ingredients should avoid contact with the veterinary medicinal product. Avoid contact with skin and eyes. In case of contact with the skin or eyes, rinse immediately with water. After use, wash hands and skin that have come into contact with the product. Do not eat, drink or smoke during use.

TO BE PRESCRIBED BY REGISTERED VETERINARY SURGEONS ONLY.

Interaction with Other Medicinal Product
An antagonistic effect has been demonstrated in vitro when fluoroquinolones are combined with bacteriostatic active substances such as macrolides, tetracyclines or phenicols. Concomitant administration of magnesium- or aluminium-containing substances can impair the absorption of enrofloxacin.

Use during pregnancy, lactation or lay
Not authorized for use in laying birds producing eggs for human consumption. Do not use in pullets within 14 days of coming into lay.

Adverse reactions
None known.

Baytril is sold by Elanco or its affiliates and is not a product of Bayer. The Product Name Baytril is owned by Bayer and used under license.
© 2022 Elanco or its affiliates

Base Label - 221x120mm

221mm

26mm

181.5mm

Elanco

Baytril™

10% w/v

Oral Solution

Symptoms and Treatment of Overdose

No symptoms of clinical intolerance were observed in turkeys and chickens following treatment with 6 and 10 times the recommended dose respectively. During the growth phase, the use of fluoroquinolones combined with a prolonged period of significantly increased drinking water intake and therefore increased active substance intake, e.g. as a result of high ambient temperatures, can potentially lead to damage to the joint cartilage.

Withdrawal period

Edible tissue:

Chickens: 7 days

Turkeys: 13 days

Rabbits: 15 days

Not authorised for use in poultry producing eggs for human consumption. Do not use in pullets within 14 days of coming into lay.

Storage condition

Do not store above 30°C.

After first opening of the bottle, the product must be used up within 24 hours.

Shelf life after preparation of the ready-to-use solution: 24 hours

Container should be disposed in accordance with local requirements.

This product does not contain ingredients of animal origin.
Keep out of reach of children.
Jauhkan daripada kanak-kanak.

CONTROLLED MEDICINE

MAL16105039HA

Marketing Authorisation Holder: **Elanco Malaysia Sdn Bhd**
Unit 5.04, Level 5 & 6, Tower Block, The Bousteador,
No. 10, Jalan PJU 7/6, Mutiara Damansara,
47800 Petaling Jaya, Selangor, Malaysia

Manufactured by:

KVP Pharma + Veterinär Produkte GmbH
Projensdorfer Straße 324, 24106 Kiel, Germany.

Date of revision: April 2022

Elanco

90208654

Batch No:
Manufacturing Date:
Expiry Date:

PharmacoCode
Read Direction

7mm

6mm

* Please note: Item code and pharma code position must match the front page.

- GTIN code placement is flexible but where possible should be placed close to the variable data info. Remove placeholder if not required.
- Ensure VD Headers remain RIGHT aligned and that the longest Header Text starts 2mm from the edge

For 3 VD Headers - Use Lines (2, 3 & 4)
For 2 VD Headers - Use Lines (3 & 4)
For 1 VD Header - Use Line (4)

120mm
Note: A minimum of 20mm is required for Print Free VD area. This can be extended as required. VD Headers can be any length.

20mm min.
2mm