

SDQL Salinomycin 12% (w/w) Granule Premix

Active substance & strength: Salinomycin sodium 12% (120g/kg)

Product Description: Brown to yellowish- brown granule

Pharmacodynamics: The anticoccidial effect of Salinomycin is aimed mainly against the free merozoites and schizonts of first and second generation. Like all other Polyene antibiotics, acts as an Ionophore Salinomycin and conveys the influx of cations into the cell with this property. By Salinomycin sodium and potassium ions are transported in Exchange for protons in the cell. This leads to a breakdown of the natural transmembrane ion gradient to a pH shift and the entry of water into the cell. The resulting increase in pressure causes a destruction of intracellular structures and thus cell death. Salinomycin significantly reduced the lesions caused by coccidia.

Pharmacokinetics: The active ingredient is absorbed quickly. Salinomycin is quickly distributed and eliminated. The kinetics of serum concentration is most likely a two compartment model with a distribution phase which is finished after one hour. Since the active ingredient is lipophilic and can easily diffuse across cell membranes, the distribution of Salinomycin in tissue is greater than in the serum. The highest concentrations are in liver, followed by heart, kidneys, muscles, fat and skin found. In the tissue, the active ingredient can no longer be detected, except in the liver, where tracks are measured up to 72 h after application. The active ingredient is eliminated quickly. Salinomycin showed a high rate of Body clearance.

Indication: For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati* in chicken broilers

Recommended Dosage: Chickens (broilers) Salinomycin sodium 40 to 60 grams per ton of feed (which equivalent to 333 grams to 500 grams of product per ton of feed)

Mode of Administration: orally into feed

Contraindications: Feed continuously as sole ration. Do not feed to layers. Not approved for use with pellet binders. May be fatal if accidentally fed to adult turkeys or horses.

Warnings and Precautions: Do not feed to laying chickens.

(1) Do not allow dogs, horses or turkeys access to this drug as it is known to be toxic to these species. Extra care should be taken to avoid contamination of feeds for these animals. (2) A weight gain reduction may occur in cockerels fed this drug in the absence of coccidial exposure. (3) Do not administer tiamulin to birds receiving this drug. (4) Do not use in feeds containing pellet binding agents. (5) Do not use this drug for the treatment of coccidiosis. (6) Consult a veterinarian or poultry pathologist if losses exceed 0.5% in a 2-day period. (7) Do not feed to replacement or breeding chickens.

Interactions with Other Medicaments: Basically the Salinomycin combination leads to interactions and significant growth depression in chickens and on the other hand to enhanced anti coccidial activity against *Eimeria tenella* and *E. mitis*. The effect of growth depression was eliminated by a reduced Salinomycinosis (12% reduction) and the anticoccidial activity remained intact. At the same time 120ppm should be administered.

Erythromycin, Sulfachlorpyrazin, Sulfadoxine: Together with Salinomycin could especially a reduction of the feed and water intake and are therefore detected a significant growth depression.

Sodium bentonite: Administered as a feed additive in chickens to bind mycotoxins and to reduce the severity of aflatoxin - and Mycotoxicosis. An investigation shows that sodium bentonite can affect the effect of Salinomycin, if this ingredient is used in the half recommended dosage. But the recommended Salinomycin is sufficient, so that in practice no problems should occur.

Roxarsone: The effects of Salinomycin and Roxarsone are additive. There be fewer lesions caused by coccidia, improved body weight gain and feed conversion is easier.

Tiamulin: There is a pronounced toxic interaction between Salinomycin and Tiamulin. Tiamulin interferes with the metabolism of Salinomycin in chicken and leads to weight loss.

Pregnancy and Lactation: Do not use in laying birds producing eggs for human consumption

Adverse Effects/Undesirable Effects:

-Egg production and quality-Salinomycin and other Ionophore affecting the egg and the ability for chicks to hatch. They are also fetotoxic.

-Heart and circulation-Salinomycin increases the cardiac output, blood pressure, the force of contraction of the left ventricle in dogs, improves the myocardial blood perfusion and the mechanical efficiency of the heart.

-Rumen flora-Salinomycin shows no effect on the cellulolytischen and lactatverwertenden bacteria, but increases the share of the amylolytischen bacteria. The total number of protozoa decreases initial.

-Immune system-Administered with 60 mg/kg Salinomycin during 49 days to chickens, had no notable effect on the humoral and cellular immune response to vaccine of Newcastle disease. A drastic immune suppressive effect but occurred after 120 mg/kg food

Overdose and Treatment: Poultry should have access to drinking water at all times. Drug overdosage or lack of water intake may result in leg weakness or paralysis.

Withdrawal Period(s): Zero day **Packaging available:** 25kg bag

Storage condition: Store around 30 degree Celsius (room temperature)

Name and address of manufacturer:

Shandong Qilu King-Phar Pharmaceutical Co. Ltd.

No 21 Qinglong Road, Pingyin, Jinan, China

Name and address of marketing authorisation holder:

QL Pacific Vet Group Sdn Bhd

No 886-C8, Jalan Subang 9, Taman Perindustrian Subang, 47600 Subang Jaya Selangor, Malaysia

Date of Revision of Package Insert : 11 October 2016