

FMP Insert (VN)
Draft (1) 05 Aug 19

FEMENTIC

PREMIX POWDER 8% w/w

Ingredient(s):

Each g contains:

Fenbendazole 80mg

Pharmacodynamics:

Fenbendazole is a methylcarbamate benzimidazole antiparasitic agent and has a broad spectrum of activity against a variety of pathogenic internal parasites. In susceptible parasites, benzimidazole mechanism of action is believed due to disrupting intracellular microtubular transport system by binding selectively and damaging tubulin, preventing tubulin polymerization, and inhibiting microtubule formation. Benzimidazoles also act at higher concentrations to disrupt metabolic pathways within the helminth, and inhibit metabolic enzymes, including malate dehydrogenase and fumarate reductase. Benzimidazoles may be considered time-dependent antiparasitic agents.

Pharmacokinetics:

Fenbendazole is only marginally absorbed after oral administration. The amount absorbed from the gut is apparently more associated with the solubility of the drug and not the dose given. After oral dosing in calves, peak blood levels of 0.11 micrograms/mL, was measured. Absorbed fenbendazole is metabolized (and vice-versa) to the active compound, oxfendazole (sulfoxide) and the sulfone. In cattle and swine, 44-50% of a dose of fenbendazole is excreted unchanged in the faeces, and <1% in the urine.

Indication:

In cattle, treatment of infections with the following parasites:

Gastrointestinal nematodes (adult and L4 larvae): *Ostertagia ostertagi* (including inhibited larvae) *Trichostrongylus* spp., *Cooperia onchophora*, *Nematodirus helvetianus*, *Bunostomum phlebotomum*, *Strongyloides papillosus*, *Oesophagostomum radiatum*;
Lungworms (adults and L4 larvae): *Dictyocaulus viviparus*.

In swines, treatment of infections with the following parasites:

Gastrointestinal nematodes: *Ascaris suum*, *Strongyloides ransomi*, *Oesophagostomum* spp.; Lungworms: *Metastrongylus* spp.

Target Species:

Cattle, swine

Recommended Dose:

Cattle:

7.5 mg fenbendazole per kg body weight, single dose orally, 9.375g of product per 100 kg body weight to be mixed with the food.

Swine:

5 mg fenbendazole per kg body weight, single dose orally, 6.25g of product per 100 kg body weight to be mixed with the food.

Mode of Administration:

Oral administration via feeds.

Contraindication:

Do not use in animals with known hypersensitivity to the active substance.

Do not use if you suspect resistance to benzimidazole.

Warning and Precautions:

Body weight should be assessed as accurately as possible before calculating the dosage.

Resistance to all benzimidazoles can develop during frequent and repeated use of a pest control of this family.

Interaction with Other Medicaments:

Oxbendazole or fenbendazole should not be given concurrently with the bromsalan flukicides; abortions in cattle have been reported after using these compounds.

Pregnancy and Lactation:

Fenbendazole is generally considered safe to use in pregnancy for all species.

Side Effects:

At usual doses, fenbendazole generally does not cause any adverse effects. Hypersensitivity reactions secondary to antigen release by dying parasites may occur, particularly at high dosages.

Environmental Property:

Fenbendazole is water insoluble; exhibits rapid degradation under photolysis (less than one day). Fenbendazole adsorbed tightly to particles of soil and is not expected to migrate from application sites into runoff or leachate water.

Symptoms and Treatment of Overdose:

Fenbendazole is apparently well tolerated at doses up to 100X recommended. The LD50 in laboratory animals exceeds 10g/kg when administered PO. It is unlikely an acute overdosage would lead to clinical signs.

Storage Condition:

Store in a cool, dry place, below temperature 30°C.

Shelf Life:

3 years.

For food producing animals product**Withdrawal Periods:**

Animals must not be slaughtered for human consumption during treatment.

Cattle: Meat – 11 days

Milk – 6 days

Swine: Meat – 8 days

Maximum Residual Limit (MRL):

Cattle:

Milk, muscle, kidney, fat: 100µg/kg

Liver: 500µg/kg

Swine:

Muscle, kidney, fat: 100µg/kg

Liver: 500µg/kg

Product Description:

A white to off white color powder.

10kg in paper bag



Made by YSP

Manufacturer:

Y.S.P INDUSTRIES VIETNAM CO., LTD

Road No.6, Nhon Trach 3 Industrial zone, Nhon Trach,
Dong Nai province, Vietnam Tel: +8461 3569 993

Product Registration Holder:

Y.S.P. INDUSTRIES (M) SDN. BHD. (192593 U)

Lot 3, 5 & 7, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor Darul Ehsan, Malaysia.

Ordering Line: 1 800 88 3027

Product Info: 1 800 88 3679

xxxxxx printing code xxxxxx

Date of revision: 05 Aug 2019