

Defuzin™

Fusidic Acid Cream 2% w/w
Sodium Fusidate Ointment 2% w/w

COMPOSITION

Defuzin Cream

Contains Fusidic Acid 2%w/w.

Methyl Hydroxybenzoate 0.18%w/w and Propyl Hydroxybenzoate 0.02%w/w as preservatives.

Defuzin Ointment

Contains Sodium Fusidate 2% w/w

PHARMACODYNAMICS

Fusidic acid/Sodium Fusidate is a steroidal antibacterial with a bacteriostatic or bactericidal activity, mainly against gram-positive bacteria. It restrains bacterial protein synthesis by inhibiting a factor necessary for translocation of peptide subunits and elongation of the peptide chain.

Fusidic acid/Sodium Fusidate is very active against gram-positive bacteria. Staphylococci, in particular *Staphylococcus aureus* and *Staphylococcus epidermis* (including methicillin-resistant strains), *Nocardia asteroides* and many clostridial strains. The streptococci and enterococci are less susceptible.

Fusidic acid/Sodium Fusidate has some activity against gram-negative bacteria, namely *Neisseria spp.*, *Bacteroides fragilis*, *Mycobacterium tuberculosis* and *M. leprae*; a range of protozoa including *Giardia lamblia* and *Plasmodium falciparum*.

PHARMACOKINETICS

Following topical application, the degree of Fusidic acid/Sodium Fusidate penetration on intact human skin depends on factors such as the duration of exposure to Fusidic acid/Sodium Fusidate and the condition of the skin.

Fusidic acid/Sodium Fusidate is excreted mainly in the bile, almost entirely as metabolites, some of which have weak antimicrobial activity. Little is excreted in the urine.

INDICATIONS

Defuzin is indicated for treatment of skin infections caused by staphylococci, streptococci, *Propionibacterium acnes*, *Corynebacterium minutissimum*, and other organisms sensitive to Fusidic acid/Sodium Fusidate. The most important indications are impetigo, infected wounds, folliculitis, boils, sycosis barbae, carbuncles, hidradenitis, paronychia and erythrasma.

DOSAGE AND ADMINISTRATION

For external use only. Apply to the affected area 2-3 times daily, generally for a period of 7 days, except in treatment of acne where therapy should be extended for a longer period according to the nature of this disease.

CONTRAINDICATIONS

Infection caused by non-susceptible organisms. Known hypersensitivity to Fusidic acid/Sodium Fusidate or to any of the excipients.

WARNINGS AND PRECAUTIONS

As with all antibiotics, extended or recurrent use may increase the risk of developing antibiotic resistance.

Extended or recurrent use may increase the risk of developing contact sensitisation.

Caution should still be exercised when Fusidic acid/Sodium Fusidate is used near the eyes.

DRUG INTERACTIONS

No drug interactions have been identified.

PREGNANCY AND LACTATION

Use in pregnancy: There is inadequate evidence of safety in human pregnancy. Animal studies and many years of clinical experience have suggested that Fusidic acid/Sodium Fusidate is devoid of teratogenic effect. The topical use of Fusidic acid/Sodium Fusidate requires that the potential benefits be weighed against the possible hazards to the foetus.

Use in lactation: Safety in nursing mothers has not been established. When Fusidic acid/Sodium Fusidate has been given systemically, levels have been detected in breast milk, but with topical use the possible amount of drug present is unlikely to affect the infant.

SIDE EFFECTS

Fusidic acid/Sodium Fusidate is well tolerated.

Hypersensitivity reactions in the form of rashes and irritation may occur with topical treatment.

SYMPTOMS AND TREATMENT FOR OVERDOSAGE

Overdose is unlikely to occur. Unless hypersensitivity to Fusidic acid/Sodium Fusidate or any of the excipients exists, accidental ingestion of Fusidic acid/Sodium Fusidate is unlikely to cause any harm.

STORAGE CONDITION

Store below 30°C.

SHELF LIFE

The expiry date is indicated on the packaging.

PRODUCT DESCRIPTION, DOSAGE FORMS AND PACKAGING AVAILABLE

Defuzin Cream

White, smooth, homogeneous cream in aluminium collapsible tube of 5g and 15g.

Defuzin Ointment

Off white to yellow, smooth, homogeneous fatty basis ointment in aluminium collapsible tube of 5g and 15g.

KEEP OUT OF REACH OF CHILDREN / JAUHI DARI KANAK-KANAK

For further information, please consult your pharmacist or physician.

Revision Date : 5-May-2016

Manufactured and Marketed by:

Xepa-Soul Pattinson (Malaysia) Sdn Bhd

1-5, Cheng Industrial Estate, 75250 Melaka, Malaysia.